ACETAMINOPHEN- acetaminophen tablet Contract Pharmacy Services-PA

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Acetaminophen Tablets

DRUG FACTS

ACTIVE INGREDIENT (IN EACH TABLET)

Acetaminophen Tablets, 325mg

PURPOSE

Pain reliever/fever reducer

USES

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - 1. headache
 - 2. muscular aches
 - 3. the common cold
 - 4. minor pain of arthritis
 - 5. backache
 - 6. toothache
 - 7. premenstrual and menstrual cramps

WARNINGS

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you

have liver disease

Ask a doctor or pharmacist before use if

you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DIRECTIONS

adults and children 12 years and over

- take 2 tablets every 4 to 6 hours while symptoms last
- children do not take more than 8 tablets in 24 hours
 - do not take for more than 10 days unless directed by a doctor

do not use adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

OTHER INFORMATION

store at 15° to 30°C (59° to 86°F)

INACTIVE INGREDIENTS

povidone, pregelatinized starch, stearic acid

QUESTIONS OR COMMENTS?

Distributed by: Marlex Pharmaceuticals, 50 McCullough Drive, New Castle, DE 19720, U.S.A.


call toll free **1-866-820-7381**

Repackaged by:

Contract Pharmacy Services-PA 125 Titus Ave Suite 200 Warrington, PA 18976 USA

Original--08/2010--NJW

Package Label



Blister of 30 Tablets

ACETAMINOPHEN

acetaminophen tablet

Product Information	t Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67046-025(NDC:10135-123)	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg	

		Product Characteristics				
Color	WHITE	Score	no score			
Shape	ROUND	Size	10 mm			
Flavor		Imprint Code				
Contains						

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:67046-025-30	30 in 1 BLISTER PACK			
2	NDC:67046-025-60	60 in 1 BLISTER PACK			
3	NDC:67046-025-62	60 in 1 BLISTER PACK			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	08/02/2010		

Labeler - Contract Pharmacy Services-PA (945429777)

Revised: 7/2010 Contract Pharmacy Services-PA