

TOKISAN SORE THROAT RELIEF- benzocaine liquid
GOYESCAS OF FLA

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dose

Directions

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

Warnings

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Inactive Ingredients

Inactive ingredients

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol

Uses

Uses for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

Purpose

DRUG FACTS PURPOSE

Benzocaine 5% Anesthetic

Warnings

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Active Ingredient

DRUG FACTS

ACTIVE INGREDIENTS PURPOSE

Benzocaine 5% Anesthetic

DRUG FACTS

ACTIVE INGREDIENTS PURPOSE

Benzocaine 5% Anesthetic

Uses for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Directions

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

Other Information

Store at room temperature (68-77F) (20-25C)

Do not freeze

Alcohol Free

Inactive ingredients

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol.

TOKISAN[®]

TOKISAN[®]

Drug Facts

| Active Ingredients | Purpose |
|--------------------|------------|
| Benzocaine 5% | Anesthetic |

Uses For the temporary relief of:

- occasional minor irritation, pain, sore mouth and sore throat
- pain due to canker sores ■ pain due to minor irritation or injury of the mouth and gums ■ minor discomfort and protection of irritated areas in sore mouth and sore throat

Warnings

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not exceed recommended dosage.

Stop use and ask a doctor if:

- sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting ■ sore mouth symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens

If pregnant or breast feeding, ask a health professional before use.

Directions

Adults and children 2 years of age and older apply to affected area for at least one minute and then spit out.

Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the use of the product. Children under 2 years of age. Consult a doctor.

Other Information

- store at room temperature (69-77°F) (20-25°C)
- do not freeze
- alcohol free

Inactive Ingredients

Benzalkonium Chloride, Eucalyptus Oil, FD & C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and Sorbitol.

ORIGINAL THROAT SPRAY

For the Temporary Relief of
• MINOR SORE THROAT
and other minor irritations.



DISTRIBUTED BY:
GOYESCAS CORP. OF FLA.
MIAMI, FLORIDA 33178
GOYESCASUSA.COM
1 FL. OZ.



Drug Facts

Active Ingredients

Uses for the temporary relief of: ■ occasional minor sore throat ■ pain due to minor irritation or injury of the throat ■ relief of minor pain, discomfort and irritation.

Directions: Adults and children 2 years of age and older: spray 1 to 2 sprays into the mouth 4 to 6 times a day. Children under 2 years of age: Consult a doctor.

Warnings: ■ do not freeze ■ alcohol free

TOKISAN[®]
ORIGINAL SPRAY

For the temporary relief of MINOR SORE THROAT, MOUTH SORES, CANCER OF THE THROAT and other minor oral irritation.

Ingredients:
GENTIANIC ACID, P-FLUORINATED BENZOIC ACID, HYDROXYETHYLCELLULOSE, POLYETHYLENE GLYCOL, PURIFIED WATER, SODIUM BICARBONATE, SODIUM CITRATE AND SODIUM CHLORIDE.

Warnings

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. See the back of the package if you have history of allergy to "red" antibiotics.

Do not exceed recommended dosage.

Stop use and ask a doctor if:

- sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, skin peeling, or swollen lymph nodes in neck.
- symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens.

If pregnant or breast feeding, ask a health professional for advice before using.

Inactive Ingredients:
Benzalkonium chloride, octopylates oil, PEG-400, propylene glycol, purified water, sodium bicarbonate, sodium lauryl sulfate and sorbitol.

TOKISAN SORE THROAT RELIEF

benzocaine liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11856-5001 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------|--------------------------|-----------------|
| BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) | BENZOCAINE | 50 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|------------------------------------------|-----------------|
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) | |
| EUCALYPTUS OIL (UNII: 2R04ONI662) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM BORATE (UNII: 91MBZ8H3QO) | |
| SORBITOL (UNII: 506T60A25R) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|-------------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:11856-5001-1 | 1 in 1 BOX | | |
| 1 | | 33 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|-------------------------------------------------|-----------------------------|---------------------------|
| OTC monograph final | part346 | 08/01/1958 | |

Labeler - GOYESCAS OF FLA (032506248)