

GOYESCAS BRONCOVAC 5 SORE THROAT RELIEF- benzocaine liquid
GOYESCAS OF FLA

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dosage & Administration

Directions

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

Warnings

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Inactive Ingredients

Inactive ingredients

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol

Uses

Uses for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

Purpose

DRUG FACTS PURPOSE

Benzocaine 5% Anesthetic

Warnings

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Active Ingredients

DRUG FACTS

ACTIVE INGREDIENTS PURPOSE

Benzocaine 5% Anesthetic

DRUG FACTS

ACTIVE INGREDIENTS PURPOSE

Benzocaine 5% Anesthetic

Uses for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

Warnings

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

Directions

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

Other Information


Store at room temperature (68-77F) (20-25C)

Do not freeze

Alcohol Free

Inactive ingredients

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol.



Goyescas
#5
BRONCOVAC®

Goyescas
#5
BRONCOVAC®

Drug Facts

Active Ingredients	Purpose
Benzocaine 5%	Anesthetic

Uses: for the temporary relief of:
■ occasional minor irritation, pain, sore mouth and sore throat
■ pain due to canker sores ■ pain due to minor irritation or injury of the oral and nasal cavities ■ minor discomfort and protection of irritated areas in sore mouth and sore throat

Warnings
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Do not exceed recommended dosage.

Stop use and ask a doctor if:
■ sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting ■ sore mouth symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens.
If pregnant or breast feeding, ask a health professional before use.

Directions
Adults and children 2 years of age and older: apply to affected area for at least one minute and spit out.
Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the use of the product. Children under 2 years of age: Consult a doctor.

Other Information
■ Store at room temperature (68-77°F) (20-25°C)
■ do not freeze
■ alcohol free

Inactive Ingredients
Benzalkonium Chloride, Eucalyptus Oil, FD & C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and Sorbitol.

ORIGINAL THROAT SPRAY

For the Temporary Relief of
• **MINOR SORE THROAT**
and other minor irritations.



DISTRIBUTED BY:
GOYESCAS CORP. OF FLA.
MIAMI, FLORIDA 33178
GOYESCASUSA.COM
1 FL. OZ.



Drug Facts

Active Ingredients:
 Benzocaine 5%
 Diphenhydramine Hydrochloride 1%
 Thymol 0.1%

Use for the temporary relief of:

- sore throat
- minor oral irritation
- canker sores
- pain due to minor irritation or injury of the lips and gums
- minor oral itching and irritation
- itching and sore throat

Directions: Adults and children 2 years of age and older: apply to affected area for at least one minute and then spit out. Do not swallow. Children under 2 years of age: Consult a doctor.

Warnings:

- do not freeze
- alcohol free

Other information:

- Contains 100% alcohol-free ingredients.
- Contains 100% alcohol-free ingredients.

Keep out of reach of children. In case of excessive use, stop using immediately. If symptoms do not improve, get away. Do not use this product if you have history of allergy to "local" anesthetics.

Do not exceed recommended dosage.

Stop use and ask a doctor if:

- symptoms do not improve after 3 days
- symptoms worsen or are followed by fever, hives, rash, swelling, nausea or vomiting
- some mouth symptoms do not improve in 7 days, or if irritation, pain or redness persists
- if pregnant or breast feeding, ask a health professional before use.

Inactive Ingredients:
 Benzalkonium chloride, cetylalcohol, FD&C red #40, hydroxyethylcellulose, hydroxypropyl methylcellulose, potassium sorbate, sodium benzoate and sodium butyrate and sodium

BRONCOVAC®
Soyescszs

ORIGINAL THROAT SPRAY

For the temporary relief of MINOR SORE THROAT, CANKER SORES and other minor oral irritation.

1 fl. oz. (28 mL)

DISTRIBUTED BY: SOYESCZS CORP. OF FLA.,
 MIAMI, FLORIDA 33174
 www.soyesczs.com

GOYESCAS BRONCOVAC 5 SORE THROAT RELIEF

benzocaine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11856-5002
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	
SORBITOL (UNII: 506T60A25R)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
EUCALYPTUS OIL (UNII: 2R04ONI662)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11856-5002-1	1 in 1 BOX		
1		33 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part346	10/01/2006	

Labeler - GOYESCAS OF FLA (032506248)