

**GOYESCAS 7 SYRUPS SORE THROAT RELIEF- benzocaine liquid**  
**GOYESCAS OF FLA**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Dosage & Administration**

**Directions**

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

**Warnings**

**Warnings**

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

**Inactive Ingredients**

**Inactive ingredients**

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol

**Uses**

**Uses** for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

**Purpose**

**DRUG FACTS PURPOSE**

**Benzocaine 5% Anesthetic**

**Warnings**

**Warnings**

**Keep away of reach from children.** In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

## **Active Ingredients**

### **DRUG FACTS**

#### **ACTIVE INGREDIENTS PURPOSE**

**Benzocaine 5% Anesthetic**

### **DRUG FACTS**

#### **ACTIVE INGREDIENTS PURPOSE**

**Benzocaine 5% Anesthetic**

**Uses** for the temporary relief of:

Occasional minor irritation, pain, sore mouth and throat. Pain due to canker sores. Pain due to minor irritation of injury of the mouth and gums, minor discomfort and protection of irritated area in sore mouth and sore throat.

### **Warnings**

Keep away of reach from children. In case of overdose, get medical help or contact a Poison Control Center right away. Do not exceed recommended dosage. Stop use and ask a Doctor if: Sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, sore mouth symptoms do not improve in 7 days or if irritation, pain or redness persist or worsens. If pregnant or breast feeding, ask a health professional before use.

### **Directions**

Adults and children 2 years of age and older, apply to affected area for at least one minute and then spit out. Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the used of the product. Children 2 years of age: Consult a doctor.

### **Other Information**

Store at room temperature (68-77F) (20-25C)

Do not freeze

Alcohol Free

### **Inactive ingredients**

Benzalkonium chloride, Eucalyptus oil, FD&C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and sorbitol.



**Drug Facts**

**Active Ingredients**  
Benzocaine 9%

**Purpose**  
Anesthetic.

**Uses** for the temporary relief of:

- occasional minor irritation, pain, sore mouth and sore throat
- pain due to canker sores
- pain due to minor irritation or injury of the mouth and gums
- minor discomfort and protection of irritated areas in sore mouth and sore throat

**Warnings**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not exceed recommended dosage.

**Stop use and ask a doctor if:**

- sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, hoarseness or vomiting
- sore mouth symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens

If pregnant or breast feeding, ask a health professional before use.

**Directions**

Adults and children 2 years of age and older: apply to affected area for at least one minute and then spit out.

Use up to 4 times daily or as directed by doctor. Children under 12 years of age should be supervised in the use of the product. Children under 2 years of age: consult a doctor.

**Other information**

- stored at room temperature (68-77°F) (20-25°C)
- do not freeze
- alcohol free

**Inactive ingredients**

Benzalkonium Chloride, Eucalyptus Oil, FD & C Red #40, Propylene Glycol, Purified Water, Sodium Bicarbonate, Sodium Borate and Sorbitol.

**ORIGINAL THROAT SPRAY**

For the Temporary Relief of  
• MINOR SORE THROAT  
and other minor irritations.



DISTRIBUTED BY:  
GOYESCAS CORP. OF FLA.  
MIAMI, FLORIDA 33178  
GOYESCASUSA.COM  
**1 FL. OZ.**



**Drug Facts**

**Active Ingredients**  
Benzocaine 5%  
Dry Aromatized Analgesic

**Purpose**  
Temporarily relieves oral pain associated with:  
canker sores ■ pain due to minor irritation or injury of the mouth ■ sore throat ■ minor oral discomfort  
Use only for the relief of minor oral discomfort and pain.

**Directions** Adults and children 2 years of age and older:  
Apply to the area for relief of minor oral discomfort and pain. Use only for the relief of minor oral discomfort and pain. Do not use for more than 7 days. If symptoms persist, consult a doctor.

**Warnings**  
Do not use if you have a history of allergy to "numbing" anesthetics.  
Do not exceed recommended dosage.  
Stop use and ask a doctor if:  
■ sore throat becomes persistent for more than 2 days, is severe, or is accompanied by fever, rash, or swelling, or if you are unable to swallow.  
■ symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens.  
■ you experience dizziness, lightheadedness, or fainting.  
■ you experience numbness or tingling in your mouth or throat.  
■ you experience numbness or tingling in your fingers or toes.  
■ you experience numbness or tingling in your face or neck.  
■ you experience numbness or tingling in your arms or legs.  
■ you experience numbness or tingling in your hands or feet.  
■ you experience numbness or tingling in your feet or toes.  
■ you experience numbness or tingling in your legs or feet.  
■ you experience numbness or tingling in your feet or toes.



**7 JARBES®**

For the temporary relief of MINOR SORE THROAT, CANKER SORES and other minor oral discomforts.

Contains 5% Benzocaine and other active ingredients.

© 2008 Joy Excess, Inc. All rights reserved.

**Warnings**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Tell the doctor if you have a history of allergy to "numbing" anesthetics.  
Do not exceed recommended dosage.  
Stop use and ask a doctor if:  
■ sore throat becomes persistent for more than 2 days, is severe, or is accompanied by fever, rash, or swelling, or if you are unable to swallow.  
■ symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens.  
■ you experience dizziness, lightheadedness, or fainting.  
■ you experience numbness or tingling in your mouth or throat.  
■ you experience numbness or tingling in your fingers or toes.  
■ you experience numbness or tingling in your face or neck.  
■ you experience numbness or tingling in your arms or legs.  
■ you experience numbness or tingling in your feet or toes.  
■ you experience numbness or tingling in your legs or feet.  
■ you experience numbness or tingling in your feet or toes.

**Active Ingredients**  
Benzocaine (ethyl p-aminobenzoate), methylated oil, FD&C red #40, propylene glycol, purified water, sodium lactate, sodium benzoate and paraffin.

# GOYESCAS 7 SYRUPS SORE THROAT RELIEF

benzocaine liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11856-5003
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	50 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
EUCALYPTUS OIL (UNII: 2R04ONI662)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	
SORBITOL (UNII: 506T60A25R)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11856-5003-1	1 in 1 BOX		
1		33 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part346	10/01/2015	

**Labeler** - GOYESCAS OF FLA (032506248)

Revised: 10/2015

GOYESCAS OF FLA