

LAXATIVE- bisacodyl tablet, delayed release
Walgreen Company

Walgreens 44-676A

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

Walgreens

Walgreens Pharmacist Recommended[†]

NDC 0363-6760-01

Laxative Tablets

BISACODYL USP, 5 mg / STIMULANT LAXATIVE

For Sensitive Stomachs

- Gentle, dependable overnight relief

30
ENTERIC -
COATED
TABLETS

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844

ORG011967601

†Our pharmacists recommend the Walgreens brand.

DISTRIBUTED BY: **WALGREEN CO.**

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com

©2021 Walgreen Co.



Walgreens 44-676A

LAXATIVE

NDC 0363-6760-01

Walgreens



Laxative Tablets

BISACODYL USP, 5 mg / STIMULANT LAXATIVE

For Sensitive Stomachs

- Gentle, dependable overnight relief

30
ENTERIC-COATED
TABLETS



ACTUAL SIZE

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient

(in each tablet)

Bisacodyl USP, 5 mg. Stimulant laxative

Purpose

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)

Drug Facts (continued)

Warnings

Do not use if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, flatness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

(Questions or comments? 1-800-426-9391)

Our pharmacists recommend the Walgreens brand.

Inactive ingredients carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcystalline cellulose, polydextrose, polyethylene glycol, stearic acid, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ avoid excessive humidity
- see end flap for expiration date and lot number

Directions ■ take with a glass of water	adults and children 12	take 1 to 3 tablets in a single daily dose
	children 6 to under 12	take 1 tablet in a single daily dose
	children under 6 years	ask a doctor

Drug Facts (continued)

Directions ■ take with a glass of water

adults and children 12 take 1 to 3 tablets in a single daily dose

children 6 to under 12 take 1 tablet in a single daily dose

children under 6 years ask a doctor

Drug Facts (continued)

Warnings

Do not use if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, flatness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcystalline cellulose, polydextrose, polyethylene glycol, stearic acid, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Drug Facts (continued)

(Questions or comments? 1-800-426-9391)

Our pharmacists recommend the Walgreens brand.

how2recycle.info

50844
ORG011967601

W30ORG0821-F

Do not print Area
Lot & Exp. Date



ITEM 504569 W00000-0000-0

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-2201-676A-01-HR
ORG011967601

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2021 Walgreen Co.

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6760
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B710)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6760-01	2 in 1 CARTON	09/16/2015	

1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-6760-52	6 in 1 CARTON	09/16/2015	
2		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	09/16/2015	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-6760)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-6760) , pack(0363-6760)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-6760)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-6760)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-6760)

Revised: 9/2023

Walgreen Company