

BAYER GENUINE ASPIRIN- aspirin tablet
R J General Corporation

Bayer Genuine Aspirin®

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature

Inactive ingredients

carnauba wax ², corn starch, hypromellose, powdered cellulose, triacetin

² may contain this ingredient

Questions or comments?**1-800-331-4536**

(Mon - Fri 9AM - 5PM EST) or www.bayeraspirin.com

PRINCIPAL DISPLAY PANEL - 100 Tablet Pouch Carton***GENUINE ASPIRIN******BAYER***®***ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER*****THE WONDER DRUG**®***Safe Pain Relief Plus
Lifesaving Benefits******325 mg******COATED TABLETS***



This product is repackaged by RJ General Corp, 2350 Burnet Ave., Cincinnati, Ohio 45219 (513)621-3738 (Mon-Fri 9AM-3PM EST) from a product manufactured by Bayer. Bayer®, the Bayer Cross® and The Wonder Drug are registered trademarks of Bayer, used with permission.

Questions? 1-800-331-4536 or www.bayeraspirin.com

Inactive ingredients: carnauba wax, corn starch, hypromellose, powdered cellulose, tracetin. *may contain this ingredient.

Other information: Do not use if packet is torn, cut or opened. Store at controlled room temperature 15° to 30° C (59° to 86° F). Avoid excessive heat and humidity.

Children under 12 years	Consult a doctor.
Adults and children 12 years of age and older.	As directed by a doctor.

Directions: Drink a full glass of water with each dose. Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

Warnings: Children and teenagers should not use this medicine for fever, pain or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. **Reye's syndrome:** Children and teenagers should not use this medicine for fever, pain or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. **Alcohol Warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain reliever/fever reducer. **Do not use:** Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock. **Ask a doctor before using if you have:** asthma, stomach problems (such as heartburn, upset stomach or stomach pain) that last or come back, gastric ulcers, bleeding problems. **Ask a doctor or pharmacist before use if you are taking a prescription drug for:** anticoagulation (thinning of blood), diabetes, gout, arthritis. **When using this product do not exceed recommended dose. Stop use and ask a doctor if:** an allergic reaction occurs, Seek medical help right away, pain gets worse or last more than 10 days, redness or swelling is present, fever last more than 3 days, ringing in the ears or loss of hearing occurs. **If pregnant or breast-feeding baby, ask a health professional before use. It is especially important not to use Aspirin During The Last 3 Months Of Pregnancy Unless Specifically Directed To Do So By A Doctor Because It May Cause Problems In The Unborn Child Or Complications During Delivery. KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts

Active ingredients (in each tablet): Aspirin 325 mg, pain reliever/fever reducer.

Purpose: temporarily relieves minor aches and pains associated with: muscle pain, menstrual pain, headaches, pain and fever of colds, toothache, minor arthritis pain.

Uses: temporarily relieves minor aches and pains associated with: muscle pain, menstrual pain, headaches, pain and fever of colds, toothache, minor arthritis pain.



GENUINE ASPIRIN
BAYER
 ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER



THE WONDER DRUG®



Safe Pain Relief Plus
 Lifesaving Benefits

COATED TABLETS





GENUINE ASPIRIN
BAYER[®] 
ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

THE WONDER DRUG[®]



**Safe Pain Relief Plus
Lifesaving Benefits**

325^{mg}

COATED TABLETS

GENUINE ASPIRIN
BAYER[®] 
ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

THE WONDER DRUG[®]



**Safe Pain Relief Plus
Lifesaving Benefits**

325^{mg}

COATED TABLETS


Bayer HealthCare
 Bayer HealthCare LLC
 Whippany, NJ 07981 USA
 Bayer, the Bayer Cross and The Wonder Drug
 are registered trademarks of Bayer, used with permission.

BAYER GENUINE ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70264-013(NDC:0280-2000)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	BAYER
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70264-013-01	50 in 1 CARTON	07/31/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	07/31/2014	

Labeler - R J General Corporation (122542830)

Establishment

Name	Address	ID/FEI	Business Operations
R J General Corporation		122542830	repack(70264-013)

Revised: 10/2023

R J General Corporation