

BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid bandage liquid
Chain Drug Marketing Association

Quality Choice Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

- do not store at temperature above 120 °F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- ***cap bottle tightly***
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

Quality Choice

For Small Cuts & Wounds

Liquid Bandage

Antiseptic to Help Prevent Infection

Invisible | Flexible | Waterproof | Easy to use

Use Liquid Bandage For:

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks, & help in preventing the formulation of calluses.

Also Used By:

Bowlers, dancers, fishermen, golfers, hikers, musicians, runners, and tennis players

0.3 FL OZ. (9 mL)



*Compare to SKIN SHIELD®

For Small Cuts & Wounds

Liquid Bandage

Antiseptic To Help Prevent Infection

Invisible | Flexible | Waterproof
Easy to use

Use Liquid Bandage For:

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

Also Used By:

Bowlers, dancers, fishermen, golfers, hikers, musicians, Runners, and tennis players.

LB00BT



For Small Cuts & Wounds

Liquid Bandage

Antiseptic To Help Prevent Infection

0.3 FL. OZ. (9mL)

NO VARNISH AREA



©Distributed by C.D.M.A., Inc.
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300

* This product is not manufactured or distributed by Insight Pharmaceuticals LLC, owner of the registered trademark Skin Shield®.

MADE IN USA



DO NOT USE IF NECKBAND IS BROKEN

Drug Facts

Purpose
First aid antiseptic
Topical antiseptic

Active ingredients
Benzethonium chloride 0.75%
Dyclonine hydrochloride 0.75%

Uses
First aid to prevent bacterial contamination in minor cuts, scrapes and burns

Warnings
For external use only.

Flammable: do not use while smoking or near heat or flame.

Do not use
■ in the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor
■ on infected areas or wounds that are draining ■ with other first aid products such as balm and cream ■ over sutures ■ on mucous membranes

Ask a doctor before use if you have
■ deep puncture wounds ■ deep cuts ■ arterial lacerations ■ serious bleeding ■ diabetes ■ poor circulation ■ serious burns

When using this product
■ do not store at temperature above 120°F
■ avoid use by deliberately concentrating and shaking the contents can be harmful or fatal

Stop use and ask a doctor if
■ condition persists or gets worse ■ reaction occurs

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions
■ clean affected area ■ apply a small amount on the area 1-5 times daily
■ let dry ■ a second coating may be applied for extra protection ■ to remove, apply more liquid bandage and quickly wipe off ■ fingernail polish remover may dissolve Liquid Bandage

Other information
■ cap bottle tightly ■ store at room temperature away from heat
■ may temporarily sting upon application ■ do not allow to come in contact with hair, countertops, or other finished surfaces — will stain

Inactive ingredients
acetone, myristic acid, castor oil, ethyl acetate, riboflavin, SD alcohol 40

Questions? Call 1-248-449-9300

PRINT SIDE

BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE
liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-045
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q30O)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-045-03	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	12/19/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	12/19/2017	

Labeler - Chain Drug Marketing Association (011920774)