

BAYTRIL- enrofloxacin injection, solution
Elanco US Inc.

Elanco™ Baytril® 100

(enrofloxacin)

**100 mg/mL Antimicrobial
Injectable Solution**

For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle

For Intramuscular Or Subcutaneous Use In Swine

Not For Use In Female Dairy Cattle 20 Months Of Age Or Older

Or In Calves To Be Processed For Veal

CAUTION:

Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.


PRODUCT DESCRIPTION:

Baytril 100 is a sterile, ready-to-use injectable antimicrobial solution that contains enrofloxacin, a broad-spectrum fluoroquinolone antimicrobial agent.

Each mL of Baytril 100 contains 100 mg of enrofloxacin. Excipients are L-arginine base 200 mg, n-butyl alcohol 30 mg, benzyl alcohol (as a preservative) 20 mg and water for injection q.s.

CHEMICAL NOMENCLATURE AND STRUCTURE:

1-cyclopropyl-7-(4-ethyl-1-piperazinyl)-6-fluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid.



INDICATIONS:

Cattle - Single-Dose Therapy: Baytril 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella*

multocida, *Histophilus somni* and *Mycoplasma bovis* in beef and non-lactating dairy cattle; and for the control of BRD in beef and non-lactating dairy cattle at high risk of developing BRD associated with *M. haemolytica*, *P. multocida*, *H. somni* and *M. bovis*.

Cattle - Multiple-Day Therapy: Baytril 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* in beef and non-lactating dairy cattle.

Swine: Baytril 100 is indicated for the treatment and control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Haemophilus parasuis*, *Streptococcus suis*, *Bordetella bronchiseptica* and *Mycoplasma hyopneumoniae*. Baytril 100 is indicated for the control of colibacillosis in groups or pens of weaned pigs where colibacillosis associated with *Escherichia coli* has been diagnosed.

DOSAGE AND ADMINISTRATION:

Baytril 100 provides flexible dosages and durations of therapy.

Baytril 100 may be administered as a single dose for one day for treatment and control of BRD (cattle), for treatment and control of SRD or for control of colibacillosis (swine), or for multiple days for BRD treatment (cattle). Selection of the appropriate dose and duration of therapy for BRD treatment in cattle should be based on an assessment of the severity of the disease, pathogen susceptibility and clinical response.

Cattle:

Single-Dose Therapy (BRD Treatment): Administer, by subcutaneous injection, a single dose of 7.5-12.5 mg/kg of body weight (3.4-5.7 mL/100 lb).

Multiple-Day Therapy (BRD Treatment): Administer daily, a subcutaneous dose of 2.5-5 mg/kg of body weight (1.1-2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for three days. Additional treatments may be given on Days 4 and 5 to animals that have shown clinical improvement but not total recovery.

Single-Dose Therapy (BRD Control): Administer, by subcutaneous injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb).

Examples of conditions that may contribute to calves being at high risk of developing BRD include, but are not limited to, the following:

- Transportation with animals from two or more farm origins.
- An extended transport time with few to no rest stops.
- An environmental temperature change of $\geq 30^{\circ}\text{F}$ during transportation.
- A $\geq 30^{\circ}\text{F}$ range in temperature fluctuation within a 24-hour period.
- Exposure to wet or cold weather conditions.
- Excessive shrink (more than would be expected with a normal load of cattle).
- Stressful arrival processing procedures (e.g., castration or dehorning).
- Exposure within the prior 72 hours to animals showing clinical signs of BRD.

Administered dose volume should not exceed 20 mL per injection site.

Table 1 - Baytril 100 Dose and Treatment Schedule for Cattle*

Weight (lb)	Treatment		Control
	Single-Dose Therapy 7.5 - 12.5 mg/kg Dose Volume (mL)	Multiple-Day Therapy 2.5 - 5.0 mg/kg Dose Volume (mL)	Single-Dose Therapy 7.5 mg/kg Dose Volume (mL)
100	3.5 - 5.5	1.5 - 2.0	3.5
200	7.0 - 11.0	2.5 - 4.5	7.0
300	10.5 - 17.0	3.5 - 6.5	10.5
400	14.0 - 22.5	4.5 - 9.0	14.0
500	17.0 - 28.5	5.5 - 11.5	17.0
600	20.5 - 34.0	7.0 - 13.5	20.5
700	24.0 - 39.5	8.0 - 16.0	24.0
800	27.5 - 45.5	9.0 - 18.0	27.5
900	31.0 - 51.0	10.0 - 20.5	31.0
1000	34.0 - 57.0	11.0 - 23.0	34.0
1100	37.5 - 62.5	12.5 - 25.0	37.5

*Dose volumes have been rounded to the nearest 0.5 mL within the dose range.

Swine:

Administer, either by intramuscular or subcutaneous (behind the ear) injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). Administered dose volume should not exceed 5 mL per injection site.

For the control of colibacillosis, administration should be initiated within the first 60 days post-weaning when clinical signs are present in at least 2% of the animals in the group. If no improvement is noted within 48 hours, the diagnosis should be reevaluated.

Table 2 - Baytril 100 Dose Schedule for Swine

Weight (lb)	Dose Volume (mL)
15	0.5
30	1.0
50	1.7
100	3.4
150	5.1
200	6.8
250	8.5

Dilution of Baytril 100: Baytril 100 may be diluted with sterile water prior to injection. The diluted product should be used within 24 hours. Store diluted solution in amber glass bottles between 4-40°C (36-104°F).

Table 3 - Dilution Schedule*

Swine Weight	mL of Baytril® 100	mL of sterile water	Number of doses
10 lb	34 mL	66 mL	100
15 lb	51 mL	49 mL	100
20 lb	68 mL	32 mL	100

25 lb	85 mL	15 mL	100
-------	-------	-------	-----

*For 1 mL dose volume from diluted solution

Use within 30 days of first puncture and puncture a maximum of 30 times with a needle or 4 times with a dosage delivery device. Any product remaining beyond these parameters should be discarded.

RESIDUE WARNINGS:

Cattle: Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. **Swine:** Animals intended for human consumption must not be slaughtered within 5 days of receiving a single-injection dose.

HUMAN WARNINGS:

Not for use in humans. Keep out of reach of children. Avoid contact with eyes. In case of contact, immediately flush eyes with copious amounts of water for 15 minutes. In case of dermal contact, wash skin with soap and water. Consult a physician if irritation persists following ocular or dermal exposures. Individuals with a history of hypersensitivity to quinolones should avoid this product. In humans, there is a risk of user photosensitization within a few hours after excessive exposure to quinolones. If excessive accidental exposure occurs, avoid direct sunlight. For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

PRECAUTIONS:

The effects of enrofloxacin on cattle or swine reproductive performance, pregnancy and lactation have not been adequately determined. The long-term effects on articular joint cartilage have not been determined in pigs above market weight.

Subcutaneous injection in cattle and swine, or intramuscular injection in swine, can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

Baytril 100 contains different excipients than other Baytril products. The safety and efficacy of this formulation in species other than cattle and swine have not been determined.

Quinolone-class drugs should be used with caution in animals with known or suspected Central Nervous System (CNS) disorders. In such animals, quinolones have, in rare instances, been associated with CNS stimulation which may lead to convulsive seizures. Quinolone-class drugs have been shown to produce erosions of cartilage of weight-bearing joints and other signs of arthropathy in immature animals of various species. See Animal Safety section for additional information.

ADVERSE REACTIONS:

No adverse reactions were observed during clinical trials.

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

MICROBIOLOGY:

Enrofloxacin is bactericidal and exerts its antibacterial effect by inhibiting bacterial DNA gyrase (a type II topoisomerase) thereby preventing DNA supercoiling and replication which leads to cell death.¹ Enrofloxacin is active against Gram-negative and Gram-positive bacteria.

EFFECTIVENESS:

Cattle: A total of 845 calves with naturally-occurring BRD were treated with Baytril 100 in eight field trials located in five cattle-feeding states. Response to treatment was compared to non-treated controls. Single-dose and multiple-day therapy regimens were evaluated. BRD and mortality were significantly reduced in enrofloxacin-treated calves. No adverse reactions were reported in treated animals.

The effectiveness of Baytril 100 for the control of respiratory disease in cattle at high risk of developing BRD was evaluated in a six-location study in the U.S. and Canada. A total of 1,150 crossbred beef calves at high risk of developing BRD were enrolled in the study. Baytril 100 (7.5 mg/kg BW) or an equivalent volume of sterile saline was administered as a single subcutaneous injection within two days after arrival. Cattle were observed daily for clinical signs of BRD and were evaluated for success on Day 14 post-treatment. Treatment success in the Baytril 100 group (497/573, 87.83%) was significantly higher ($P = 0.0013$) than success in the saline control group (455/571, 80.92%). In addition, there were more treatment successes ($n = 13$) than failures ($n = 3$) in the group of animals positive for *M. bovis* on Day 0 that were treated with Baytril 100. No product-related adverse reactions were reported.

Swine: A total of 590 pigs were treated with Baytril 100 or saline in two separate natural infection SRD field trials. For the treatment of SRD, the success rate of enrofloxacin-treated pigs that were defined as “sick and febrile” (increased respiratory rate, labored or dyspneic breathing, depressed attitude and a rectal temperature $\geq 104^{\circ}\text{F}$) was statistically significantly greater than the success rate of saline-treated “sick and febrile” pigs. For the control of SRD, mean rectal temperature, mortality (one trial) and morbidity were statistically significantly lower for enrofloxacin-treated pigs in pens containing a percentage of “sick and febrile” pigs compared to saline-treated pigs.

The effectiveness of Baytril 100 administered as a single SC dose of 7.5 mg/kg BW for the treatment and control of SRD associated with *M. hyopneumoniae* was demonstrated using an induced infection model study and three single-site natural infection field studies. In the model study, 72 healthy pigs were challenged with a representative *M. hyopneumoniae* isolate and treated with Baytril 100 or saline. A statistically significant ($P < 0.0001$) decrease in the mean total lung lesion score was observed in the Baytril 100-treated group (4%) compared with the saline-treated group (27%) at 10 days post-treatment. In two field studies evaluating effectiveness for treatment of SRD, a total of 300 pigs with clinical signs of SRD (moderate depression, moderately increased

respiratory rate, and a rectal temperature of $\geq 104^{\circ}\text{F}$) were enrolled and treated with Baytril 100 or saline. At 7 days post-treatment, the cure rate was statistically significantly higher at each site ($P < 0.0001$) in the Baytril 100-treated groups (61.3% and 92%) compared with the saline-treated groups (26.7% and 33.3%). In one field study evaluating effectiveness for control of SRD, a group of 400 pigs in which $> 15\%$ had clinical signs of SRD (moderate depression score, moderately increased respiratory rate, and a rectal temperature of $\geq 104^{\circ}\text{F}$) was enrolled and treated with Baytril 100 or saline. At 7 days post-treatment, the cure rate was statistically significantly higher ($P < 0.0002$) in the Baytril 100-treated group (70.0%) compared with the saline-treated group (48.5%). In addition to *M. hyopneumoniae*, *B. bronchiseptica* was also isolated in sufficient numbers from these field studies to be included in the SRD treatment and control indications.

The effectiveness of Baytril 100 for the control of colibacillosis associated with *E. coli* was evaluated in a multi-site natural infection field study. At each site, when at least 5% of the pigs were defined as “clinically affected” (presence of diarrhea and either depression or gauntness), all pigs were administered Baytril 100 as a single IM dose of 7.5 mg/kg BW or an equivalent dose volume of saline. At 7 days post-treatment, the success rate was statistically significantly higher ($P = 0.0350$) in the Baytril 100-treated group (61.5%) compared with the saline-treated group (44.7%).

The effectiveness of Baytril 100 administered as a single IM dose of 7.5 mg/kg BW for the treatment and control of SRD or as a single SC dose of 7.5 mg/kg BW for the control of colibacillosis was confirmed by demonstrating comparable serum enrofloxacin concentrations following IM or SC injection into the neck of healthy male and female pigs.

TOXICOLOGY:

The oral LD50 for laboratory rats was greater than 5000 mg/kg of body weight. Ninety-day feeding studies in dogs and rats revealed no observable adverse effects at treatment rates of 3 and 40 mg/kg respectively. Chronic studies in rats and mice revealed no observable adverse effects at 5.3 and 323 mg/kg respectively. There was no evidence of carcinogenic effect in laboratory animal models. A two-generation rat reproduction study revealed no effect with 10 mg/kg treatments. No teratogenic effects were observed in rabbits at doses of 25 mg/kg or in rats at 50 mg/kg.

ANIMAL SAFETY:

Cattle: Safety studies were conducted in feeder calves using single doses of 5, 15 and 25 mg/kg for 15 consecutive days and 50 mg/kg for 5 consecutive days. No clinical signs of toxicity were observed when a dose of 5 mg/kg was administered for 15 days. Clinical signs of depression, incoordination and muscle fasciculation were observed in calves when doses of 15 or 25 mg/kg were administered for 10 to 15 days. Clinical signs of depression, inappetance and incoordination were observed when a dose of 50 mg/kg was administered for 3 days. No drug-related abnormalities in clinical pathology parameters were identified. No articular cartilage lesions were observed after examination of stifle joints from animals administered 25 mg/kg for 15 days.

A safety study was conducted in 23-day-old calves using doses of 5, 15 and 25 mg/kg for 15 consecutive days. No clinical signs of toxicity or changes in clinical pathology parameters were observed. No articular cartilage lesions were observed in the stifle

joints at any dose level at 2 days and 9 days following 15 days of drug administration.

An injection site study conducted in feeder calves demonstrated that the formulation may induce a transient reaction in the subcutaneous tissue and underlying muscle. No painful responses to administration were observed.

Swine: Subcutaneous Safety: A safety study was conducted in 32 pigs weighing approximately 57 kg (125 lb) using single doses of 5, 15 or 25 mg/kg daily for 15 consecutive days. Incidental lameness of short duration was observed in all groups, including the saline-treated controls. Musculoskeletal stiffness was observed following the 15 and 25 mg/kg treatments with clinical signs appearing during the second week of treatment. Clinical signs of lameness improved after treatment ceased and most animals were clinically normal at necropsy.

A second study was conducted in two pigs weighing approximately 23 kg (50 lb), treated with 50 mg/kg for 5 consecutive days. There were no clinical signs of toxicity or pathological changes.

An injection site study conducted in pigs demonstrated that the formulation may induce a transient reaction in the subcutaneous tissue. No painful responses to administration were observed.

Intramuscular Safety: A safety study was conducted in 48 weaned, 20- to 22-day-old pigs. Pigs were administered Baytril 100, at 7.5, 22.5 and 37.5 mg/kg BW by IM injection into the neck once weekly for 3 consecutive weeks. All pigs remained clinically normal throughout the study. Transient decreases in feed and water consumption were observed after each treatment. Mild, transient, post-treatment injection site swellings were observed in pigs receiving the 37.5 mg/kg BW dose. Injection site inflammation was found on post-mortem examination in all enrofloxacin-treated groups.

STORAGE CONDITIONS:

Protect from direct sunlight. Do not refrigerate or freeze. Store at 20-30°C (68-86°F), excursions permitted up to 40°C (104°F). Precipitation may occur due to cold temperature. To redissolve, warm and then shake the vial.

HOW SUPPLIED:

Baytril® 100:

100 mg/mL	100 mL Bottle
100 mg/mL	250 mL Bottle
100 mg/mL	500 mL Bottle

REFERENCES:

1. Hooper, D. C., Wolfson, J. S., Quinolone Antimicrobial Agents, 2nd ed, 59 - 75, 1993.

For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS),
call Elanco Product & Veterinary Support at 1-800-428-4441.

Elanco

90206956_PA600217X

Revised: January 2022

Baytril is sold by Elanco or its affiliates and is not a product of Bayer. The product name Baytril is owned by Bayer and used under license.

Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2022 Elanco or its affiliates.

Baytril® 100

Approved by FDA under NADA # 141-068

Manufactured for: Elanco US Inc.

Greenfield, IN 46140 U.S.A

Made in Germany

Principal Display Panel

Baytril® 100

(enrofloxacin)

**100 mg/mL Antimicrobial
Injectable Solution**

**For Subcutaneous Use In Beef Cattle And
Non-Lactating Dairy Cattle**

For Intramuscular Or Subcutaneous Use In Swine

**Not For Use In Female Dairy Cattle
20 Months Of Age Or Older
Or In Calves To Be Processed For Veal**

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.


100 mL

Manufactured for: Elanco US Inc.

Greenfield, IN 46140 U.S.A

Made in Germany

Approved by FDA under NADA # 141-068



Baytril® 100

(enrofloxacin)

100 mg/mL Antimicrobial Injectable Solution

For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle

For Intramuscular Or Subcutaneous Use In Swine

Not For Use In Female Dairy Cattle 20 Months Of Age Or Older

Or In Calves To Be Processed For Veal

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian. Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A Made in Germany
Approved by FDA under NADA # 141-068

100 mL

Elanco

Baytril® 100
(enrofloxacin)

100 mg/mL Antimicrobial Injectable Solution

**For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle
For Intramuscular Or Subcutaneous Use In Swine
Not For Use In Female Dairy Cattle 20 Months Of Age Or Older
Or In Calves To Be Processed For Veal**

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian. Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals. To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A Made in Germany
Approved by FDA under NADA # 141-068

100 mL

Elanco

INDICATIONS: For the treatment and control of respiratory diseases in cattle and swine. For the control of colibacillosis in swine. See package insert for complete indications for use.
CONTAINS PER mL: Enrofloxacin 100 mg. Excipients: L-arginine base 200 mg, n-butyl alcohol 30 mg, benzyl alcohol (as a preservative) 20 mg and water for injection q.s.
DOSAGE AND ADMINISTRATION:
Cattle: Single-Dose Therapy (BRD Treatment): Administer, by subcutaneous injection, a single dose of 7.5-12.5 mg/kg of body weight (3.4-5.7 mL/100 lb).
Multiple-Day Therapy (BRD Treatment): Administer daily, a subcutaneous dose of 2.5-5 mg/kg of body weight (1.1-2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for 3-5 days.
Single-Dose Therapy (BRD Control): Administer, by subcutaneous injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). See insert for examples of conditions that contribute to high risk. Administered dose volume should not exceed 20 mL per injection site.
Swine: Administer either by intramuscular or subcutaneous (behind the ear) injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). Administered dose volume should not exceed 5 mL per injection site. For the control of colibacillosis, administration should be initiated within the first 60 days post-weaning when clinical signs are present in at least 2% of the animals in the group. See package insert for full dosage and administration information. Use within 30 days of first puncture and puncture a maximum of 30 times with a needle or 4 times with a dosage delivery device. Any product remaining beyond these parameters should be discarded.

RESIDUE WARNINGS:
Cattle: Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.
Swine: Animals intended for human consumption must not be slaughtered within 5 days of receiving a single-injection dose.

HUMAN WARNINGS: Not for use in humans. Keep out of reach of children.
For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

STORAGE CONDITIONS: Protect from direct sunlight. Do not refrigerate or freeze. Store at 20-30°C (68-86°F), excursions permitted up to 40°C (104°F). Precipitation may occur due to cold temperature. To redissolve, warm and then shake the vial. Read package insert carefully for complete details.

90206951_YL600217A

B8U7
Lot No.:
Exp. Date:

Baytril® 100

(enrofloxacin)

**100 mg/mL Antimicrobial
Injectable Solution**

**For Subcutaneous Use In Beef Cattle And
Non-Lactating Dairy Cattle**

For Intramuscular Or Subcutaneous Use In Swine

**Not For Use In Female Dairy Cattle
20 Months Of Age Or Older
Or In Calves To Be Processed For Veal**

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

250 mL

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A
Made in Germany

Approved by FDA under NADA # 141-068

90206958_SH600217D

Baytril® 100

(enrofloxacin)



100 mg/mL Antimicrobial
Injectable Solution

For Subcutaneous Use in Beef Cattle And
Non-Lactating Dairy Cattle

For Intramuscular Or Subcutaneous Use In Swine

Not For Use In Female Dairy Cattle
20 Months Of Age Or Older
Or In Calves To Be Processed For Veal

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

250 mL

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A.
Made In Germany
Approved by FDA under NADA # 141-058

Elanco



Lot No.:
Exp. Date:

Baytril 100 Dose and Treatment Schedule for Cattle*

Weight (lb)	Treatment		Control
	Single-Dose Therapy 7.5 - 12.5 mg/kg Dose Volume (mL)	Multiple-Day Therapy 2.5 - 5.0 mg/kg Dose Volume (mL)	
100	3.5 - 5.5	1.5 - 2.0	3.5
200	7.0 - 11.0	2.5 - 4.5	7.0
300	10.5 - 17.0	3.5 - 6.5	10.5
400	14.0 - 22.5	4.5 - 9.0	14.0
500	17.0 - 28.5	5.5 - 11.5	17.0
600	20.5 - 34.0	7.0 - 13.5	20.5
700	24.0 - 39.5	8.0 - 16.0	24.0
800	27.5 - 45.5	9.0 - 18.0	27.5
900	31.0 - 51.0	10.0 - 20.5	31.0
1000	34.0 - 57.0	11.0 - 23.0	34.0
1100	37.5 - 62.5	12.5 - 25.0	37.5

*Dose volumes have been rounded to the nearest 0.5 mL within the dose range.

Baytril 100 Dose Schedule for Swine

Weight (lb)	Dose Volume (mL)
35	0.5
39	1.0
53	1.7
100	3.4
150	5.1
200	6.8
250	8.5

100 mg/mL Antimicrobial
Injectable Solution



Baytril® 100

(enrofloxacin)

Baytril® 100

(enrofloxacin)



100 mg/mL Antimicrobial
Injectable Solution

DOSAGE AND ADMINISTRATION:

Cattle:

Single-Dose Therapy (BRD Treatment): Administer, by subcutaneous injection, a single dose of 7.5-12.5 mg/kg of body weight (3.4-5.7 mL/100 lb).

Multiple-Day Therapy (BRD Treatment): Administer daily, a subcutaneous dose of 2.5-5 mg/kg of body weight (1.1-2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for three days. Additional treatments may be given on Days 4 and 5 to animals that have shown clinical improvement but not total recovery.

Single-Dose Therapy (BRD Control): Administer, by subcutaneous injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). See insert for examples of conditions that contribute to high risk. Administered dose volume should not exceed 20 mL per injection site.

Swine:

Administer, either by intramuscular or subcutaneous (behind the ear) injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). Administered dose volume should not exceed 5 mL per injection site. For the control of colibacillosis, administration should be initiated within the first 60 days post-weaning when clinical signs are present in at least 2% of the animals in the group. See package insert for full dosage and administration information.

INDICATIONS:

Cattle: Baytril 100 is indicated in beef and non-lactating dairy cattle for:

Single-Dose Therapy: the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* in beef and non-lactating dairy cattle; and for the control of BRD in beef and non-lactating dairy cattle at high risk of developing BRD associated with *M. haemolytica*, *P. multocida*, *H. somni* and *M. bovis*.

Multiple-Day Therapy: the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* in beef and non-lactating dairy cattle.

Swine: Baytril 100 is indicated for:

The treatment and control of swine respiratory disease (SRD) associated with *Aerobacillus pleuropneumoniae*, *Pasteurella multocida*, *Aeromonas parasuis*, *Streptococcus suis*, *Bordetella bronchiseptica* and *Mycoplasma hyopneumoniae*. The control of colibacillosis in groups or pens of weaned pigs where colibacillosis associated with *Escherichia coli* has been diagnosed.

CONTAINS PER mL: Enrofloxacin 100 mg, Excipients: L-arginine base 200 mg, n-butyl alcohol 30 mg, benzyl alcohol (as a preservative) 20 mg and water for injection q.s.

Use within 30 days of first puncture and puncture a maximum of 30 times with a needle or 4 times with a dosage delivery device. Any product remaining beyond these parameters should be discarded.

RESIDUE WARNINGS:

Cattle: Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Swine: Animals intended for human consumption must not be slaughtered within 5 days of receiving a single-injection dose.

HUMAN WARNINGS: Not for use in humans. Keep out of reach of children. See package insert.

For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

STORAGE CONDITIONS: Protect from direct sunlight. Do not refrigerate or freeze. Store at 20-30°C (68-86°F), excursions permitted up to 40°C (104°F). Precipitation may occur due to cold temperature. To redissolve, warm and then shake vial.

Read package insert carefully for complete details.

Baytril® 100

(enrofloxacin)

100 mg/mL Antimicrobial Injectable Solution

**For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle
For Intramuscular Or Subcutaneous Use In Swine
Not For Use In Female Dairy Cattle 20 Months Of Age Or Older
Or In Calves To Be Processed For Veal**

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian. Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A Made in Germany
Approved by FDA under NADA # 141-068

250 mL

Elanco

Baytril® 100

(enrofloxacin)

100 mg/mL Antimicrobial Injectable Solution

For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle

For Intramuscular Or Subcutaneous Use In Swine

Not For Use In Female Dairy Cattle 20 Months Of Age Or Older Or In Calves To Be Processed For Veal

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A
Made in Germany
Approved by FDA under NADA # 141-068

500 mL

INDICATIONS: For the treatment and control of respiratory diseases in cattle and swine. For the control of colibacillosis in swine. See package insert for complete indication use.

Baytril[®] 100
(enrofloxacin)

100 mg/mL Antimicrobial Injectable Solution

500 mL

For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle
For Intramuscular Or Subcutaneous Use In Swine
Not For Use In Female Dairy Cattle 20 Months Of Age Or Older Or In Calves To Be Processed For Veal

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.
Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.
To assure responsible antimicrobial drug use, enrofloxacin should only be used as a second-line drug for colibacillosis in swine following consideration of other therapeutic options.

Manufactured for: Elanco US Inc.
Greenfield, IN 46140 U.S.A
Made in Germany
Approved by FDA under NADA # 141-068

INDICATIONS: For the treatment and control of respiratory diseases in cattle and swine. For the control of colibacillosis in swine. See package insert for complete indications for use.

CONTAINS PER mL: Enrofloxacin 100 mg. Excipients: L-arginine base 200 mg, n-butyl alcohol 30 mg, benzyl alcohol (as a preservative) 20 mg and water for injection q.s.

DOSAGE AND ADMINISTRATION:
Cattle: Single-Dose Therapy (BRD Treatment): Administer, by subcutaneous injection, a single dose of 7.5–12.5 mg/kg of body weight (3.4–5.7 mL/100 lb).
Multiple-Day Therapy (BRD Treatment): Administer daily, a subcutaneous dose of 2.5–5 mg/kg of body weight (1.1–2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for 3–5 days.
Single-Dose Therapy (BRD Control): Administer, by subcutaneous injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). See insert for examples of conditions that contribute to high risk. Administered dose volume should not exceed 20 mL per injection site.
Swine: Administer, either by intramuscular or subcutaneous (behind the ear) injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). Administered dose volume should not exceed 5 mL per injection site. For the control of colibacillosis, administration should be initiated within the first 60 days post-weaning when clinical signs are present in at least 2% of the animals in the group. See package insert for full dosage and administration information.

RESIDUE WARNINGS:
Cattle: Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.
Swine: Animals intended for human consumption must not be slaughtered within 5 days of receiving a single-injection dose.

HUMAN WARNINGS: Not for use in humans. Keep out of reach of children.
For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

STORAGE CONDITIONS: Protect from direct sunlight. Do not refrigerate or freeze. Store at 20–30°C (68–86°F), excursions permitted up to 40°C (104°F). Precipitation may occur due to cold temperature. To redissolve, warm and then shake the vial.

Read attached booklet carefully for complete details.

90206988_YL600217E

Elanco

BAYTRIL

enrofloxacin injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:58198-0021
Route of Administration	SUBCUTANEOUS, INTRAMUSCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ENROFLOXACIN (UNII: 3DX3XEK1BN) (ENROFLOXACIN - UNII:3DX3XEK1BN)	ENROFLOXACIN	100 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0021-1	1 in 1 CARTON		
1	NDC:58198-0021-3	100 mL in 1 BOTTLE		
2	NDC:58198-0021-2	1 in 1 CARTON		
2	NDC:58198-0021-4	250 mL in 1 BOTTLE		
3	NDC:58198-0021-5	500 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141068	07/24/1998	

Labeler - Elanco US Inc. (966985624)

Establishment

Name	Address	ID/FEI	Business Operations
TriRx Shawnee LLC		118187894	MANUFACTURE, ANALYSIS, PACK, LABEL

Establishment

Name	Address	ID/FEI	Business Operations
Bayer AG		323208116	API MANUFACTURE

Revised: 10/2022

Elanco US Inc.