

GM COLLIN PURACNE- salicylic acid gel
Laboratoires Dermo-Cosmetik Inc.

G.M. COLLIN ® PURACNE GEL

Product / Drug Facts

Active ingredient

Salicylic Acid 2.0%

Purpose

Acne Treatment

Uses

- Helps treat acne pimples.
- Dries and helps clear up acne pimples.
- Helps prevent new acne pimples.

Warnings

For external use only.

Do not use if you have very sensitive skin.

When using this product

- Avoid contact with the eyes, lips and mouth; if contact occurs, rinse thoroughly with water.
- Using other topical acne drugs at the same time or right after use of this product may increase dryness or irritation of the skin. If this occurs, only one product should be used unless otherwise directed by a doctor.
- **Skin reactions:** This product may cause irritation, burning, itching, swelling and blistering of the skin. If any of the above noted symptoms occur, discontinue use.

Stop use and ask a doctor if

- The skin reaction becomes severe or spreads beyond the site of application.
- You develop severe itching, hives, swelling of eyes or mouth, blistering or difficulty breathing.
- No substantial improvement occurs within 6 to 8 weeks.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash hands with a non-medicinal soap.

- Clean the skin thoroughly with the appropriate G.M. Collin cleanser before applying this product.
- For New Users: Apply product to a small area once a day for three days to test if you are sensitive to this product.
- Apply morning and/or evening on the affected areas, or as recommended by your skin care professional.
- Start with one application daily, then increase if needed.
- If bothersome dryness or peeling occurs, reduce application to once a day or every other day.

Other information

Store at 15-30°C (59-86°F).

Inactive ingredients

AMMONIUM ACRYLOYLDIMETHYLTAURATE/VP COPOLYMER, AQUA/WATER/EAU, BIOSACCHARIDE GUM-1, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, CARBOMER, CINNAMOMUM CAMPHORA (CAMPHOR) BARK OIL, CUPRESSUS SEMPERVIRENS OIL, DIMETHICONE, ETHOXYDIGLYCOL, GLYCERIN, LAVANDULA ANGUSTIFOLIA (LAVENDER) OIL, MADECASSOSIDE, NIACINAMIDE, NORDIHYDROGUAIARETIC ACID, OLEANOLIC ACID, PEG-40 HYDROGENATED CASTOR OIL, PEG-60 ALMOND GLYCERIDES, PEG-8, PELARGONIUM GRAVEOLENS FLOWER OIL, PENTYLENE GLYCOL, PEUMUS BOLDUS LEAF EXTRACT, ROSMARINUS OFFICINALIS (ROSEMARY) LEAF OIL, SCLEROTIUM GUM, SODIUM HYALURONATE, TRIETHANOLAMINE, TROPAEOLUM MAJUS FLOWER/LEAF/STEM EXTRACT, XANTHAN GUM.

Questions?

USA: 1 (800) 341-1531 /
CANADA: 1 (800) 361-1263

www.gmcollin.com
MADE IN CANADA
DISTRIBUÉ EXCLUSIVEMENT PAR /
EXCLUSIVELY DISTRIBUTED BY:
® LABORATOIRES DERMO-COSMETIK INC.
68 STINSON, MONTREAL, QUEBEC,
CANADA H4N 2E7

PRINCIPAL DISPLAY PANEL - 50 ml Tube Carton

G.M. COLLIN

PURACNE
GEL

Acne Treatment

NDC 64127-195-01

NPN 80086999

50 ml - 1.7 fl. oz

PURACNE GEL
6V1143

G.M. COLLIN

COLLIN[®]
G.M.

Information sur le produit

Ingédient actif :
Acide salicylique 2.0 %
Utilité : Traitement acné

Usages

- aide à traiter les boutons d'acné.
- assèche et aide à éliminer les boutons d'acné.
- aide à prévenir de nouveaux boutons d'acné.

Mises en garde

Pour usage externe seulement.
Ne pas utiliser si vous avez la peau très sensible.

Lorsque vous utilisez ce produit :

- éviter le contact avec les yeux, les lèvres et la bouche; En cas de contact, rincer abondamment à l'eau.
- l'utilisation d'autres médicaments topiques contre l'acné en même temps ou juste après l'utilisation de ce produit peut augmenter la sécheresse ou l'irritation de la peau. Si cela se produit, un seul produit doit être utilisé, sauf indication contraire d'un médecin.
- **Réactions cutanées :** Ce produit peut provoquer une irritation, des brûlures, des démangeaisons, un gonflement et des cloques sur la peau. Si l'un des symptômes mentionnés ci-dessus apparaît, arrêtez l'utilisation.

Cesser d'utiliser et consulter un médecin si :

- la réaction cutanée devient sévère ou s'étend au-delà du site d'application.
- vous développez de fortes démangeaisons, de l'urticaire, un gonflement des yeux ou de la bouche, des cloques ou des difficultés respiratoires.
- aucune amélioration substantielle ne se produit dans les 6 à 8 semaines.

Garder hors de la portée des enfants. En cas d'ingestion, demander de l'aide médicale ou contacter immédiatement un centre antipoison.

Mode d'emploi

- Se laver les mains avec un savon non médicamenteux.
- Bien nettoyer la peau avec le nettoyant G.M. Collin approprié avant d'appliquer le produit.
- Pour les nouveaux utilisateurs: appliquer le produit sur une petite surface, une fois par jour pendant trois jours afin de tester la sensibilité à ce produit.
- Appliquer matin et/ou soir sur les zones affectées, ou selon les recommandations de votre professionnel en soins de la peau.



PURACNE

GEL
Traitement pour l'acné

GEL
Acne Treatment

NDC 64127-195-01
NPN 80086999

50 ml - 1.7 fl. oz.

Information sur le produit (suite)

- Commencer par une application par jour, puis augmenter au besoin.
- En cas de sécheresse ou de desquamation, réduire l'application à une fois par jour ou à une fois tous les deux jours.

Autres renseignements
• Conserver à 15-30°C (59-86°F)

Ingédients inactifs :
voir « Product Facts / Drug Facts »

Questions ?
USA: 1 (800) 341-1631
CANADA: 1 (800) 361-1263

www.gmcollin.com
MADE IN CANADA
DISTRIBUÉ EXCLUSIVEMENT PAR /
EXCLUSIVELY DISTRIBUTED BY:
© LABORATOIRES DERMO-COSMETIK INC.
66 STINSON, MONTREAL, QUEBEC,
CANADA H4N 2E7



Product Facts / Drug Facts

Active ingredient: Salicylic Acid 2.0% **Purpose:** Acne Treatment

Uses

- helps treat acne pimples.
- dries and helps clear up acne pimples.
- helps prevent new acne pimples.

Warnings

For external use only.

Do not use if you have very sensitive skin.

When using this product:

- avoid contact with the eyes, lips and mouth; if contact occurs, rinse thoroughly with water.
- using other topical acne drugs at the same time or right after use of this product may increase dryness or irritation of the skin. If this occurs, only one product should be used unless otherwise directed by a doctor.
- **Skin reactions:** This product may cause irritation, burning, itching, swelling and blistering of the skin. If any of the above noted symptoms occur, discontinue use.

Stop use and ask a doctor if:

- the skin reaction becomes severe or spreads beyond the site of application.
- you develop severe itching, hives, swelling of eyes or mouth, blistering or difficulty breathing.
- no substantial improvement occurs within 6 to 8 weeks.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash hands with a non-medical soap.
- Clean the skin thoroughly with the appropriate G.M. Collin cleanser before applying this product.
- For New Users: Apply product to a small area once a day for three days to test if you are sensitive to this product.
- Apply morning and/or evening on the affected areas, or as recommended by your skin care professional.
- Start with one application daily, then increase if needed.
- If bothersome dryness or peeling occurs, reduce application to once a day or every other day.

Other information
• Store at 15-30°C (59-86°F)

Inactive ingredients

AMMONIUM ACRYLOYLDIMETHYLTAURATE/VP COPOLYMER, AQUA/WATER/EAU, BISACRYLATE DIMETHACRYLATE, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, CARBOMER, CINNAMOMUM CAMPHORA (CAMPHOR) BARK OIL, CUPRESSUS SEMPERVIRENS OIL, DIMETHICONE, ETHOXYGLYCOL, GLYCERIN, LAVANDULA ANGUSTIFOLIA (LAVENDER) OIL, MADECASSOSIDE, NIACINAMIDE, NORDIHYDROGUAIARIC ACID, OLEFANOLIC ACID, PEG-40 HYDROGENATED CASTOR OIL, PEG-60 ALMOND GLYCERIDES, PEG-8, PELARGONIUM GRAVEOLENS FLOWER OIL, PENTYLENE GLYCOL, PEUMUS BOLDUS LEAF EXTRACT, ROSMARINUS OFFICINALIS (ROSEMARY) LEAF OIL, SCLEROTIUM BUII, SODIUM HYALURONATE, TRIETHANOLAMINE, TROPACOLUM MAJUS FLOWER/LEAF/STEM EXTRACT, XANTHAN GUM.



GM COLLIN PURACNE

salicylic acid gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:64127-195
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: 0414PZ4LPZ) (SALICYLIC ACID - UNII:0414PZ4LPZ)	SALICYLIC ACID	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
AMMONIUM ACRYLOYLDIMETHYLTAURATE/VP COPOLYMER (UNII: W59H9296ZG)	
BIOSACCHARIDE GUM-1 (UNII: BB4PU4V09H)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)	
CAMPHOR OIL (UNII: 75IZZ8Y727)	
DIMETHICONE (UNII: 92RU3N3Y10)	
DIETHYLENE GLYCOL MONOETHYL ETHER (UNII: A1A1I8X02B)	
GLYCERIN (UNII: PDC6A3C0OX)	
MADECASSOSIDE (UNII: CQ2F5O6YIY)	
NIACINAMIDE (UNII: 25X51I8RD4)	
OLEANOLIC ACID (UNII: 6SMK8R7TGJ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
PEG-60 ALMOND GLYCERIDES (UNII: 4Y0E651N0F)	
PELARGONIUM GRAVEOLENS FLOWER OIL (UNII: 3K0J1S7QGC)	
PENTYLENE GLYCOL (UNII: 50C1307PZG)	
BETASIZOFIRAN (UNII: 2X51AD1X3T)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
TROLAMINE (UNII: 9O3K93S3TK)	
XANTHAN GUM (UNII: TTV12P4NEE)	
WATER (UNII: 059QF0KO0R)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
BIOTIN (UNII: 6S06U10H04)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
PEUMUS BOLDUS LEAF (UNII: Q4EWM09M3O)	
TROPAEOLUM MAJUS FLOWERING TOP (UNII: RGT30824HY)	

ROSEMARY OIL (UNII: 8LGU7VM393)	
MASOPROCOL (UNII: 7BO8G1BYQU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
LAVENDER OIL (UNII: ZBP1YXW0H8)	
CUPRESSUS SEMPERVIRENS LEAF OIL (UNII: M7QUY89S4O)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64127-195-01	1 in 1 CARTON	11/26/2018	
1		50 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M006	11/26/2018	

Labeler - Laboratoires Dermo-Cosmetik Inc. (249335480)

Revised: 10/2023

Laboratoires Dermo-Cosmetik Inc.