

**ICY HOT WITH LIDOCAINE- lidocaine and menthol patch
Chattem, Inc.**

Icy Hot with Lidocaine Patch

ICY HOT with 4% LIDOCAINE PATCH

Drug Facts

Active ingredients

Lidocaine 4%

Menthol 1%

Purposes

Topical anesthetic

Topical analgesic

Use

for the temporary relief of pain

Warnings

For external use only

Do not use

- on puncture wounds, cuts, irritated, damaged or swollen skin
- more than 1 patch on your body at a time or with other topical analgesics at the same time
- with a heating pad or apply local heat to the area of use

When using this product

- use only as directed
- do not bandage tightly
- avoid contact with the eyes and mucous membranes
- rare cases of serious burns have been reported with products of this type
- a transient burning sensation may occur upon application but generally disappears in several days
- dispose of used patch in manner that always keeps product away from children and

pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- severe burning sensation, redness or irritation develops
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and older:

- clean and dry affected area
- remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
- carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area
- once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area
- use 1 patch at a time and not more than 3 to 4 times daily

children under 12 years of age: consult a doctor

Inactive ingredients

aluminum glycinate, aluminum hydroxide, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, methylparaben, nonoxynol-30, polyacrylic acid, polysorbate 80, propylene glycol, silica, sodium polyacrylate, tartaric acid, titanium dioxide, urea, water

Child-resistant packaging.

Keep carton as it contains important information.

PRINCIPAL DISPLAY PANEL

ICYHOT

LIDOCAINE PATCH PLUS MENTHOL



ICY HOT WITH LIDOCAINE

lidocaine and menthol patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41167-1720
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	240 mg
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	60 mg

Inactive Ingredients

Ingredient Name	Strength
2-ETHYLHEXYL ACRYLATE (UNII: HR49R9S6XG)	
METHYL ACRYLATE (UNII: WC487PR91H)	
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IU00)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	

GLYCERIN (UNII: PDC6A3C00X)
METHYLPARABEN (UNII: A2I8C7HI9T)
NONOXYNOL-30 (UNII: JJX07DG188)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)
TARTARIC ACID (UNII: W4888I119H)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
UREA (UNII: 8W8T17847W)
WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41167-1720-1	5 in 1 CARTON; Type 0: Not a Combination Product	02/18/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/18/2018	

Labeler - Chattem, Inc. (003336013)

Revised: 10/2023

Chattem, Inc.