

MUCOSAN MULTI-SYMPTOM CHERRY- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid
Menper Distributors, Inc.

Mucosan Multi-Symptom Cherry

Active ingredients (in each 5 mL or teaspoon) Purpose

Dextromethorphan HBr, 10 mg.....Cough suppressant

Guaifenesin, 100 mg.....Expectorant

Phenylephrine HCl, 5 mg.....Nasal decongestant

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

Do not exceed recommended dosage.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast - feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- TSP=teaspoon

| | |
|---------------------------------------|----------------------------------|
| age | dose |
| adults and children 12 years and over | 2 teaspoons (10mL) every 4 hours |
| children under 12 years | do not use |

Other information

- **each teaspoon (5mL) contains:** sodium 3 mg
- store at 15-30°C (59-86°F). Do not refrigerate.

anhydrous citric acid, FD&C red #40, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

1-800-560-5223 Monday to Friday 9am-4pm, Eastern Time

MucoSan
multi-sympton
multi-sintoma

Dextromethorphan HBr
Guafenesin
Phenylephrine HCl
Dextrometorfano HBr
Guafenesina
Fenilefrina HCl

COUGH & CONGESTION
TOS Y CONGESTIÓN

RELIEVES / ALIVIA:
Cough/Tos
Nasal Congestion/
Congestión Nasal
Chest Congestion/
Congestión de Pecho

12+
years/
años

4 FL oz (118 mL)

relief · alivio

Drug Facts

Active Ingredients (in each 5 mL or 1 teaspoon)
Dextromethorphan HBr, 10 mg..... Cough Suppressant
Guafenesin, 100 mg..... Expectorant
Phenylephrine HCl, 5 mg..... Nasal Decongestant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes ■ temporarily relieves these symptoms occurring with a cold ■ nasal congestion ■ cough due to minor throat and bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.
Do not exceed recommended dosage.
Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ symptoms do not get better within 7 days or are accompanied by fever ■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24 hour period ■ take adult strength product is not intended for use in children under 12 years of age ■ measure only with dosing cup provided ■ keep dosing cup with product ■ TSP = teaspoon

| | |
|---------------------------------------|-----------------------------------|
| age | dose |
| adults and children 12 years and over | 2 teaspoons (10 mL) every 4 hours |
| children under 12 years | do not use |

Other information ■ each teaspoon (5mL) contains: sodium 3 mg ■ store at 15-30°C (59-86°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, FD&C red #40, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?
1-800-560-5223 Monday to Friday 9am-4pm, Eastern Time

Distributed by / Distribuido por: The Menper Group, 11421 NW107th ST, Miami, FL 33178, USA
Include información en español

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING
NO SE USE SI EL SELLO IMPRESO DEBAJO DE LA TAPA ESTÁ ROTO O FALTA

Rev. 1
PEEL HERE
LEVANTE AQUÍ

Lot
Exp

Muco san

multi-sympton
multi-sintoma

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING / NO SE USE SI EL SELLO
IMPRESO DEBAJO DE LA TAPA ESTA ROTO O FALTA

Muco san

multi-sympton
multi-sintoma

Dextromethorphan HBr/Guaifenesin/
Phenylephrine HCl Dextrometorfano HBr/
Guaifenesina/Fenilefrina HCl



**COUGH &
CONGESTION**
TOS Y CONGESTIÓN

RELIEVES / ALIVIA:

Cough/Tos

Nasal Congestion/
Congestión Nasal

Chest Congestion/
Congestión de Pecho

12+
years/
años

4 FL oz (118 mL)

Drug Facts

Active ingredients (in each 5 mL or 1 teaspoon) Purposes
Dextromethorphan HBr, 10 mg.....Cough Suppressant
Guaifenesin, 100 mg.....Expectorant
Phenylephrine HCl, 5 mg.....Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Do not exceed recommended dosage.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- TSP = teaspoon

| age | dose |
|---------------------------------------|-----------------------------------|
| adults and children 12 years and over | 2 teaspoons (10 mL) every 4 hours |
| children under 12 years | do not use |

Drug Facts (continued)

Other information

- each teaspoon (5 mL) contains: sodium 3 mg
- store at 15-30°C (59-86°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

1-800-560-5223 Monday to Friday 9am-4pm, Eastern Time

Información del Medicamento

Ingredientes activos (en cada 5 mL o 1 cucharadita) Propósitos

Dextrometorfano HBr, 10 mg.....Supresor de la Tos
Guaifenesina, 100 mg.....Expectorante
Fenilefrina HCl, 5 mg.....Descongestionante Nasal

Usos

- ayuda a desprender las flemas (mucosidad) y a adelgazar las secreciones bronquiales para drenar las vías respiratorias.
- alivia temporalmente los siguientes síntomas que se presentan con el resfriado:
 - congestión nasal
 - tos debida a irritación menor de garganta y bronquios

Advertencias

No utilizarlo si usted está tomando un inhibidor de la monoamino oxidasa (IMAO) de prescripción (ciertos medicamentos para la depresión, condiciones psiquiátricas o emocionales, o enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento IMAO. Si usted desconoce si su medicamento de prescripción contiene un IMAO, consulte a su médico o farmacéutico antes de tomar este producto.

Consulte a su médico antes de usar, si padece

- enfermedad cardíaca
- hipertensión
- enfermedad de la tiroides
- diabetes
- dificultad al orinar debido a dilatación de la glándula prostática
- tos que se presenta con exceso de flema (mucosidad)
- tos persistente o crónica tal como ocurre con el tabaquismo, asma, bronquitis crónica o enfisema

Información del Medicamento (continuación)

Consulte a su médico o farmacéutico antes de utilizarlo si usted está tomando cualquier otro descongestionante o estimulante nasal oral.

No exceda la dosis recomendada.

- Suspense su uso y consulte a su médico si
 - se siente nervioso, mareado o con insomnio
 - los síntomas no mejoran en los siguientes 7 días o se acompañan de fiebre
 - la tos persiste por más de 7 días, regresa, o se presenta acompañada por fiebre, erupción o dolor de cabeza persistente. Estos podrían ser signos de una condición grave.

En caso de embarazo o lactancia, consulte a un profesional médico antes de su uso. **Manténgase fuera del alcance de los niños.** En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento de manera inmediata.

Indicaciones

- no tomar más de 6 dosis en un período de 24 horas
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad
- medir solo con la copa dosificadora incluida
- mantenga la copa dosificadora con este producto
- TSP = cucharadita

| edad | dosis |
|---|-------------------------------------|
| adultos y niños 12 años de edad y mayores | 2 cucharaditas (10 mL) cada 4 horas |
| niños menores de 12 años | no utilizar |

Información adicional

- cada cucharadita (5 mL) contiene: sodio 3 mg
- almacenar entre 15-30 °C (59-86 °F). No refrigerar.

Ingredientes inactivos


ácido cítrico anhidro, FD&C rojo #40, sabor, glicerina, mentol, propilenglicol, agua purificada, benzoato sódico, solución de sorbitol, sucralosa

¿Preguntas o comentarios?

1-800-560-5223 Lunes a Viernes 9am-4pm, Hora del Este

Distributed by / Distribuido por:
The Menper Group
11421 NW107th ST.
Miami, FL 33178, USA


MADE IN USA
HECHO EN EEUU



EXP:

LOT# NO./LOTE:

MUCOSAN MULTI-SYMPATOM CHERRY

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:53145-693

Route of Administration

ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|-------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 5 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 5 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 100 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| WATER (UNII: 059QF0KO0R) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| MENTHOL (UNII: L7T10EIP3A) | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:53145-693-04 | 1 in 1 CARTON | 11/01/2024 | |
| 1 | | 118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 11/01/2024 | |

Labeler - Menper Distributors, Inc. (101947166)

Registrant - Menper Distributors, Inc. (101947166)

Revised: 11/2024

Menper Distributors, Inc.