

**NUMB520- lidocaine liquid**  
**Clinical Resolution Laboratory, Inc.**

-----  
**Numb520 Spray**

***DRUG FACTS***

**Active Ingredient**

Lidocaine 5%

***Purpose***

Local Anesthetic

***Uses:***

Temporarily relieves pain, itching, or swelling associated with anorectal disorders.

***Warnings***

**(For external use only)**

**Do not use this product if**

- Pregnant or breastfeeding, ask a health professional before use
- Tamper-evident shrink band around the container is broken.

**When using this product**

- Do not exceed the recommended daily unless directed by a doctor.
- Certain persons can develop allergic reactions to ingredients in this product.
- Do not put this product into the rectum by using fingers or any medical device or applicator.
- In case of accidental overdose, contact a doctor or Poison Control Center immediately.

**Stop use and ask a doctor if**

The symptom being treated does not subside, or if redness, irritation, swelling, pain, or other symptoms develop or increase.

**Keep out of reach of children**

In case of accidental ingestion, seek medical attention immediately.

***Directions***

- Sensitivity test advised prior to use. Apply sparingly to affected area after thoroughly cleansing. Wait until anesthetic effect occurs (90 seconds). You may reapply to

continue numbing effect.

- Apply up to 6 times a day.
- Children under 12 year of age: consult a doctor.

### Other Information

- Keep away from direct sunlight or heat.
- Store in room temperature (59-86°F / 15-30°C).

### Inactive Ingredients

Water, Aloe Barbadensis (Aloe Vera) Leaf Juice, Propylene Glycol, Benzyl Alcohol, Epinephrine HCL, Citric Acid, Disodium EDTA, Cholesterol, Sodium Metabisulfite, Lecithin, Polysorbate 80, Phenoxyethanol

### Package Labeling:

	<table border="1"><tr><td colspan="2"><b>DRUG FACTS</b></td></tr><tr><td><b>Active Ingredient</b></td><td><b>Purpose</b></td></tr><tr><td>Lidocaine 5%</td><td>Local Anesthetic</td></tr><tr><td colspan="2"><b>Uses:</b> Temporarily relieves pain, itching, or swelling associated with anorectal disorders.</td></tr><tr><td colspan="2"><b>Warnings (For external use only)</b></td></tr><tr><td colspan="2"><b>Do not use this product if</b></td></tr><tr><td colspan="2">• Pregnant or breastfeeding, ask a health professional before use.</td></tr><tr><td colspan="2">• Tamper-evident shrink band around the container is broken.</td></tr><tr><td colspan="2"><b>When using this product</b></td></tr><tr><td colspan="2">• Do not exceed the recommended daily dosage unless directed by a doctor.</td></tr><tr><td colspan="2">• Certain persons can develop allergic reactions to ingredients in this product.</td></tr><tr><td colspan="2">• Do not put this product into the rectum by using fingers or any medical device or applicator.</td></tr><tr><td colspan="2">• In case of accidental overdose, contact a doctor or Poison Control Center immediately.</td></tr><tr><td colspan="2"><b>Stop use and ask a doctor if</b></td></tr><tr><td colspan="2">The symptom being treated does not subside, or if redness, irritation, swelling, pain, or other symptoms develop or increase.</td></tr><tr><td colspan="2"><b>Keep out of reach of children</b></td></tr><tr><td colspan="2">In case of accidental ingestion, seek medical attention immediately.</td></tr><tr><td colspan="2"><b>Directions</b></td></tr><tr><td colspan="2">• Sensitivity test advised prior to use. Apply sparingly to affected area after thoroughly cleansing. Wait until anesthetic effect occurs (90 seconds). You may reapply to continue numbing effect.</td></tr><tr><td colspan="2">• Apply up to 6 times a day.</td></tr><tr><td colspan="2">• Children under 12 years of age: consult a doctor.</td></tr><tr><td colspan="2"><b>Other Information</b></td></tr><tr><td colspan="2">• Keep away from direct sunlight or heat.</td></tr><tr><td colspan="2">• Store in room temperature (59-86°F / 15-30°C).</td></tr><tr><td colspan="2"><b>Inactive Ingredients</b></td></tr><tr><td colspan="2">Water, Aloe Barbadensis (Aloe Vera) Leaf Juice, Propylene Glycol, Benzyl Alcohol, Epinephrine HCl, Citric Acid, Disodium EDTA, Cholesterol, Sodium Metabisulfite, Lecithin, Polysorbate 80, Phenoxyethanol</td></tr></table>	<b>DRUG FACTS</b>		<b>Active Ingredient</b>	<b>Purpose</b>	Lidocaine 5%	Local Anesthetic	<b>Uses:</b> Temporarily relieves pain, itching, or swelling associated with anorectal disorders.		<b>Warnings (For external use only)</b>		<b>Do not use this product if</b>		• Pregnant or breastfeeding, ask a health professional before use.		• Tamper-evident shrink band around the container is broken.		<b>When using this product</b>		• Do not exceed the recommended daily dosage unless directed by a doctor.		• Certain persons can develop allergic reactions to ingredients in this product.		• Do not put this product into the rectum by using fingers or any medical device or applicator.		• In case of accidental overdose, contact a doctor or Poison Control Center immediately.		<b>Stop use and ask a doctor if</b>		The symptom being treated does not subside, or if redness, irritation, swelling, pain, or other symptoms develop or increase.		<b>Keep out of reach of children</b>		In case of accidental ingestion, seek medical attention immediately.		<b>Directions</b>		• Sensitivity test advised prior to use. Apply sparingly to affected area after thoroughly cleansing. Wait until anesthetic effect occurs (90 seconds). You may reapply to continue numbing effect.		• Apply up to 6 times a day.		• Children under 12 years of age: consult a doctor.		<b>Other Information</b>		• Keep away from direct sunlight or heat.		• Store in room temperature (59-86°F / 15-30°C).		<b>Inactive Ingredients</b>		Water, Aloe Barbadensis (Aloe Vera) Leaf Juice, Propylene Glycol, Benzyl Alcohol, Epinephrine HCl, Citric Acid, Disodium EDTA, Cholesterol, Sodium Metabisulfite, Lecithin, Polysorbate 80, Phenoxyethanol	
<b>DRUG FACTS</b>																																																					
<b>Active Ingredient</b>	<b>Purpose</b>																																																				
Lidocaine 5%	Local Anesthetic																																																				
<b>Uses:</b> Temporarily relieves pain, itching, or swelling associated with anorectal disorders.																																																					
<b>Warnings (For external use only)</b>																																																					
<b>Do not use this product if</b>																																																					
• Pregnant or breastfeeding, ask a health professional before use.																																																					
• Tamper-evident shrink band around the container is broken.																																																					
<b>When using this product</b>																																																					
• Do not exceed the recommended daily dosage unless directed by a doctor.																																																					
• Certain persons can develop allergic reactions to ingredients in this product.																																																					
• Do not put this product into the rectum by using fingers or any medical device or applicator.																																																					
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.																																																					
<b>Stop use and ask a doctor if</b>																																																					
The symptom being treated does not subside, or if redness, irritation, swelling, pain, or other symptoms develop or increase.																																																					
<b>Keep out of reach of children</b>																																																					
In case of accidental ingestion, seek medical attention immediately.																																																					
<b>Directions</b>																																																					
• Sensitivity test advised prior to use. Apply sparingly to affected area after thoroughly cleansing. Wait until anesthetic effect occurs (90 seconds). You may reapply to continue numbing effect.																																																					
• Apply up to 6 times a day.																																																					
• Children under 12 years of age: consult a doctor.																																																					
<b>Other Information</b>																																																					
• Keep away from direct sunlight or heat.																																																					
• Store in room temperature (59-86°F / 15-30°C).																																																					
<b>Inactive Ingredients</b>																																																					
Water, Aloe Barbadensis (Aloe Vera) Leaf Juice, Propylene Glycol, Benzyl Alcohol, Epinephrine HCl, Citric Acid, Disodium EDTA, Cholesterol, Sodium Metabisulfite, Lecithin, Polysorbate 80, Phenoxyethanol																																																					
 <p>Fast Acting Numbing Effect</p> <p><b>Topical Anesthetic Spray</b></p> <p>4.15 fl.oz e 123ml</p>	 <p>11005 83438 4</p> <p>@EbanelSkincare</p>  <p>Manufactured for Ebanel Laboratories, Inc. www.Ebanel.com • Made in USA</p>																																																				

Ebanel™

# NUMB520 SPRAY



Fast Acting Numbing Effect

## Topical Anesthetic Spray

8.5 fl.oz e 251ml

### DRUG FACTS

Active Ingredient	Purpose
Lidocaine 5%	Local Anesthetic

**Uses:** Temporarily relieves pain, itching, or swelling associated with anorectal disorders.

#### Warnings (For external use only)

##### Do not use this product if

- Pregnant or breastfeeding, ask a health professional before use.
- Tamper-evident shrink band around the container is broken.

##### When using this product

- Do not exceed the recommended daily dosage unless directed by a doctor.
- Certain persons can develop allergic reactions to ingredients in this product.
- Do not put this product into the rectum by using fingers or any medical device or applicator.
- In case of accidental overdose, contact a doctor or Poison Control Center immediately.

##### Stop use and ask a doctor if

The symptom being treated does not subside, or if redness, irritation, swelling, pain, or other symptoms develop or increase.

##### Keep out of reach of children

In case of accidental ingestion, seek medical attention immediately.

##### Directions

- Sensitivity test advised prior to use. Apply sparingly to affected area after thoroughly cleansing. Wait until anesthetic effect occurs (90 seconds). You may reapply to continue numbing effect.
- Apply up to 6 times a day.
- Children under 12 years of age: consult a doctor.

##### Other Information

- Keep away from direct sunlight or heat.
- Store in room temperature (59-86°F / 15-30°C).

##### Inactive Ingredients

Water, Aloe Barbadensis (Aloe Vera) Leaf Juice, Propylene Glycol, Benzyl Alcohol, Epinephrine HCl, Citric Acid, Disodium EDTA, Cholesterol, Sodium Metabisulfite, Lecithin, Polysorbate 80, Phenoxyethanol



@EbanelSkincare



Manufactured for Ebanel Laboratories, Inc.  
www.Ebanel.com • Made in USA

## NUMB520

lidocaine liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63742-006
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	50 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
CHOLESTEROL (UNII: 97C5T2UQ7J)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>EPINEPHRINE HYDROCHLORIDE</b> (UNII: WBB047OO38)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63742-006-00	123 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/15/2018	
2	NDC:63742-006-01	251 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/15/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M015	05/15/2018	

**Labeler** - Clinical Resolution Laboratory, Inc. (825047942)

Revised: 12/2023

Clinical Resolution Laboratory, Inc.