

ASPIRIN LOW DOSE- aspirin tablet, chewable
Better Living Brands, LLC

Signature Care 44-218

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

Signature™

care

Quality Guaranteed

COMPARE TO

Bayer® Chewable Low Dose Aspirin
active ingredient†

NDC 21130-918-07

Low Dose

Aspirin 81 mg

Chewable Tablets

Pain Reliever (**NSAID**)

• Aspirin regimen**

ORANGE

FLAVOR

**Chew or crush
tablets completely
before swallowing.**

Actual Size

36 TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

****Talk to your doctor or other healthcare provider
before using this product for your heart.**

†This product is not manufactured or distributed by
Bayer AG, distributors of Bayer® Chewable Low Dose
Aspirin. 50844 REV0122B21807

**DISTRIBUTED BY BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0009
1-888-723-3929
www.betterlivingbrandsLLC.com**

**QUALITY & SATISFACTION 100% GUARANTEED
OR YOUR MONEY BACK**

1This product is not manufactured or distributed by Bayer AG, distributors of Bayer® Chewable Low Dose Aspirin. 50844 REV0122B21807

DISTRIBUTED BY BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0009
1-888-723-3929
www.betterlivingbrandsLLC.com
QUALITY & SATISFACTION 100% GUARANTEED
OR YOUR MONEY BACK



Scan here for more information



COMPARE TO Bayer® Chewable Low Dose Aspirin active ingredient¹

NDC 21130-918-07

Low Dose Aspirin 81 mg

Chewable Tablets Pain Reliever (NSAID)

Chew or crush tablets completely before swallowing.

Actual Size

36 TABLETS

• Aspirin regimen**




ORANGE FLAVOR

**Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

RD 19094



3 121130 77942 1

S1714

**No Print / No Varnish
Lot no. & Exp. date**

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?
1-800-426-9391

Drug Facts

Active ingredient (in each chewable tablet)
Aspirin 81 mg (NSAID) Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Sign use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur

These could be signs of a serious condition.

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Sign use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur

These could be signs of a serious condition.

Signature Care 44-218A

ASPIRIN LOW DOSE

aspirin tablet, chewable

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-918
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange (light)	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	44;218
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-918-07	1 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	

Labeler - Better Living Brands, LLC (009137209)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(21130-918)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(21130-918)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(21130-918)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(21130-918)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(21130-918)

Revised: 5/2024

Better Living Brands, LLC