

**NAPROXEN SODIUM- naproxen sodium tablet, coated**  
**Harris Teeter**

-----  
**HTE - 1144 - 2019-1010**

***Drug Facts***

**Active ingredient (in each tablet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID <sup>1</sup>)

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

**Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

adults and children 12 years and older	<ul style="list-style-type: none"> <li>▪ take 1 tablet every 8 to 12 hours while symptoms last</li> <li>▪ for the first dose you may take 2 tablets within the first hour</li> <li>▪ do not exceed 2 tablets in any 8- to 12-hour period</li> <li>▪ do not exceed 3 tablets in a 24-hour period</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>▪ ask a doctor</li> </ul>

### Other information

- **each tablet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information

### Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

### PRINCIPAL DISPLAY PANEL

Harris Teeter


Naproxen Sodium

Compare to the Active Ingredient in Aleve(R)

Naproxen Sodium Tablets, 220 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

24 Tablets



**NAPROXEN SODIUM**  
naproxen sodium tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-144
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:5Y76R9ATQ)		NAPROXEN SODIUM	220 mg

Inactive Ingredients		Strength
Ingredient Name		

<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10 mm
<b>Flavor</b>		<b>Imprint Code</b>	I3
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72036-144-01	1 in 1 CARTON	07/17/2012	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:72036-144-02	1 in 1 CARTON	12/01/2011	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:72036-144-03	1 in 1 CARTON	12/01/2011	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	12/01/2011	

**Labeler** - Harris Teeter (047279351)