

MOTION SICKNESS RELIEF- meclizine hcl tablet
Walgreen Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens 44-403-Motion sickness

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity
- to treat motion sickness, take at first signs of symptoms
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

Questions or comments?

1-800-426-9391

Principal display panel

Compare to the active ingredient in
Dramamine® All Day Less Drowsy††

NDC 0363-4030-21

Motion Sickness Relief

MECLIZINE HCl TABLETS, 25 mg / ANTIEMETIC

All Day Relief Less Drowsy Formula

- Helps prevent nausea, dizziness & vomiting due to motion sickness
- All-day relief of motion sickness

16 TABLETS

ACTUAL SIZE

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Medtech Products Inc.,

owner of the registered trademark
Dramamine® All Day Less Drowsy.

50844 ORG021940321

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED

walgreens.com

©2021 Walgreen Co.



Walgreens 44-403

MOTION SICKNESS RELIEF

meclizine hcl tablet

W30RG1021-F2

Drug Facts (continued)
(Questions or comments? 1-800-426-9391)

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® All Day Less Drowsy. 50844 ORG021940321

Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet) Purpose
 Meclizine HCl 25 mg, Antiemetic

Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

Warnings
 Do not use for children under 12 years of age unless directed by a doctor.
 Ask a doctor before use if you have ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ breathing problem such as emphysema or chronic bronchitis
 Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
When using this product
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ drowsiness may occur ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity
 ■ to treat motion sickness, take at first signs of symptoms
 ■ adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ protect from heat and humidity
 ■ see end flap for expiration date and lot number

Inactive ingredients corn starch, D & C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Do not print Lot & Exp

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com
 ©2021 Walgreen Co.

ITEM 501197 W00000-0000-0
 3 1191704695 2



B-2201-403-21-HR
 ORG021940321

NDC 0363-4030-21



Motion Sickness Relief
 MECLIZINE HCl TABLETS, 25 mg / ANTIEMETIC

All Day Relief Less Drowsy Formula

- Helps prevent nausea, dizziness & vomiting due to motion sickness
- All-day relief of motion sickness

16 TABLETS



ACTUAL SIZE

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-4030
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4030-21	2 in 1 CARTON	10/21/2021	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	10/21/2021	

Labeler - Walgreen Company (008965063)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-4030)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-4030) , pack(0363-4030)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-4030)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-4030)

Revised: 10/2022

Walgreen Company