

A1043-17 SINGLE SHOT EPIDURAL 17G TUOHY- anesthesia conduction kit
Smiths Medical ASD, Inc.

Lidocaine Rx only
Hydrochloride
Injection, USP

AQUEOUS SOLUTIONS FOR
INFILTRATION AND NERVE BLOCK

Ampul

Plastic Multiple-dose Fliptop Vial

Glass Teartop Vial


Lidocaine Hydrochloride Injection, USP is a sterile, nonpyrogenic solution of lidocaine hydrochloride in water for injection for parenteral administration in various concentrations with characteristics as follows:

Concentration	0.5%	1%	1.5%	2%
mg/mL lidocaine HCl (anhyd.)	5	10	15	20
mg/mL sodium chloride	8	7	6.5	6

Multiple-dose vials contain 0.1% of methylparaben added as preservative. May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. The pH is 6.5 (5.0 to 7.0). See HOW SUPPLIED section for various sizes and strengths.

Lidocaine is a local anesthetic of the amide type.

Lidocaine Hydrochloride, USP is chemically designated 2-(diethylamino)-N-(2,6-dimethylphenyl)-acetamide monohydrochloride monohydrate, a white powder freely soluble in water. The molecular weight is 288.82. It has the following structural formula:



The semi-rigid vial used for the plastic vials is fabricated from a specially formulated polyolefin. It is a copolymer of ethylene and propylene. The safety of the plastic has been confirmed by tests in animals according to USP biological standards for plastic containers. The container requires no vapor barrier to maintain the proper drug concentration.

Mechanism of action: Lidocaine stabilizes the neuronal membrane by inhibiting the ionic fluxes required for the initiation and conduction of impulses, thereby effecting local anesthetic action.

Hemodynamics: Excessive blood levels may cause changes in cardiac output, total peripheral resistance, and mean arterial pressure. With central neural blockade these changes may be attributable to block of autonomic fibers, a direct depressant effect of the local anesthetic agent on various

components of the cardiovascular system and/or the beta-adrenergic receptor stimulating action of epinephrine when present. The net effect is normally a modest hypotension when the recommended dosages are not exceeded.

Pharmacokinetics and metabolism: Information derived from diverse formulations, concentrations and usages reveals that lidocaine is completely absorbed following parenteral administration, its rate of absorption depending, for example, upon various factors such as the site of administration and the presence or absence of a vasoconstrictor agent. Except for intravascular administration, the highest blood levels are obtained following intercostal nerve block and the lowest after subcutaneous administration.

The plasma binding of lidocaine is dependent on drug concentration, and the fraction bound decreases with increasing concentration. At concentrations of 1 to 4 mcg of free base per mL, 60 to 80 percent of lidocaine is protein bound. Binding is also dependent on the plasma concentration of the alpha-1-acid glycoprotein.

Lidocaine crosses the blood-brain and placental barriers, presumably by passive diffusion.

Lidocaine is metabolized rapidly by the liver, and metabolites and unchanged drug are excreted by the kidneys. Biotransformation includes oxidative N-dealkylation, ring hydroxylation, cleavage of the amide linkage, and conjugation. N-dealkylation, a major pathway of biotransformation, yields the metabolites monoethylglycinexylidide and glycinexylidide. The pharmacological/toxicological actions of these metabolites are similar to, but less potent than, those of lidocaine. Approximately 90% of lidocaine administered is excreted in the form of various metabolites, and less than 10% is excreted unchanged. The primary metabolite in urine is a conjugate of 4-hydroxy-2, 6-dimethylaniline.

The elimination half-life of lidocaine following an intravenous bolus injection is typically 1.5 to 2.0 hours. Because of the rapid rate at which lidocaine is metabolized, any condition that affects liver function may alter lidocaine kinetics. The half-life may be prolonged two-fold or more in patients with liver dysfunction. Renal dysfunction does not affect lidocaine kinetics but may increase the accumulation of metabolites.

Factors such as acidosis and the use of CNS stimulants and depressants affect the CNS levels of lidocaine required to produce overt systemic effects. Objective adverse manifestations become increasingly apparent with increasing venous plasma levels above 6.0 mcg free base per mL. In the rhesus monkey arterial blood levels of 18-21 mcg/mL have been shown to be threshold for convulsive activity.

Lidocaine Hydrochloride Injection, USP is indicated for production of local or regional anesthesia by infiltration techniques such as percutaneous injection and intravenous regional anesthesia by peripheral nerve block techniques such as brachial plexus and intercostal and by central neural techniques such as lumbar and caudal epidural blocks, when the accepted procedures for these techniques as described in standard textbooks are observed.

Lidocaine is contraindicated in patients with a known history of hypersensitivity to local anesthetics of the amide type.

LIDOCAINE HYDROCHLORIDE INJECTION, FOR INFILTRATION AND NERVE BLOCK, SHOULD BE EMPLOYED ONLY BY CLINICIANS WHO ARE WELL VERSED IN DIAGNOSIS AND MANAGEMENT OF DOSE-RELATED TOXICITY AND OTHER ACUTE EMERGENCIES THAT MIGHT ARISE FROM THE BLOCK TO BE EMPLOYED AND THEN ONLY AFTER ENSURING THE IMMEDIATE AVAILABILITY OF OXYGEN, OTHER RESUSCITATIVE DRUGS, CARDIOPULMONARY EQUIPMENT, AND THE PERSONNEL NEEDED FOR PROPER MANAGEMENT OF TOXIC REACTIONS AND RELATED EMERGENCIES (See also ADVERSE REACTIONS and PRECAUTIONS). DELAY IN PROPER MANAGEMENT OF DOSE-RELATED TOXICITY, UNDERVENTILATION FROM ANY CAUSE AND/OR ALTERED SENSITIVITY MAY LEAD TO THE DEVELOPMENT OF ACIDOSIS, CARDIAC ARREST AND, POSSIBLY, DEATH.

Intra-articular infusions of local anesthetics following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. The majority of reported cases of chondrolysis have involved the shoulder joint; cases of gleno-humeral chondrolysis have been described in pediatric and adult patients following intra-articular infusions of local anesthetics with and without epinephrine for periods of 48 to 72 hours. There is insufficient information to determine whether shorter infusion periods are not associated with these findings. The time of onset of symptoms, such as joint pain, stiffness and loss of motion can be variable, but may begin as early as the 2nd month after surgery. Currently, there is no effective treatment for chondrolysis; patients who experienced chondrolysis have required additional diagnostic and therapeutic procedures and some required arthroplasty or shoulder replacement.

To avoid intravascular injection, aspiration should be performed before the local anesthetic solution is injected. The needle must be repositioned until no return of blood can be elicited by aspiration. Note, however, that the absence of blood in the syringe does not guarantee that intravascular injection has been avoided.

Local anesthetic solutions containing antimicrobial preservatives (e.g., methylparaben) should not be used for epidural or spinal anesthesia because the safety of these agents has not been established with regard to intrathecal injection, either intentional or accidental.

General: The safety and effectiveness of lidocaine depend on proper dosage, correct technique, adequate precautions, and readiness for emergencies. Standard textbooks should be consulted for specific techniques and precautions for various regional anesthetic procedures.

Resuscitative equipment, oxygen, and other resuscitative drugs should be available for immediate use. (See WARNINGS and ADVERSE REACTIONS). The lowest dosage that results in effective anesthesia should be used to avoid high plasma levels and serious adverse effects. Syringe aspirations should also be performed before and during each supplemental injection when using indwelling catheter techniques. During the administration of epidural anesthesia, it is recommended that a test dose be administered initially and that the patient be monitored for central nervous system toxicity and cardiovascular toxicity, as well as for signs of unintended intrathecal administration before proceeding. When clinical conditions permit, consideration should be given to employing local anesthetic solutions that contain epinephrine for the test dose because circulatory changes compatible with epinephrine may also serve as a warning sign of unintended intravascular injection. An intravascular injection is still possible even if aspirations for blood are negative. Repeated doses of lidocaine may cause significant increases in blood levels with each repeated dose because of slow accumulation of the drug or its metabolites. Tolerance to elevated blood levels varies with the status of the patient. Debilitated, elderly patients, acutely ill patients and children should be given reduced doses commensurate with their age and physical condition. Lidocaine should also be used with caution in patients with severe shock or heart block. Lumbar and caudal epidural anesthesia should be used with extreme caution in persons with the following conditions: existing neurological disease, spinal deformities, septicemia and severe hypertension.

Local anesthetic solutions containing a vasoconstrictor should be used cautiously and in carefully circumscribed quantities in areas of the body supplied by end arteries or having otherwise compromised blood supply. Patients with peripheral vascular disease and those with hypertensive vascular disease may exhibit exaggerated vasoconstrictor response. Ischemic injury or necrosis may result. Preparations containing a vasoconstrictor should be used with caution in patients during or following the administration of potent general anesthetic agents, since cardiac arrhythmias may occur under such conditions.

Careful and constant monitoring of cardiovascular and respiratory (adequacy of ventilation) vital signs and the patient's state of consciousness should be accomplished after each local anesthetic injection. It should be kept in mind at such times that restlessness, anxiety, tinnitus, dizziness, blurred vision, tremors, depression or drowsiness may be early warning signs of central nervous system toxicity.

Since amide-type local anesthetics are metabolized by the liver, lidocaine should be used with caution in patients with hepatic disease. Patients with severe hepatic disease, because of their inability to metabolize local anesthetics normally, are at greater risk of developing toxic plasma concentrations. Lidocaine should also be used with caution in patients with impaired cardiovascular function since they may be less able to compensate for functional changes associated with the prolongation of A-V conduction produced by these drugs. Many drugs used during the conduct of anesthesia are considered potential triggering agents for familial malignant hyperthermia. Since it is not known whether amide-type local anesthetics may trigger this reaction and since the need for supplemental general anesthesia cannot be predicted in advance, it is suggested that a standard protocol for the management of malignant hyperthermia should be available. Early unexplained signs of tachycardia, tachypnea, labile blood pressure and metabolic acidosis may precede temperature elevation. Successful outcome is dependent on early diagnosis, prompt discontinuance of the suspect triggering agent(s) and institution of treatment, including oxygen therapy, indicated supportive measures and dantrolene (consult dantrolene sodium intravenous package insert before using).

Proper tourniquet technique, as described in publications and standard textbooks, is essential in the performance of intravenous regional anesthesia. Solutions containing epinephrine or other vasoconstrictors should not be used for this technique.

Lidocaine should be used with caution in persons with known drug sensitivities. Patients allergic to para-aminobenzoic acid derivatives (procaine, tetracaine, benzocaine, etc.) have not shown cross sensitivity to lidocaine.

Use in the Head and Neck Area: Small doses of local anesthetics injected into the head and neck area, including retrobulbar, dental and stellate ganglion blocks, may produce adverse reactions similar to systemic toxicity seen with unintentional intravascular injections of larger doses. Confusion, convulsions, respiratory depression and/or respiratory arrest and cardiovascular stimulation or depression have been reported. These reactions may be due to intra-arterial injections of the local anesthetic with retrograde flow to the cerebral circulation. Patients receiving these blocks should have their circulation and respiration monitored and be constantly observed. Resuscitative equipment and personnel for treating adverse reactions should be immediately available. Dosage recommendations should not be exceeded. (See DOSAGE AND ADMINISTRATION).

Information for Patients: When appropriate, patients should be informed in advance that they may experience temporary loss of sensation and motor activity, usually in the lower half of the body following proper administration of epidural anesthesia.

Clinically Significant Drug Interactions: The administration of local anesthetic solutions containing epinephrine or norepinephrine to patients receiving monoamine oxidase inhibitors or tricyclic antidepressants may produce severe prolonged hypertension.

Phenothiazines and butyrophenones may reduce or reverse the pressor effect of epinephrine.

Concurrent use of these agents should generally be avoided. In situations when concurrent therapy is necessary, careful patient monitoring is essential.

Concurrent administration of vasopressor drugs (for the treatment of hypotension related to obstetric blocks) and ergot-type oxytoxic drugs may cause severe persistent hypertension or cerebrovascular accidents.

Drug Laboratory Test Interactions: The intramuscular injection of lidocaine may result in an increase in creatine phosphokinase levels. Thus, the use of this enzyme determination without isoenzyme separation as a diagnostic test for the presence of acute myocardial infarction may be compromised by the intramuscular injection of lidocaine.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Studies of lidocaine in animals to evaluate the carcinogenic and mutagenic potential or the effect on fertility have not been conducted.

Pregnancy: Teratogenic Effects. Pregnancy Category B. Reproduction studies have been performed in

rats at doses up to 6.6 times the human dose and have revealed no evidence of harm to the fetus caused by lidocaine. There are, however, no adequate and well-controlled studies in pregnant women. Animal reproduction studies are not always predictive of human response. General consideration should be given to this fact before administering lidocaine to women of childbearing potential, especially during early pregnancy when maximum organogenesis takes place.

Labor and Delivery: Local anesthetics rapidly cross the placenta and when used for epidural, paracervical, pudendal or caudal block anesthesia, can cause varying degrees of maternal, fetal and neonatal toxicity (See CLINICAL PHARMACOLOGY—Pharmacokinetics). The potential for toxicity depends upon the procedure performed, the type and amount of drug used, and the technique of drug administration. Adverse reactions in the parturient, fetus and neonate involve alterations of the central nervous system peripheral vascular tone and cardiac function.

Maternal hypotension has resulted from regional anesthesia. Local anesthetics produce vasodilation by blocking sympathetic nerves. Elevating the patient's legs and positioning her on her left side will help prevent decreases in blood pressure. The fetal heart rate also should be monitored continuously, and electronic fetal monitoring is highly advisable.

Epidural, spinal, paracervical, or pudendal anesthesia may alter the forces of parturition through changes in uterine contractility or maternal expulsive efforts. In one study, paracervical block anesthesia was associated with a decrease in the mean duration of first stage labor and facilitation of cervical dilation. However, spinal and epidural anesthesia have also been reported to prolong the second stage of labor by removing the parturient's reflex urge to bear down or by interfering with motor function. The use of obstetrical anesthesia may increase the need for forceps assistance.

The use of some local anesthetic drug products during labor and delivery may be followed by diminished muscle strength and tone for the first day or two of life. The long-term significance of these observations is unknown. Fetal bradycardia may occur in 20 to 30 percent of patients receiving paracervical nerve block anesthesia with the amide-type local anesthetics and may be associated with fetal acidosis. Fetal heart rate should always be monitored during paracervical anesthesia. The physician should weigh the possible advantages against risks when considering paracervical block in prematurity, toxemia of pregnancy and fetal distress. Careful adherence to recommended dosage is of the utmost importance in obstetrical paracervical block. Failure to achieve adequate analgesia with recommended doses should arouse suspicion of intravascular or fetal intracranial injection. Cases compatible with unintended fetal intracranial injection of local anesthetic solution have been reported following intended paracervical or pudendal block or both. Babies so affected present with unexplained neonatal depression at birth, which correlates with high local anesthetic serum levels, and often manifest seizures within six hours. Prompt use of supportive measures combined with forced urinary excretion of the local anesthetic has been used successfully to manage this complication.

Case reports of maternal convulsions and cardiovascular collapse following use of some local anesthetics for paracervical block in early pregnancy (as anesthesia for elective abortion) suggest that systemic absorption under these circumstances may be rapid. The recommended maximum dose of each drug should not be exceeded. Injection should be made slowly and with frequent aspiration. Allow a 5-minute interval between sides.

Nursing Mothers: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when lidocaine is administered to a nursing woman.

Pediatric Use: Dosages in pediatric patients should be reduced, commensurate with age, body weight and physical condition. See DOSAGE AND ADMINISTRATION.

Systemic: Adverse experiences following the administration of lidocaine are similar in nature to those observed with other amide local anesthetic agents. These adverse experiences are, in general, dose-related and may result from high plasma levels caused by excessive dosage, rapid absorption or inadvertent intravascular injection, or may result from a hypersensitivity, idiosyncrasy or diminished tolerance on the part of the patient. Serious adverse experiences are generally systemic in nature. The

following types are those most commonly reported:

Central Nervous System: CNS manifestations are excitatory and/or depressant and may be characterized by lightheadedness, nervousness, apprehension, euphoria, confusion, dizziness, drowsiness, tinnitus, blurred or double vision, vomiting, sensations of heat, cold or numbness, twitching, tremors, convulsions, unconsciousness, respiratory depression and arrest. The excitatory manifestations may be very brief or may not occur at all, in which case the first manifestation of toxicity may be drowsiness merging into unconsciousness and respiratory arrest.

Drowsiness following the administration of lidocaine is usually an early sign of a high blood level of the drug and may occur as a consequence of rapid absorption.

Cardiovascular System: Cardiovascular manifestations are usually depressant and are characterized by bradycardia, hypotension, and cardiovascular collapse, which may lead to cardiac arrest.

Allergic: Allergic reactions are characterized by cutaneous lesions, urticaria, edema or anaphylactoid reactions. Allergic reactions may occur as a result of sensitivity either to local anesthetic agents or to the methylparaben used as a preservative in multiple dose vials. Allergic reactions as a result of sensitivity to lidocaine are extremely rare and, if they occur, should be managed by conventional means. The detection of sensitivity by skin testing is of doubtful value.

Neurologic: The incidences of adverse reactions associated with the use of local anesthetics may be related to the total dose of local anesthetic administered and are also dependent upon the particular drug used, the route of administration and the physical status of the patient. In a prospective review of 10,440 patients who received lidocaine for spinal anesthesia, the incidences of adverse reactions were reported to be about 3 percent each for positional headaches, hypotension and backache; 2 percent for shivering; and less than 1 percent each for peripheral nerve symptoms, nausea, respiratory inadequacy and double vision. Many of these observations may be related to local anesthetic techniques, with or without a contribution from the local anesthetic.

In the practice of caudal or lumbar epidural block, occasional unintentional penetration of the subarachnoid space by the catheter may occur. Subsequent adverse effects may depend partially on the amount of drug administered subdurally.

These may include spinal block of varying magnitude (including total spinal block), hypotension secondary to spinal block, loss of bladder and bowel control, and loss of perineal sensation and sexual function. Persistent motor, sensory and/or autonomic (sphincter control) deficit of some lower spinal segments with slow recovery (several months) or incomplete recovery have been reported in rare instances when caudal or lumbar epidural block has been attempted. Backache and headache have also been noted following use of these anesthetic procedures.

There have been reported cases of permanent injury to extraocular muscles requiring surgical repair following retrobulbar administration.

Acute emergencies from local anesthetics are generally related to high plasma levels encountered during therapeutic use of local anesthetics or to unintended subarachnoid injection of local anesthetic solution (see ADVERSE REACTIONS, WARNINGS and PRECAUTIONS).

Management of Local Anesthetic Emergencies: The first consideration is prevention, best accomplished by careful monitoring of cardiovascular and respiratory vital signs and the patient's state of consciousness after each local anesthetic injection. At the first sign of change, oxygen should be administered.

The first step in the management of convulsions, as well as underventilation or apnea due to unintended subarachnoid injection of drug solution, consists of immediate attention to the maintenance of a patent airway and assisted or controlled ventilation with oxygen and a delivery system capable of permitting immediate positive airway pressure by mask. Immediately after the institution of these ventilatory measures, the adequacy of the circulation should be evaluated, keeping in mind that drugs used to treat convulsions sometimes depress the circulation when administered intravenously. Should convulsions

persist despite adequate respiratory support, and if the status of the circulation permits, small increments of an ultra-short acting barbiturate (such as thiopental or thiamylal) or a benzodiazepine (such as diazepam) may be administered intravenously. The clinician should be familiar, prior to use of local anesthetics, with these anticonvulsant drugs. Supportive treatment of circulatory depression may require administration of intravenous fluids and, when appropriate, a vasopressor as directed by the clinical situation (e.g., ephedrine).

If not treated immediately, both convulsions and cardiovascular depression can result in hypoxia, acidosis, bradycardia, arrhythmias and cardiac arrest. Underventilation or apnea due to unintentional subarachnoid injection of local anesthetic solution may produce these same signs and also lead to cardiac arrest if ventilatory support is not instituted. If cardiac arrest should occur standard cardiopulmonary resuscitative measures should be instituted.

Endotracheal intubation, employing drugs and techniques familiar to the clinician, may be indicated, after initial administration of oxygen by mask, if difficulty is encountered in the maintenance of a patent airway or if prolonged ventilatory support (assisted or controlled) is indicated.

Dialysis is of negligible value in the treatment of acute overdosage with lidocaine.

The oral LD50 of lidocaine HCl in non-fasted female rats is 459 (346–773) mg/kg (as the salt) and 214 (159–324) mg/kg (as the salt) in fasted female rats.

Table 1 (Recommended Dosages) summarizes the recommended volumes and concentrations of Lidocaine Hydrochloride Injection, USP for various types of anesthetic procedures. The dosages suggested in this table are for normal healthy adults and refer to the use of epinephrine-free solutions. When larger volumes are required only solutions containing epinephrine should be used, except in those cases where vasopressor drugs may be contraindicated.

There have been adverse event reports of chondrolysis in patients receiving intra-articular infusions of local anesthetics following arthroscopic and other surgical procedures. Lidocaine is not approved for this use (see WARNINGS and DOSAGE AND ADMINISTRATION).

These recommended doses serve only as a guide to the amount of anesthetic required for most routine procedures. The actual volumes and concentrations to be used depend on a number of factors such as type and extent of surgical procedure, depth of anesthesia and degree of muscular relaxation required, duration of anesthesia required, and the physical condition of the patient. In all cases the lowest concentration and smallest dose that will produce the desired result should be given. Dosages should be reduced for children and for elderly and debilitated patients and patients with cardiac and/or liver disease.

The onset of anesthesia, the duration of anesthesia and the degree of muscular relaxation are proportional to the volume and concentration (i.e., total dose) of local anesthetic used. Thus, an increase in volume and concentration of Lidocaine Hydrochloride Injection will decrease the onset of anesthesia, prolong the duration of anesthesia, provide a greater degree of muscular relaxation and increase the segmental spread of anesthesia. However, increasing the volume and concentration of Lidocaine Hydrochloride Injection may result in a more profound fall in blood pressure when used in epidural anesthesia. Although the incidence of side effects with lidocaine is quite low, caution should be exercised when employing large volumes and concentrations, since the incidence of side effects is directly proportional to the total dose of local anesthetic agent injected.

For intravenous regional anesthesia, only the 50 mL single-dose vial containing 0.5% Lidocaine Hydrochloride Injection, USP should be used.

Epidural Anesthesia

For epidural anesthesia, only the following available specific products of Lidocaine Hydrochloride Injection by Hospira are recommended:

1%. 30 mL single-dose teartop vials

1.5%..... 20 mL single-dose ampuls
2%..... 10 mL single-dose ampuls

Although these solutions are intended specifically for epidural anesthesia, they may also be used for infiltration and peripheral nerve block provided they are employed as single dose units. These solutions contain no bacteriostatic agent. In epidural anesthesia, the dosage varies with the number of dermatomes to be anesthetized (generally 2–3 mL of the indicated concentration per dermatome).

Caudal and Lumbar Epidural Block: As a precaution against the adverse experiences sometimes observed following unintentional penetration of the subarachnoid space, a test dose such as 2–3 mL of 1.5% lidocaine hydrochloride should be administered at least 5 minutes prior to injecting the total volume required for a lumbar or caudal epidural block. The test dose should be repeated if the patient is moved in a manner that may have displaced the catheter. Epinephrine, if contained in the test dose (10–15 mcg have been suggested), may serve as a warning of unintentional intravascular injection. If injected into a blood vessel, this amount of epinephrine is likely to produce a transient "epinephrine response" within 45 seconds, consisting of an increase in heart rate and systolic blood pressure, circumoral pallor, palpitations and nervousness in the unsedated patient. The sedated patient may exhibit only a pulse rate increase of 20 or more beats per minute for 15 or more seconds. Patients on beta-blockers may not manifest changes in heart rate, but blood pressure monitoring can detect an evanescent rise in systolic blood pressure. Adequate time should be allowed for onset of anesthesia after administration of each test dose. The rapid injection of a large volume of Lidocaine Hydrochloride Injection through the catheter should be avoided, and, when feasible, fractional doses should be administered.

In the event of the known injection of a large volume of local anesthetic solutions into the subarachnoid space, after suitable resuscitation and if the catheter is in place, consider attempting the recovery of drug by draining a moderate amount of cerebrospinal fluid (such as 10 mL) through the epidural catheter.

Maximum Recommended Dosages

NOTE: The products accompanying this insert do not contain epinephrine.

Adults: For normal healthy adults, the individual maximum recommended dose of lidocaine HCl with epinephrine should not exceed 7 mg/kg (3.5 mg/lb) of body weight and in general it is recommended that the maximum total dose not exceed 500 mg. When used without epinephrine, the maximum individual dose should not exceed 4.5 mg/kg (2 mg/lb) of body weight and in general it is recommended that the maximum total dose does not exceed 300 mg. For continuous epidural or caudal anesthesia, the maximum recommended dosage should not be administered at intervals of less than 90 minutes. When continuous lumbar or caudal epidural anesthesia is used for non-obstetrical procedures, more drug may be administered if required to produce adequate anesthesia.

The maximum recommended dose per 90 minute period of lidocaine hydrochloride for paracervical block in obstetrical patients and non-obstetrical patients is 200 mg total. One-half of the total dose is usually administered to each side. Inject slowly five minutes between sides. (See also discussion of paracervical block in PRECAUTIONS).

For intravenous regional anesthesia, the dose administered should not exceed 4 mg/kg in adults.

Children: It is difficult to recommend a maximum dose of any drug for children, since this varies as a function of age and weight. For children over 3 years of age who have a normal lean body mass and normal body development, the maximum dose is determined by the child's age and weight. For example, in a child of 5 years weighing 50 lbs., the dose of lidocaine HCl should not exceed 75 — 100 mg (1.5 — 2 mg/lb). The use of even more dilute solutions (i.e., 0.25 — 0.5%) and total dosages not to exceed 3 mg/kg (1.4 mg/lb) are recommended for induction of intravenous regional anesthesia in children.

In order to guard against systemic toxicity, the lowest effective concentration and lowest effective dose should be used at all times. In some cases it will be necessary to dilute available concentrations with 0.9% sodium chloride injection in order to obtain the required final concentration.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever the solution and container permit. Solutions that are discolored and/or contain particulate matter should not be used.

Table 1
Recommended Dosages of Lidocaine Hydrochloride Injection, USP for Various Anesthetic Procedures in Normal Healthy Adults

Procedure	Lidocaine Hydrochloride Injection, USP (without Epinephrine)		
	Conc. (%)	Vol. (mL)	Total Dose (mg)
Infiltration			
Percutaneous	0.5 or 1.0	1–60	5–300
Intravenous Regional	0.5	10–60	50–300
Peripheral Nerve Blocks, e.g.			
Brachial	1.5	15–20	225–300
Dental	2.0	1–5	20–100
Intercostal	1.0	3	30
Paravertebral	1.0	3–5	30–50
Pudendal (each side)	1.0	10	100
Paracervical			
Obstetrical Analgesia (each side)	1.0	10	100
Sympathetic Nerve Blocks, e.g.			
Cervical (stellate ganglion)	1.0	5	50
Lumbar	1.0	5–10	50–100
Central Neural Blocks			
Epidural*			
Thoracic	1.0	20–30	200–300
Lumbar			
Analgesia	1.0	25–30	250–300
Anesthesia	1.5	15–20	225–300
	2.0	10–15	200–300
Caudal			
Obstetrical Analgesia	1.0	20–30	200–300
Surgical Anesthesia	1.5	15–20	225–300

*Dose determined by number of dermatomes to be anesthetized (2 to 3 mL/dermatome).

THE ABOVE SUGGESTED CONCENTRATIONS AND VOLUMES SERVE ONLY AS A GUIDE. OTHER VOLUMES AND CONCENTRATIONS MAY BE USED PROVIDED THE TOTAL MAXIMUM RECOMMENDED DOSE IS NOT EXCEEDED.

Sterilization, Storage and Technical Procedures: Disinfecting agents containing heavy metals, which cause release of respective ions (mercury, zinc, copper, etc.) should not be used for skin or mucous membrane disinfection as they have been related to incidence of swelling and edema. When chemical disinfection of multi-dose vials is desired, either isopropyl alcohol (91%) or 70% ethyl alcohol is recommended. Many commercially available brands of rubbing alcohol, as well as solutions of ethyl alcohol not of USP grade, contain denaturants which are injurious to rubber and, therefore, are not to be used. It is recommended that chemical disinfection be accomplished by wiping the vial stopper

thoroughly with cotton or gauze that has been moistened with the recommended alcohol just prior to use.

Lidocaine Hydrochloride Injection, USP is supplied as follows:

Unit of Sale	Concentration	Each
Single-dose		
NDC 0409-4278-01 Tray of 25	0.5% 250 mg/50 mL (5 mg/mL)	NDC 0409-4278-16 Glass Teartop Vial
NDC 0409-4713-02 Carton of 25	1% 50 mg/5 mL (10 mg/mL)	NDC 0409-4713-12 Glass Ampul
NDC 0409-4713-32 Carton of 50	1% 20 mg/2 mL (10 mg/mL)	NDC 0409-4713-42 Glass Ampul
NDC 0409-4713-62 Case of 800	1% 20 mg/2 mL (10 mg/mL)	NDC 0409-4713-72 Glass Ampul
NDC 0409-4713-65 Case of 800	1% 50 mg/5 mL (10 mg/mL)	NDC 0409-4713-75 Glass Ampul
NDC 0409-4279-02 Carton of 25	1% 300 mg/30 mL (10 mg/mL)	NDC 0409-4279-16 Glass Teartop Vial
NDC 0409-4270-01 Case of 25	1% 300 mg/30 mL (10 mg/mL)	NDC 0409-4270-11 Sterile Glass Teartop Vial
NDC 0409-4776-01 Carton of 25	1.5% 300 mg/20 mL (15 mg/mL)	NDC 0409-4776-10 Glass Ampul
NDC 0409-4056-01 Carton of 5	1.5% 300 mg/20 mL (15 mg/mL)	NDC 0409-4056-11 Sterile Glass Ampul
NDC 0409-4282-01 Bundle of 25	2% 40 mg/2 mL (20mg/mL)	NDC 0409-4282-11 Glass Ampul
NDC 0409-4282-02 Carton of 25	2% 200 mg/10 mL (20 mg/mL)	NDC 0409-4282-12 Glass Ampul

Unit of Sale	Concentration	Each
Multiple-dose:		
NDC 0409-4275-01 Tray of 25	0.5% 250 mg/50 mL (5 mg/mL)	NDC 0409-4275-16 Plastic Fliptop Vial
NDC 0409-4276-01 Tray of 25	1% 200 mg/20 mL (10 mg/mL)	NDC 0409-4276-16 Plastic Fliptop Vial
NDC 0409-4276-02	1% 500 mg/50 mL	NDC 0409-4276-17

Tray of 25	(10 mg/mL)	Plastic Fliptop Vial
NDC 0409-4277-01 Tray of 25	2% 400 mg/20 mL (20 mg/mL)	NDC 0409-4277-16 Plastic Fliptop bottle
NDC 0409-4277-02 Tray of 25	2% 1000 mg/50 mL (20 mg/mL)	NDC 0409-4277-17 Plastic Fliptop Vial

Single-dose products are preservative-free.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Lidocaine Hydrochloride Injection, USP solutions packaged in ampuls and glass teartop vials may be autoclaved one time only. Autoclave at 15 pounds pressure, 121°C (250°F) for 15 minutes. DO NOT AUTOCLAVE PRODUCT IN PLASTIC VIALS.

Revised: 1/2015

EN-3679

Hospira, Inc., Lake Forest, IL 60045 USA

Sodium Chloride Injection, USP is a sterile, nonpyrogenic, isotonic solution of sodium chloride 0.9% (9 mg/mL) in Water for Injection containing no antimicrobial agent or other added substance. The pH is between 4.5 and 7.0. Its chloride and sodium ion concentrates are approximately 0.154 mEq of each per milliliter and its calculated osmolality is 0.308 milliosmols per mL.

Sodium chloride occurs as colorless cubic crystals or white crystalline powder and has a saline taste. Sodium Chloride is freely soluble in water. It is soluble in glycerin and slightly soluble in alcohol. The empirical formula for sodium chloride is NaCl, and the molecular weight is 58.44.

Sodium chloride comprises over 90% of the inorganic constituents of the blood serum. Sodium chloride in water dissociates to provide sodium (Na⁺) and chloride (Cl⁻) ions. These ions are normal constituents of the body fluids (principally extracellular) and are essential for maintaining electrolyte balance. The small volume of Fluid and amount of sodium chloride provided by Sodium Chloride Injection, USP, 0.9% when used only as a vehicle for parenteral injection of drugs, is unlikely to exert a significant effect on fluid and electrolyte balance except possibly in very small infants.

Sodium Chloride Injection is used to flush intravascular catheters or as a sterile, isotonic single dose vehicle, solvent, or diluent for substances to be administered intravenously, intramuscularly or subcutaneously and for other extemporaneously prepared single dose sterile solutions according to instructions of the manufacture of the drug to be administered.

Sodium Chloride must be used with caution in the presence of congestive heart failure, circulatory insufficiency, kidney dysfunction or hypoproteinemia. Excessive amounts of sodium chloride by any route may cause hypokalemia and acidosis.

Excessive amounts by parental routes may precipitate congestive heart failure and acute pulmonary edema, especially seen in patients with preexisting cardiovascular disease and those receiving corticosteroids, corticotropin or other drugs that may give rise to sodium retention. For use in newborns, when a Sodium Chloride solution is required for preparation or diluting medications, or in flushing intravenous catheters, only preservative-free Sodium Chloride Injection, USP, 0.9% should be used.

Since Sodium Chloride Injection does not contain antimicrobial agents and is intended for single use, any unused amount must be discarded immediately following withdrawal of any portion of the contents of the vial or ampul. Do not open ampul until it is to be used.

Consult the manufactures instructions for choice of vehicle, appropriate dilution or volume for dissolving the drug to be injected, including the route and rate of injection.

Pregnancy Category C: Animal reproductive studies have not been conducted with Sodium Chloride Injection USP 0.9%. It is also not known whether Sodium Chloride Injection USP 0.9% can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Chloride Injection USP 0.9% should be given to a pregnant woman only if clearly needed.

Reactions which may occur because of this solution, added drugs or the technique of reconstitution or administration include febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasation.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate countermeasures, and if possible, retrieve and save the remainder of the unused vehicle for examination.

When used as a diluent, solvent or intravascular flushing solution, this parental preparation is unlikely to pose a threat of sodium chloride or fluid overload except possible in very small infants. In the event these should occur, reevaluate the patient and institute appropriate corrective measures.

Before Sodium Chloride Injection, USP, 0.9% is used as a vehicle for the administration of a drug, specific references should be checked for any possible incompatibility with sodium chloride. The volume of the preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration as recommended by the manufacture.

Sodium Chloride Injection, USP, 0.9% is also indicated for use in flushing intravenous catheters. Prior to and after administration of the medication, the intravenous catheter should be flushed in its entirety with Sodium Chloride Injection, USP, 0.9%. Use in accord with any warnings or precautions appropriate to the medication being administered. Parental drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

INSTRUCTIONS FOR USE

To open ampuls, using gauze, place thumb and forefinger on color line, break at constriction.

5 mL ampuls packaged in box of 25 each (NDC-65282-1505-1)

10 mL ampuls packaged in box of 25 each (NDC-65282-1510-1)

Store at controlled room temperature 15-30 C (59-86 F). Avoid freezing.

Manufactured for:

Spectra Medical Devices, Inc. 260-F Fordham Road, Wilmington, MA 01887

By: KM. Pharm. Co., LTD.

SM1500 Rev. B 12/00

Drug Facts

Povidone-iodine

Antiseptic

Antiseptic skin preparation

Do not use

-if allergic to iodine

-in the eyes

For external use only

Ask a doctor before use if injuries are

-deep or puncture wounds

-serious burns

Stop use and ask a doctor if

-redness, irritation, swelling or pain persists or increases

-infection occurs

Avoid pooling beneath patient

Avoid excessive heat. Store at room temperature.

Keep out of reach of children

In case of accidental ingestion, seek professional assistance or consult a poison control center immediately.

Apply locally as needed.

- 1% titratable iodine
- latex free
- for hospital or professional use only

citric acid, disodium phosphate, nonoxynol-9, sodium hydroxide, water


1 800 760-3236 (Mon to Fri 8:30 AM-5:00 PM EST)

APLICARE

POVIDONE-IODINE SOLUTION

ANTISEPTIC

STERILE Solution



10.75"

Drug Facts

<i>Active ingredient</i>	<i>Purpose</i>
Povidone-iodine USP 10%.....	Antiseptic

Use antiseptic skin preparation

Warnings

Do not use ■ if allergic to iodine ■ in the eyes

For external use only ▶

Reorder No. L-3001

APPLICARE, INC.
 MERIDEN, CT 06450
 U.S.A.
 DOLLARD-des-ORMEAUX,
 QUEBEC H9G 2S3
 www.applicare.com
 0609

(01) 0 03523380 00013 3



Questions or comments? ☎ 1 800 760-3236 (Mon to Fri 8:30 AM-5:00 PM EST)

Inactive ingredients citric acid, disodium phosphate, nonoxonyl-9, sodium hydroxide, water

Other information
 ■ 1% titratable iodine ■ latex free
 ■ for hospital or professional use only

Directions apply locally as needed

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or consult a poison control center immediately.

Avoid pooling beneath patient

Stop use and ask a doctor if
 ■ redness, irritation, swelling or pain persists
 or increases
 ■ infection occurs

Warnings
 Ask a doctor before use if injuries are
 ■ deep or puncture wounds ■ serious burns

Drug Facts (continued)

**TEAR HERE
 POUR**



LOT

Package Label Display Panel

RL-2178 (5/07)



2 mL NDC 0409-4713-62

Preservative-Free

1% LIDOCAINE HCl
Injection, USP
10 mg/mL

Hospira, Inc.
Lake Forest, IL 60045 USA **Rx** only

Package Label Display Panel

20 mL Single-dose
Preservative-Free **Rx** only

NDC 0409-4776-01
**FOR INFILTRATION AND NERVE
BLOCK INCLUDING EPIDURAL AND
CAUDAL.** Each mL contains lidocaine
hydrochloride 15 mg; sodium chloride
6.5 mg. May contain sodium hydroxide
and/or hydrochloric acid for pH
adjustment. pH 6.5 (5.0 to 7.0). Sterile,
nonpyrogenic. Usual dosage: See
insert. Store at 20 to 25°C (68 to 77°F).
[See USP Controlled Room
Temperature.]

**1.5% LIDOCAINE
HCl Inj., USP
15 mg/mL**

HOSPIRA, INC., LAKE FOREST, IL 60045 USA RL-0871 (11/04)



Package Label Display Panel

REF A1043-17
SINGLE SHOT EPIDURAL
17G TUOHY



DRUGS:
 1 Lidocaine HCl (1.5%) 20ml
 1 Lidocaine HCl (1%) 2ml
 1 Sodium Chloride (0.9%) 10ml

PROCEDURAL COMPONENTS:
 1 Tuohy Epidural Needle (17G x 3 1/2 in. TW)
 1 18G x 1 1/2 in. Needle
 1 25G x 1/2 in. Needle
 1 Glass Syringe (16ml, Metal Luer Slip)
 1 Plastic Syringe (20ml, Luer Lock)
 1 Plastic Syringe (3ml, Luer Lock)
 1 Medicine Cup, 50ml
 1 Needle Stick Pad
 1 Fanestrated Drape

PREP COMPONENTS:
 2 Nitrile Gloves
 4 Gauze Sponges
 3 Sponge Applicators
 1 Povidone-Iodine Solution, 1/2 oz

WARNINGS:
 • A needle stick with a contaminated needle may cause infectious disease.
 • The use of excessive force while placing needles into the stick pad may cause the needle to protrude through the bottom of the tray which may result in a contaminated needle stick.

PRECAUTIONS:
 • Use Aseptic technique.
 • To help prevent needle-stick injuries, needles should not be recapped or purposely bent. If excessive resistance is met during needle insertion, do not force the needle as damage may occur. To help avoid needle breakage, do not attempt to straighten a bent needle; discard it and complete the procedure with a replacement needle.
 • After use, place sharps in a suitable sharps container. Dispose of contaminated product in a safe manner according to Centers for Disease Control and Prevention (USA) and Federal/State/Local regulations (EPA, OSHA) and health care facility guidelines or local equivalent.
 • Do NOT Facilitate.
 • NOTE: See enclosure(s) for drug information. Confirm drug identity and integrity. Use only if solution is clear and colorless. Do not use if damaged.
 • To be used only by individuals familiar with single shot epidural procedures. For specific techniques and procedures, refer to standard textbooks.

STORE AT CONTROLLED ROOM TEMPERATURE

The Smiths Medical and Portex design marks are trademarks of Smiths Medical. The symbol ® indicates the trademark is registered in the U.S. Patent and Trademark Office and certain other countries.

REF SLUG 2D

Manufacturer:
 Smiths Medical ASD, Inc.
 10 Bowers Drive
 Keene, NH 03431 USA
 www.smiths-medical.com
MADE IN USA

© 2012, 2010 Smiths Medical. All rights reserved.

Caution - Do Not Reuse - Does not contain Natural Rubber Latex - Do not use if package is damaged - Sterilized using ethylene oxide - Caution: Federal (U.S.A.) law restricts this device to sale by or on the order of a physician

FLTA1043-17 REV: 003 04/16

smiths medical

A1043-17 SINGLE SHOT EPIDURAL 17G TUOHY

anesthesia conduction kit kit

Product Information

Product Type	MEDICAL DEVICE	Item Code (Source)	NHRIC:51688-9689
---------------------	----------------	---------------------------	------------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:51688-9689-2	10 in 1 CASE		
1		1 in 1 PACKAGE, COMBINATION; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 AMPULE	10 mL
Part 2	1 AMPULE	20 mL
Part 3	1 PACKET	22.5 mL
Part 4	1 AMPULE	2 mL

Part 1 of 4

SODIUM CHLORIDE

sodium chloride solution

Product Information

Item Code (Source)	NDC:65282-1510
Route of Administration	EPIDURAL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65282-1510-1	10 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/01/2014	

Part 2 of 4

LIDOCAINE HYDROCHLORIDE

lidocaine hydrochloride injection, solution

Product Information

Item Code (Source)	NDC:0409-4776
Route of Administration	EPIDURAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	15 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	6.5 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-4776-01	20 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080408	01/06/2006	

Part 3 of 4

APLICARE POVIDONE-IODINE

povidone-iodine solution

Product Information

Item Code (Source)	NDC:52380-0001
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
NONOXYNOL-9 (UNII: 48Q180SH9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52380-0001-3	22.5 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	03/01/1984	

Part 4 of 4

LIDOCAINE HYDROCHLORIDE

lidocaine hydrochloride injection, solution

Product Information

Item Code (Source)	NDC:0409-4713
Route of Administration	INFILTRATION, EPIDURAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	7 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-4713-62	2 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080408	09/28/1972	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
premarket notification	K965017	12/14/2005	

Labeler - Smiths Medical ASD, Inc. (137835299)

Establishment

Name	Address	ID/FEI	Business Operations
Smiths Medical ASD, Inc.		137835299	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		093132819	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Kwang Myung Pharm. Co., Ltd.		631099384	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Aplicare, Inc.		107255002	manufacture

Revised: 10/2016

Smiths Medical ASD, Inc.