

PEPCID AC MAXIMUM STRENGTH- famotidine tablet, film coated
Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division

PEPCID® AC

Maximum Strength

Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid

reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?

1-800-755-4008 (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 16837-855-14

MAXIMUM STRENGTH


Pepcid®
AC

Famotidine Tablets 20 mg
Acid Reducer

Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

25 Tablets




Peppcid[®] AC
MAXIMUM STRENGTH

Famotidine (active ingredient) made in Hungary
Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19004 USA
©McNeil Consumer Pharmaceuticals Co. 2021
PEPCID[®] and PEPCID AC[®] are registered trademarks
of McNeil-PPC.

MAXIMUM STRENGTH
Peppcid[®] AC
25 Tablets

MAXIMUM STRENGTH
Peppcid[®] AC
Famotidine Tablets 20 mg
Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion



30049137

NDC 16837-855-14

Do not use if blister unit is torn or broken

- 1 tablet relieves heartburn due to acid indigestion
- PEPCID AC[®] prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Drug Facts (continued)

Active ingredient
Famotidine 20 mg.....Acid reducer

Purpose
• relieves heartburn associated with acid indigestion
• prevents heartburn associated with acid indigestion
and sour stomach by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Don't use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.

Uses
• relieves heartburn associated with acid indigestion
and sour stomach
• prevents heartburn associated with acid indigestion
and sour stomach by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Don't use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts (continued)

Active ingredients
Famotidine 20 mg, hydroxypropyl cellulose, hypromellose, croscarmellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Other information
• read the directions and warnings before use
• children under 12 years: ask a doctor
• do not use more than 2 tablets in 24 hours
• cause heartburn
• before eating food or drinking beverages that
cause heartburn, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn.
• to prevent symptoms, swallow 1 tablet with a glass of water. Do not chew.
• to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
• adults and children 12 years and over:
• to prevent symptoms, swallow 1 tablet with a glass of water. Do not chew.
• to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.

Drug Facts (continued)

Keep the container in a cool, dry place. Store at 20°-25°C (68°-77°F).
• protect from moisture

Inactive ingredients
Famotidine 20 mg, hydroxypropyl cellulose, hypromellose, croscarmellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?
1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

Keep the container in a cool, dry place. Store at 20°-25°C (68°-77°F).
• protect from moisture

Inactive ingredients
Famotidine 20 mg, hydroxypropyl cellulose, hypromellose, croscarmellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?
1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn

- Do not lie flat or bend over after eating around the stomach.
- Do not eat before bedtime.
- Raise the head of your bed.
- Avoid heartburn-causing foods such as rich, spicy, fatty, or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables.
- Eat slowly and avoid big meals.
- If overweight, lose weight.
- Quit smoking.





PEPCID AC MAXIMUM STRENGTH

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:16837-855
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	SEMI-CIRCLE (D shaped)	Size	9mm
Flavor		Imprint Code	PAC;20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16837-855-05	1 in 1 CARTON	09/01/2003	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:16837-855-25	5 in 1 CARTON	09/01/2003	03/31/2014
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

3	NDC:16837-855-49	1 in 1 CARTON	09/01/2003	
3		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:16837-855-50	1 in 1 CARTON	09/01/2003	
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:16837-855-52	50 in 1 TRAY	09/01/2003	
5		1 in 1 POUCH; Type 0: Not a Combination Product		
6	NDC:16837-855-65	1 in 1 CARTON	09/01/2003	
6		65 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:16837-855-67	1 in 1 CARTON	09/01/2003	
7		65 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:16837-855-70	1 in 1 CARTON	09/01/2003	
8		70 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
9	NDC:16837-855-75	1 in 1 CARTON	09/01/2003	
9		75 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
10	NDC:16837-855-90	125 in 1 PACKAGE, COMBINATION; Type 0: Not a Combination Product	03/15/2021	
11	NDC:16837-855-80	1 in 1 CARTON	09/01/2003	
11		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
12	NDC:16837-855-85	1 in 1 CARTON	09/01/2003	
12		85 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
13	NDC:16837-855-09	1 in 1 PACKAGE	09/01/2003	
13		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
14	NDC:16837-855-12	1 in 1 CARTON	09/01/2003	
14		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
15	NDC:16837-855-13	1 in 1 POUCH; Type 0: Not a Combination Product	09/01/2003	
16	NDC:16837-855-14	5 in 1 CARTON	09/01/2003	
16		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
17	NDC:16837-855-15	1 in 1 CARTON	09/01/2003	
17		35 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
18	NDC:16837-855-17	1 in 1 CARTON	09/01/2003	
18		65 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
	NDC:16837-			

19	NDC:16837-855-18	1 in 1 CARTON	09/01/2003	
19		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
20	NDC:16837-855-40	5 in 1 CARTON	09/01/2003	
20		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
21	NDC:16837-855-51	12 in 1 CARTON	09/01/2003	
21		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
22	NDC:16837-855-16	1 in 1 CARTON	09/01/2003	
22		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
23	NDC:16837-855-20	2 in 1 PACKAGE	08/22/2014	
23		1 in 1 CARTON		
23		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
24	NDC:16837-855-19	1 in 1 CARTON	04/27/2015	
24		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020325	09/01/2003	

Labeler - Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division (878046358)

Revised: 3/2022

Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division