

VICETON- chlormaphenicol tablet, coated
Bimeda, Inc.

Viceton®

(chloramphenicol tablets)

FOR ORAL USE IN DOGS ONLY

250 mg Tablets

Antibiotic

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Federal law prohibits the use of this drug in food-producing animals.

NADA 055-059, Approved by FDA

Net contents: 500 Tablets

DESCRIPTION:

Chloramphenicol is a broad-spectrum antibiotic shown to have specific therapeutic activity against a wide variety of organisms. Its activity was first demonstrated in culture filtrates from a species of soil organism collected in Venezuela, later designated as ***Streptomyces venezuelae***. The antibiotic was subsequently isolated from culture filtrates (1), identified chemically (2), and later synthesized (3).

Aqueous solutions of chloramphenicol are neutral in pH. Chloramphenicol is stable for several years at room temperature and forms colorless to yellowish-white crystals in the shape of elongated plates or fine needles. It is only slightly soluble in water, but soluble in alcohol and propylene glycol.

Chloramphenicol is exceptionally stable in the presence of high pH, although it is destroyed at pH's in excess of 10. Dissolved in distilled water, it can withstand boiling for five hours (1).

ACTIONS:

At low concentrations, chloramphenicol exerts a bacteriostatic effect on a wide range of pathogenic organisms, including many Gram-positive and Gram-negative bacteria, spirochetes, several rickettsiae and certain large viruses and Mycoplasma (PPL0) (4,5,6,7,8,9,10,11,12,13, and 13a). At high concentrations, it inhibits growth of animal and plant cells.

Chloramphenicol exerts its bacteriostatic action by inhibiting protein synthesis in susceptible organisms. Complete suppression of the assimilation of ammonia and of the incorporation of amino acids, particularly glutamic acid, together with an increased formation of ribonucleic acid (RNA), lead to an inhibition of bacterial growth (4, 14, 15, 16, and 17).

Chloramphenicol antagonizes the action of such antibiotics as penicillin and streptomycin, which act only on growing cells, but is synergistic to tetracycline, which also acts by inhibiting protein synthesis. (18) It is possible the chloramphenicol would produce similar synergism with other antibiotics which act by inhibiting protein synthesis.

In this respect, the experimentally demonstrated synergistic action between chloramphenicol and gamma-globulin should be mentioned. Clinical observations in man and corresponding investigations in laboratory animals experimentally infected with various pathogenic bacteria have shown that a combination of chloramphenicol with gamma-globulin or specific antisera has a greater therapeutic effect than would be expected from a mere addition of the individual effects (19,20, 21, and 22).

Many experiments have revealed that the development of resistance to chloramphenicol is rare compared with that occurring with other important antibiotics. (23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, and 34) Bacterial resistance may develop in some strains against chloramphenicol but has been encountered only infrequently in clinical usage.

Chloramphenicol achieves maximum serum levels very rapidly following oral, intravenous and intraperitoneal administration. Intramuscular injection with chloramphenicol, except certain soluble forms, results in a somewhat delayed absorption and lower serum levels than when given by the oral, intravenous, or intraperitoneal route.

Chloramphenicol diffuses readily into all body tissues, but at different concentrations. Highest concentrations are found in the liver and kidney of dogs indicating that these organs are the main route of inactivation and excretion of the metabolites. The lungs, spleen, heart and skeletal muscles contain concentrations similar to that of the blood (27, 35, 36, 37, 38, and 39).

Chloramphenicol reaches significant concentration in the aqueous and vitreous humors of the eye from the blood (4).

A significant difference from other antibiotics is its marked ability to diffuse into the cerebrospinal fluid. Within three to four hours after administration, the concentration in the cerebrospinal fluid has reached, on the average, 50% of the concentration in the serum. If the meninges are inflamed, the percentage may be even higher (4, 27, 29, 40, 41, 42, and 43).

Chloramphenicol diffuses readily into milk, pleural and ascitic fluids and crosses the placenta attaining concentrations of about 75% of that of the maternal blood (36 and 44).

Chloramphenicol is rather rapidly metabolized, mainly in the liver, by conjugation with glucuronic acid.

Approximately 55% of a single daily dose can be recovered from the urine of a treated dog. A small fraction of this is in the form of unchanged chloramphenicol (36).

A single intravenous dose of 150 mg/kg (approximately 68 mg/lb) in propylene glycol is the maximum dose tolerated by the dog. (45) No toxic effect was observed when dogs were administered orally, 200 mg/kg (approximately 91 mg/lb) daily for over four months. (40) In the mouse, the LD-50 is 150-250 mg/kg (68 to 114 mg/lb) body weight by intravenous injection and 1,500 mg/kg (approximately 681 mg/lb) by the oral administration (1 and 46).

INDICATIONS:

Chloramphenicol Tablets are recommended for oral treatment of the following conditions in dogs:

- Bacterial pulmonary infections caused by susceptible microorganisms such as: *Staphylococcus aureus*, *Streptococcus pyogenes* and *Brucella bronchiseptica*.
- Infections of the urinary tract caused by susceptible microorganisms such as: *Escherichia coli*, *Proteus vulgaris*, *Corynebacterium renale*, *Streptococcus* spp., and *hemolytic Staphylococcus*.
- Enteritis caused by susceptible microorganisms such as: *E. coli*, *Proteus* spp., *Salmonella* spp., and *Pseudomonas* spp.
- Infections associated with canine distemper caused by susceptible microorganisms such as: *B. bronchiseptica*, *E. coli*, *P. aeruginosa*, *Proteus* spp., *Shigella* spp. and *Neisseria catarrhalis*.

Additional adjunctive therapy should be used when indicated. Most susceptible infectious disease organisms will respond to chloramphenicol therapy in three to five days when the recommended dosage regimen is followed. If no response to chloramphenicol therapy is obtained in three to five days, discontinue its use and review the diagnosis. Also, a change of therapy should be considered.

Laboratory tests should be conducted including *in vitro* culturing and susceptibility tests on samples

collected prior to treatment.

CONTRAINDICATIONS:

Because of potential antagonism, chloramphenicol should not be administered simultaneously with penicillin or streptomycin.

WARNING:

NOT FOR USE IN ANIMALS WHICH ARE RAISED FOR FOOD PRODUCTION. CHLORAMPHENICOL PRODUCTS SHOULD NOT BE ADMINISTERED IN CONJUNCTION WITH OR 2 HOURS PRIOR TO THE INDUCTION OF GENERAL ANESTHESIA WITH PENTOBARBITAL BECAUSE OF PROLONGED RECOVERY.

CHLORAMPHENICOL SHOULD NOT BE ADMINISTERED TO DOGS MAINTAINED FOR BREEDING PURPOSES. SOME EXPERIMENTS INDICATE THAT CHLORAMPHENICOL CAUSES IN EXPERIMENTAL ANIMALS, PARTICULARLY IN FEMALES, SIGNIFICANT DISORDERS IN MORPHOLOGY AS WELL AS IN FUNCTION OF THE GONADS.

HAZARDS AND PRECAUTIONS:

1. This antibiotic contains a chemical structure (nitrobenzene group) that is characteristic of a group of drugs long known to depress hematopoietic activity of the bone marrow (54).
2. *In vitro*-tissue culture studies using canine bone marrow cells have demonstrated that extremely high concentrations of chloramphenicol inhibit both uptake of iron by the nucleated red cells and incorporation of iron into heme (55).
3. Chloramphenicol products should not be administered in conjunction with or two hours prior to the induction of general anesthesia with pentobarbital because of prolonged recovery time.
4. Chloramphenicol products should not be administered to dogs maintained for breeding purposes. Some experiments indicate that chloramphenicol causes, in experimental animals, particularly females, significant disorders in morphology as well as in function of the gonads.

ADVERSE REACTIONS:

Certain individual dogs may exhibit transient vomiting or diarrhea after an oral dose of 25 mg/lb body weight (49).

DOSAGE AND ADMINISTRATION:

Dogs-25 mg/lb body weight every 6 hours for oral administration.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Keep out of reach of children.

Store at or below 25°C (77°F) in a dry place.

HOW SUPPLIED:

- 250 mg.....Bottles of 500's, 1000's
- 500 mg.....Bottles of 500's
- 1 gram.....Bottles of 100's

1. Ehrlich, J., Q.R. Bartz, R.M. Smith, D.A. Joslyn and P.R. Burkholder (1947): Science 106, 127.
2. Rebstock, M.C., H.M. Crooks, Jr., J. Controulis and Q.R. Bartz (1949): J. Amer. Chem. Soc. 71, 2458-2462.
3. Controulis, J., M.C. Rebstock and H.M. Crooks, Jr. (1949): J. Amer. Chem. Soc. 71, 2463-2468.

4. Barber, M. and L.P. Garrod (1963): Antibiotic and Chemo-therapy. E. and S. Livingston, Edinburgh, London: 116 to 128.
5. Christi, A.B. (1963): Practitioner 191, 588-593.
6. Lange, A. and E. Vogel (1958): Z. Hyg. infektionskrankh. 144, 555-563. Ref.: Landw. Zbl. Abt. IV, 5, 1960, 2188.
7. Schirmer, R.C., F.E. Eades and J.P. Newman (1952): Vet. Med. 46, 485-490 Ref.: Vet. Rec. 64, 1952, 597.
8. Smadel, J.E. and E.B. Jackson (1947): Science 106, 418-419.
9. Zettl, K. (1960): Vet.-Med. Nachr, 25-35.
10. Godzeski, C.W. and D.E. Pavey (1965): Nature 205, No. 4975, 1017 to 1018.
11. Hille, G. and E. Ludwig (1964): Mh. Vet.-Med. 20, 704-708.
12. Hubrig, Th. and D. Schimmel (1965): Mh. Vet.-Med. 20, 171-174.
13. Schimmel, D. and Th. Hubrig (1965): mh. Vet.-Med. 20, 334-338.
- 13a. Hagen, Wm. A. and Bruner, D.W. (Third Edition, 1957): The Infectious Diseases of Domestic Animals, Cornell University Press. 1479.
14. Brock, Th. D. (1963): J. Bact. 85, 527-531.
15. Brock, T.D. Chloramphenicol, Bacterial. Rev. 25: 32-48, 1961.
16. Saurat, P. and R. Lautie (1959): Rev. Med. Veterin. 22, 689-701.
17. Schaub, F. (1962): Schweiz, Arch. Tierheilk, 104, 373-384.
18. Reedy, R.J., W.W. Wright, E.J. Oswald and M. Ostrolenk (1958): Antibiot. Ann. 1957/58, 745-750.
19. Fisher, M.W. (1957) Antib. Chemoth. 7, 315-321.
20. Manning, M.C., L.A. Gagliardi and M.W. Fisher (1958): Antibiot. Ann. 1957/58, 566-571.
21. Reedy, R.J., E.J. Oswald and W.W. Wright (1958): Antibiot. Ann. 1957/58, 581-584.
22. Waisbren, B.A. (1957) Antib. Chemoth. 7, 315-321.
23. Beseth, L. (1963): Nord. Vet.-Med. 15, 745-754.
24. Gedek, W., R. Dunkel and L. Kotter (1964): Tierarztl, Umsch. 19, 337-340.
25. Kettner, H. and A. Braunsteiner (1960): Tierarztl, Umsch. 15, 249-255.
26. Kortum, H. (1963): Vet. Med. Diss. Munchen.
27. Kraft, H. (1960): Tierarztl, Umsch. 15, 58-60.
28. Kraft, H. (1962): Tierarztl, Umsch. 17, 278-280.
29. Lettow, E. (1954): Tierarztl, Umsch. 9, 227-230.
30. Manten, A., E.H. Kampelmacher and P.A.M. Guinee (1963): T. Dierganeesk. 88, 411-417.
31. Palyusik, M. (1963)" Mag. allator. Lapja 18, 356-357. Ref.: Vet Bull. 34, 1964, 317.
32. Ramsay, C.H. and P.R. Edwards (1961): Appl. Microbiolog. 9, 389-391.
33. Smith, H.W. (1954): Vet. Rec. 66, 42-43.
34. Smith, W. H. and W.E. Crabb (1956): Vet. Rec. 68, 274-277.
35. English, P.B. and A.A. Seawright (1961): Aust. Vet. J. 37, 9-13.
36. Glazko, A.J., L.M. Wolf, W.A. Dill and A.C. Bratton, Jr. (1949): J. Pharmacol. exp. Ther. 96, 445-

449.

37. Glazko, A.J., H.E. Carnes, A. Kazenko, L.M. Wolf and T.F. Teutner (1958): *Antibiot. Ann.* 1957/58, 792-802.
38. Eads, F.E., A.J. Glazko, L.M. Wolf, J. Erlich and M. Galbraith (1952): *Amer. J. Vet. Res.* 13, 204-206.
39. Leopold, I.H., Nichols, A.C. and Voegl, A.W., Penetration of Chloramphenicol U.S>P. (Chloromycetin) into the Eye. *Arch. Opth.*, 44:22, 1950.
40. Gruhzt, O.M., R.R. Risken, T.F. Reutner and E. Martino (1949): *J. Clin. Invest.* 28, 943-951.
41. McCrumb, Jr., R.R., M.J. Snyder and W.J. Hicken (1958): *Antibiot. Ann.* 1957/58, 837-841.
42. Ross S.H. Bischoff, W. Preissner and W. Orr (1949): *J. Clin. Invest.* 28, 1050.
43. Roy, T.E., E. Krieger, G. Craig, D. Cohen, G.A. McNaughton and N. Silverthorne (1952): *Antib. Chemoth.* 2, 505-516.
44. Walter, M. and L. Heilmeyer (1954): *Antibiotika-Fibel.* Georg Thieme Verlag, Stuttgart, 1954.
45. Ehrlich, J. chloramphenicol. *Encyclopedia of Chemical Technology.* 4: 928-936, 1964.
46. Cheymol, J. (1955): *Arzeimeittel Forsch.* 5, 1-9.
47. Smith, R.M., D.A. Joslyn, O.M. Gruhzt, I.W. McLean, M.A. Penner and J. Ehrlich (1948): *J. Bact.* 55, 425-448.
48. Van Looveren, L.J. (1958): *Schweiz. Arch. Tierheilk.* 100, 459-467.
49. Sogtrop, H.H. (1956): *Tierarztl, Umsch.* 11, 257-261.
50. Hrenoff, A.K. (1962): *Med. exp.* 6, 118-122.
51. Nelson, A.A. and J.L. Radomski (1954): *Antib. Chemoth* 4, 1174-1180.
52. Reutner, T.F., R.E. Maxwell, K.E. Weston and J.K. Weston (1955): *Antib. Chemoth.* 5, 679-711.
53. Weston, J.D., T.F. Reutner, R.Q. Thompson and R.E. Maxwell (1954): *Fed. Proc.* 13, 416.
54. Jones, L.M., *Veterinary Pharmacology and Therapeutics*, 3rd Edition, Iowa State University Press, Ames, Iowa, 539-542, 1965.
55. Ward, H.P., The Effect of Chloramphenicol on RNA and Heme Synthesis in Bone Marrow Cultures. *Journ. Lab and Clinical Med.*, 68: 403, 1966.
56. The Merck Veterinary Manual, 2nd Edition, Merck and Co., Inc., Rahway, N.J., 522-524, 1961.

Chloramphenicol should not be administered to dogs maintained for breeding purposes. Some experiments indicate that chloramphenicol causes in experimental animals, particularly in females, significant disorders in morphology as well as in function of the gonads.



Manufactured by:
Bimeda, Inc.,
Le Sueur, MN 56058, USA
www.bimeda.com

N.A. Corp Address:
Bimeda, Inc., One Tower Lane,
Oakbrook Terrace, IL 60181

Viceton® is a Registered Trademark of Bimeda, Inc.
Product No. 1VIC007 8VIC008 Rev. 07/17




Viceton®
(chloramphenicol tablets)
FOR ORAL USE IN DOGS ONLY
250 mg Tablets
Antibiotic
NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN
CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
Federal law prohibits the use of this drug in food-producing animals.
NADA 055-059, Approved by FDA

Net
Contents: 500 Tablets

INDICATIONS: Viceton® (chloramphenicol tablets) are indicated for treatment of the following conditions in dogs caused by chloramphenicol susceptible microorganisms: Bacterial gastroenteritis associated with bacterial diarrhea, bacterial pulmonary infections and bacterial infections of the urinary tract.

DOSAGE: Dogs-25 mg/lb body weight every 6 hours via oral administration.

WARNING: Not for use in animals which are raised for food production. Chloramphenicol products should not be administered in conjunction with or 2 hours prior to the induction of general anesthesia with pentobarbital because of prolonged recovery.

Read package insert carefully before use.
Store at or below 25°C (77°F) in a dry place.

LOT:
EXP:

Chloramphenicol should not be administered to dogs maintained for breeding purposes. Some experiments indicate that chloramphenicol causes in experimental animals, particularly in females, significant disorders in morphology as well as in function of the gonads.



Manufactured by:
Bimeda, Inc.,
Le Sueur, MN 56058, USA
www.bimeda.com

N.A. Corp Address:
Bimeda, Inc., One Tower Lane,
Oakbrook Terrace, IL 60181

Viceton® is a Registered Trademark of Bimeda, Inc.
Product No. 1VIC008 8VIC009 Rev. 07/17



Bimeda

Viceton®

(chloramphenicol tablets)

FOR ORAL USE IN DOGS ONLY

500 mg Tablets

Antibiotic

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
Federal law prohibits the use of this drug in food-producing animals.

NADA 055-059, Approved by FDA 

Net Contents: 500 Tablets

INDICATIONS: Viceton® (chloramphenicol tablets) are indicated for treatment of the following conditions in dogs caused by chloramphenicol susceptible microorganisms: Bacterial gastroenteritis associated with bacterial diarrhea, bacterial pulmonary infections and bacterial infections of the urinary tract.

DOSAGE: Dogs-25 mg/lb body weight every 6 hours via oral administration.

WARNING: Not for use in animals which are raised for food production. Chloramphenicol products should not be administered in conjunction with or 2 hours prior to the induction of general anesthesia with pentobarbital because of prolonged recovery.

Read package insert carefully before use.
Store at or below 25°C (77°F) in a dry place.

LOT:
EXP:

Bimeda

Viceton®

(chloramphenicol tablets)

FOR ORAL USE IN DOGS ONLY

1 Gram Tablets

Antibiotic

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
Federal law prohibits the use of this drug in food-producing animals.

NADA 055-059, Approved by FDA 

Net Contents: 100 Tablets

INDICATIONS: Viceton® (chloramphenicol tablets) are indicated for treatment of the following conditions in dogs caused by chloramphenicol susceptible microorganisms: Bacterial gastroenteritis associated with bacterial diarrhea, bacterial pulmonary infections and bacterial infections of the urinary tract.

DOSAGE: Dogs-25 mg/lb body weight every 6 hours via oral administration.

WARNING: Not for use in animals which are raised for food production. Chloramphenicol products should not be administered in conjunction with or 2 hours prior to the induction of general anesthesia with pentobarbital because of prolonged recovery.

Read package insert carefully before use.
Store at or below 25°C (77°F) in a dry place.

LOT:
EXP:

chlormaphenicol tablet, coated

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:6 1133-2212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORAMPHENICOL (UNII: 66974FR9Q1) (CHLORAMPHENICOL - UNII:66974FR9Q1)	CHLORAMPHENICOL	250 mg

Product Characteristics

Color	yellow	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:6 1133-2212-1	500 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055059	10/23/2017	

VICETON

chlormaphenicol tablet, coated

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:6 1133-2213
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORAMPHENICOL (UNII: 66974FR9Q1) (CHLORAMPHENICOL - UNII:66974FR9Q1)	CHLORAMPHENICOL	500 mg

Product Characteristics

Color	yellow	Score	2 pieces
Shape	ROUND	Size	12mm

Flavor		Imprint Code	
Contains			
Packaging			
#	Item Code	Package Description	Marketing Start Date
1	NDC:6 1133-2213-1	500 in 1 BOTTLE	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055059	01/01/2018	

VICETON			
chlormaphenicol tablet, coated			
Product Information			
Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:6 1133-2210
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	CHLORAMPHENICOL (UNII: 66974FR9Q1) (CHLORAMPHENICOL - UNII:66974FR9Q1)	CHLORAMPHENICOL	1 g
Product Characteristics			
Color	yellow	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	
Contains			
Packaging			
#	Item Code	Package Description	Marketing Start Date
1	NDC:6 1133-2210-1	100 in 1 BOTTLE	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055059	01/01/2018	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 12/2020

Bimeda, Inc.