

**ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE- esterified estrogens, methyltestosterone tablet**  
**Anzalp Pharmsolutions USA LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**Esterified Estrogens and Methyltestosterone Tablets**

**1- Product Name: Esterified Estrogens and Methyltestosterone Tablets**

**2- Name and Strength of the active Ingredient (s)**

Each tablet contains:

Esterified Estrogens, USP.....0.625 mg

Methyltestosterone, USP.....1.25 mg

**3- Product Description**

**Physical properties**

green, oval (oblong(caplets)), oral tablets

**4- Pharmacodynamics/**

**Pharmacokinetics**

**Pharmacotherapeutic group**

**Estrogens**

Estrogens are important in the development and maintenance of the female reproductive system and secondary sex characteristics. They promote growth and development of the vagina, uterus, and fallopian tubes, and enlargement of the breasts. Indirectly, they contribute to the shaping of the skeleton, maintenance of tone and elasticity of urogenital structures, changes in the epiphyses of the long bones that allow for the pubertal growth spurt and its termination, growth of axillary and pubic hair, and pigmentation of the nipples and genitals. Decline of estrogenic activity at the end of the menstrual cycle can bring on menstruation, although the cessation of progesterone secretion is the most important factor in the mature ovulatory cycle. However, in the preovulatory or nonovulatory cycle, estrogen is the primary determinant in the onset of menstruation. Estrogens also affect the release of pituitary gonadotropins. The pharmacologic effects of esterified estrogens are similar to those of endogenous estrogens. They are soluble in water and are well absorbed from the gastrointestinal tract.

In responsive tissues (female genital organs, breasts, hypothalamus, pituitary) estrogens enter the cell and are transported into the nucleus. As a result of estrogen action, specific RNA and protein synthesis occurs.

**Estrogen Pharmacokinetics**

Metabolism and inactivation occur primarily in the liver. Some estrogens are excreted into the bile; however, they are reabsorbed from the intestine and returned to the liver through the portal venous system. Water soluble esterified estrogens are strongly acidic

and are ionized in body fluids, which favor excretion through the kidneys since tubular reabsorption is minimal.

## **Androgens**

Endogenous androgens are responsible for the normal growth and development of the male sex organs and for maintenance of secondary sex characteristics. These effects include the growth and maturation of prostate, seminal vesicles, penis, and scrotum; the development of male hair distribution, such as beard, pubic, chest and axillary hair, laryngeal enlargement, vocal cord thickening, alterations in body musculature, and fat distribution. Drugs in this class also cause retention of nitrogen, sodium, potassium, phosphorus, and decreased urinary excretion of calcium. Androgens have been reported to increase protein anabolism and decrease protein catabolism. Nitrogen balance is improved only when there is sufficient intake of calories and protein. Androgens are responsible for the growth spurt of adolescence and for the eventual termination of linear growth centers. In children, exogenous androgens accelerate linear growth rates, but may cause a disproportionate advancement in bone maturation. Use over long periods may result in fusion of the epiphyseal growth centers and termination of growth process. Androgens have been reported to stimulate the production of red blood cells by enhancing the production of erythropoietic stimulating factor.

## **Androgen Pharmacokinetics**

Testosterone given orally is metabolized by the gut and 44 percent is cleared by the liver in the first pass. Oral doses as high as 400 mg per day are needed to achieve clinically effective blood levels for full replacement therapy. The synthetic androgens (methyltestosterone and fluoxymesterone) are less extensively metabolized by the liver and have longer half-lives. They are more suitable than testosterone for oral administration.

Testosterone in plasma is 98 percent bound to a specific testosterone estradiol binding globulin, and about 2 percent is free. Generally, the amount of this sex-hormone binding globulin in the plasma will determine the distribution of testosterone between free and bound forms, and the free testosterone concentration will determine its half-life.

About 90 percent of a dose of testosterone is excreted in the urine as glucuronic and sulfuric acid conjugates of testosterone and its metabolites; about 6 percent of a dose is excreted in the faeces, mostly in the unconjugated form. Inactivation of testosterone occurs primarily in the liver. Testosterone is metabolized to various 17-keto steroids through two different pathways. There are considerable variations of the half-life of testosterone as reported in the literature, ranging from 10 to 100 minutes.

In many tissues the activity of testosterone appears to depend on reduction to dihydrotestosterone, which binds to cytosol receptor proteins. The steroid-receptor complex is transported to the nucleus where it initiates transcription events and cellular changes related to androgen action.

## **5- Indication**

ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE HALF STRENGTH are indicated in the treatment of: Moderate to severe vasomotor symptoms associated with the menopause in those patients not improved by estrogens alone. (There is no evidence that estrogens are effective for nervous symptoms or depression without associated

vasomotor symptoms, and they should not be used to treat such conditions.)

ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE HALF STRENGTH HAVE NOT BEEN SHOWN TO BE EFFECTIVE FOR ANY PURPOSE DURING PREGNANCY AND ITS USE MAY CAUSE SEVERE HARM TO THE FETUS

## **6- Recommended Dose**

### **Usual Dosage Range**

ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE HALF STRENGTH daily as recommended by the physician.

Treated patients with an intact uterus should be monitored closely for signs of endometrial cancer and appropriate diagnostic measures should be taken to rule out malignancy in the event of persistent or recurring abnormal vaginal bleeding.

### **7- Recommended route of administration:**

Given cyclically for short-term use only:

For treatment of moderate to severe vasomotor symptoms associated with the menopause in patients not improved by estrogen alone.

The lowest dose that will control symptoms should be chosen and medication should be discontinued as promptly as possible. Administration should be cyclic (e.g., three weeks on and one week off). Attempts to discontinue or taper medication should be made at three-to-six-month intervals.

## **8- Contraindications**

Estrogens should not be used in women with any of the following conditions:

1. Known or suspected cancer of the breast except in appropriately selected patients being treated for metastatic disease.
2. Known or suspected estrogen-dependent neoplasia.
3. Known or suspected pregnancy.
4. Undiagnosed abnormal genital bleeding.
5. Active thrombophlebitis or thromboembolic disorders.
6. A past history of thrombophlebitis, thrombosis, or thromboembolic disorders associated with previous estrogen use (except when in treatment of breast malignancy).

Methyltestosterone should not be used in:

1. The presence of severe liver damage.
2. Pregnancy and in breast-feeding mothers because of the possibility of masculinization of the female fetus or breast-fed infant.

## **9- Warnings and Precautions**

Associated with Estrogens

Induction of malignant neoplasms. Long term continuous administration of natural and synthetic estrogens in certain animal species increases this frequency of carcinomas of the breast, cervix, vagina, and liver. There is now evidence that estrogens increase the risk of carcinoma of the endometrium in humans.

At the present time there is no satisfactory evidence that estrogens given to postmenopausal women increase the risk of cancer of the breast,<sup>18</sup> although a recent long-term follow-up of a single physician's practice has raised this possibility.<sup>18a</sup> Because of the animal data, there is a need for caution in prescribing estrogens for women with a strong family history of breast cancer or who have breast nodules, fibrocystic disease, or abnormal mammograms.

Gallbladder disease. A recent study has reported a 2 to 3-fold increase in the risk of surgically confirmed gallbladder disease in women receiving postmenopausal estrogens,<sup>18</sup> similar to the 2-fold increase previously noted in users of oral contraceptives as in the case of oral contraceptives the increased risk appeared after two years of use.

Effects similar to those caused by estrogen-progesterone oral contraceptives. There are several serious adverse effects of oral contraceptives, most of which have not, up to now, been documented as consequences of postmenopausal estrogen therapy. This may reflect the comparatively low doses of estrogen used in postmenopausal women. It would be expected that the larger doses of estrogen used to treat prostatic or breast cancer or postpartum breast engorgement are more likely to result in these adverse effects, and, in fact, it has been shown that there is an increased risk of thrombosis in men receiving estrogens for prostatic cancer and women for postpartum breast engorgement.

a. Thromboembolic disease. It is now well established that users of oral contraceptives have an increased risk of various thromboembolic and thrombotic vascular diseases, such as thrombophlebitis, pulmonary embolism, stroke, and myocardial infarction. Cases of retinal thrombosis, mesenteric thrombosis, and optic neuritis have been reported in oral contraceptive users. There is evidence that the risk of several of these adverse reactions is related to the dose of the drug. An increased risk of post-surgery thromboembolic complications has also been reported in users of oral contraceptives. If feasible, estrogen should be discontinued at least 4 weeks before surgery of the type associated with an increased risk of thromboembolism, or during periods of prolonged immobilization.

While an increased rate of thromboembolic and thrombotic disease in postmenopausal users of estrogens has not been found, this does not rule out the possibility that such an increase may be present or that subgroups of women who have underlying risk factors or who are receiving relatively large doses of estrogens may have increased risk. Therefore, estrogens should not be used in persons with active thrombophlebitis or thromboembolic disorders, and they should not be used (except in treatment of malignancy) in persons with a history of such disorders in association with estrogen use. They should be used with caution in patients with cerebral vascular or coronary artery disease and only for those in whom estrogens are clearly needed.

Large doses of estrogen (5 mg esterified estrogens per day), comparable to those used to treat cancer of the prostate and breast, have been shown in a large prospective clinical trial in men to increase the risk of nonfatal myocardial infarction, pulmonary embolism and thrombophlebitis. When estrogen doses of this size are used, any of the

thromboembolic and thrombotic adverse effects associated with oral contraceptive use should be considered a clear risk.

b. Hepatic adenoma. Benign hepatic adenomas appear to be associated with the use of oral contraceptives. Although benign and rare, these may rupture and may cause death through intra-abdominal hemorrhage. Such lesions have not yet been reported in association with other estrogen or progestogen preparations but should be considered in estrogen users having abdominal pain and tenderness, abdominal mass, or hypovolemic shock. Hepatocellular carcinoma has also been reported in women taking estrogen-containing oral contraceptives. The relationship of this malignancy to these drugs is not known at this time.

c. Elevated blood pressure. Increased blood pressure is not uncommon in women using oral contraceptives. There is now a report that this may occur with use of estrogens in the menopause and blood pressure should be monitored with estrogen use, especially if high doses are used.

d. Glucose tolerance. A worsening of glucose tolerance has been observed in a significant percentage of patients of estrogen-containing oral contraceptives. For this reason, diabetic patients should be carefully observed while receiving estrogens.

Hypercalcemia. Administration of estrogens may lead to severe hypercalcemia in patients with breast cancer and bone metastases. If this occurs, the drug should be stopped and appropriate measures taken to reduce the serum calcium level.

### **Associated with Methyltestosterone**

In patients with breast cancer, androgen therapy may cause hypercalcemia by stimulating osteolysis. In this case the drug should be discontinued.

Prolonged use of high doses of androgens has been associated with the development of peliosis hepatis and hepatic neoplasms including hepatocellular carcinoma. (See PRECAUTIONS – Carcinogenesis).

Peliosis hepatis can be a life-threatening or fatal complication.

Cholestatic hepatitis and jaundice occur with 17-alpha-alkyl androgens at a relatively low dose. If cholestatic hepatitis with jaundice appears or if liver function tests become abnormal, the androgen should be discontinued and the etiology should be determined. Drug-induced jaundice is reversible when the medication is discontinued.

Edema with or without heart failure may be a serious complication in patients with preexisting cardiac, renal, or hepatic disease. In addition to discontinuation of the drug, diuretic therapy may be required.

### **Precautions**

#### **Associated with Estrogens**

##### **A. General Precautions**

A complete medical and family history should be taken prior to the initiation of any estrogen therapy. The pretreatment and periodic physical examinations should include special reference to blood pressure, breasts, abdomen, and pelvic organs, and should include a Papanicolaou smear. As a general rule, estrogen should not be prescribed for longer than one year without another physical examination being performed.

Fluid retention– Because estrogens may cause some degree of fluid retention,

conditions which might be influenced by this factor such as asthma, epilepsy, migraine, and cardiac or renal dysfunction, require careful observation.

Certain patients may develop undesirable manifestations of excessive estrogenic stimulation, such as abnormal or excessive uterine bleeding, mastodynia, etc.

Oral contraceptives appear to be associated with an increased incidence of mental depression. Although it is not clear whether this is due to the estrogenic or progestogenic component of the contraceptive, patients with a history of depression should be carefully observed.

Preexisting uterine leiomyomata may increase in size during estrogen use.

The pathologist should be advised of estrogen therapy when relevant specimens are submitted.

Patients with a past history of jaundice during pregnancy have an increased risk of recurrence of jaundice while receiving estrogen-containing oral contraceptive therapy. If jaundice develops in any patient receiving estrogen, the medication should be discontinued while the cause is investigated.

Estrogens may be poorly metabolized in patients with impaired liver function and they should be administered with caution in such patients.

Because estrogens influence the metabolism of calcium and phosphorus, they should be used with caution in patients with metabolic bone diseases that are associated with hypercalcemia or in patients with renal insufficiency.

Because of the effects of estrogens on epiphyseal closure, they should be used judiciously in young patients in whom bone growth is not complete.

Certain endocrine and liver function tests may be affected by estrogen-containing oral contraceptives. The following similar changes may be expected with larger doses of estrogen:

- a. Increased sulfobromophthalein retention.
- b. Increased prothrombin and factors VII, VIII, IX and X; decreased antithrombin 3; increased norepinephrine induced platelet aggregability.
- c. Increased thyroid binding globulin (TBG) leading to increased circulating total thyroid hormone, as measured by PBI, T4 by column, or T4 by radioimmunoassay. Free T3 resin uptakes is decreased, reflecting the elevated TBG; free T4 concentration is unaltered.
- d. Impaired glucose tolerance.
- e. Decreased pregnanediol excretion.
- f. Reduced response to metyrapone test.
- g. Reduced serum folate concentration.
- h. Increased serum triglyceride and phospholipid concentration.

## B. Pregnancy Category X

See CONTRAINDICATIONS.

## C. Nursing Mothers

As a general principle, the administration of any drug to nursing mothers should be done only when clearly necessary since many drugs are excreted in human milk.

Associated with Methyltestosterone

## A. General Precautions

Women should be observed for signs of virilization (deepening of the voice, hirsutism, acne, clitoromegaly, and menstrual irregularities). Discontinuation of drug therapy at the

time of evidence of mild virilism is necessary to prevent irreversible virilization. Such virilization is usual following androgen use at high doses.

Prolonged dosage of androgen may result in sodium and fluid retention. This may present a problem, especially in patients with compromised cardiac reserve or renal disease.

Hypersensitivity may occur rarely.

PBI may be decreased in patients taking androgens.

Hypercalcemia may occur. If this does occur, the drug should be discontinued.

## B. Information for the Patient

The physician should instruct patients to report any of the following side effects of androgens:

Women: Hoarseness, acne, changes in menstrual periods, or more hair on the face.

All Patients: Any nausea, vomiting, changes in skin color or ankle swelling.

## C. Laboratory Tests

Women with disseminated breast carcinoma should have frequent determination of urine and serum calcium levels during the course of androgen therapy.

Because of the hepatotoxicity associated with the use of 17-alpha-alkylated androgens, liver function tests should be obtained periodically.

Hemoglobin and hematocrit should be checked periodically for polycythemia in patients who are receiving high doses of androgens.

## D. Drug Interactions

Anticoagulants C-17 substituted derivatives of testosterone, such as methandrostenolone, have been reported to decrease the anticoagulant requirements of patients receiving oral anticoagulants. Patients receiving oral anticoagulant therapy require close monitoring, especially when androgens are started or stopped.

Oxyphenbutazone. Concurrent administration of oxyphenbutazone and androgens may result in elevated serum levels of oxyphenbutazone.

Insulin. In diabetic patients the metabolic effects of androgens may decrease blood glucose and insulin requirements.

## E. Drug/Laboratory Test Interferences

Androgens may decrease levels of thyroxine-binding globulin, resulting in decreased T4 serum levels and increased resin uptake of T3 and T4. Free thyroid hormone levels remain unchanged, however, and there is no clinical evidence of thyroid dysfunction.

## F. Carcinogenesis

Animal Data. Testosterone has been tested by subcutaneous injection and implantation in mice and rats. The implant induced cervical-uterine tumours in mice, which metastasized in some cases. There is suggestive evidence that injection of testosterone into some strains of female mice increases their susceptibility to hepatoma.

Testosterone is also known to increase the number of tumours and decrease the degree of differentiation of chemically induced carcinomas of the liver in rats.

Human Data. There are rare reports of hepatocellular carcinoma in patients receiving

long-term therapy with androgens in high doses. Withdrawal of the drugs did not lead to regression of the tumours in all cases.

Geriatric Patients treated with androgens may be at an increased risk for the development of prostatic hypertrophy and prostatic carcinoma.

#### G. Pregnancy

Teratogenic Effects. Pregnancy Category X (see CONTRAINDICATIONS).

#### H. Nursing Mothers

It is not known whether androgens are excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from androgens, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

### **10- Undesirable Effects**

#### Associated with Estrogens

(See Warnings regarding induction of neoplasia, adverse effects on the fetus, increased incidence of gallbladder disease, and adverse effects similar to those of oral contraceptives, including thromboembolism). The following additional adverse reactions have been reported with estrogenic therapy, including oral contraceptives:

Genitourinary system.

Breakthrough bleeding, spotting, change in menstrual flow.

Dysmenorrhea.

Premenstrual-like syndrome.

Amenorrhea during and after treatment.

Increase in size of uterine fibromyomata.

Vaginal candidiasis.

Change in cervical erosion and in degree of cervical secretion.

Cystitis-like syndrome.

#### **Breasts.**

Tenderness, enlargement, secretion.

#### **Gastrointestinal.**

Nausea, vomiting.

Abdominal cramps, bloating.

Cholestatic jaundice.

#### **Skin.**

Chloasma or melasma which may persist when drug is discontinued.



Erythema multiforme.

Erythema nodosum.

Hemorrhagic eruption.

Loss of scalp hair.

Hirsutism.

### **Eyes.**

Steepening of corneal curvature.

Intolerance to contact lenses.

### **CNS.**

Headache, migraine, dizziness.

Mental depression.

Chorea.

### **Miscellaneous.**

Increase or decrease in weight.

Reduced carbohydrate tolerance.

Aggravation of porphyria.

Edema.

Changes in libido.

## **Associated with Methyltestosterone**

### **A. Endocrine and Urogenital.**

Female: The most common side effects of androgen therapy are amenorrhea and other menstrual irregularities, inhibition of gonadotropin secretion, and virilization, including deepening of the voice and clitoral enlargement. The latter usually is not reversible after androgens are discontinued. When administered to a pregnant woman androgens cause virilization of external genitalia of the female fetus.

Skin and Appendages: Hirsutism, male pattern of baldness, and acne.

Fluid and Electrolyte Disturbances: Retention of sodium, chloride, water, potassium, calcium, and inorganic phosphates.

Gastrointestinal: Nausea, cholestatic jaundice, alterations in liver function test, rarely hepatocellular neoplasms, and peliosis hepatis.

Hematologic: Suppression of clotting factors, II, V, VII, and X, bleeding in patients on concomitant anticoagulant therapy, and polycythemia.

Nervous System: Increased or decreased libido, headache, anxiety, depression, and generalized paresthesia.

Metabolic: Increased serum cholesterol.

Miscellaneous: Inflammation and pain at site of intramuscular injection or subcutaneous implantation of testosterone containing pellets, stomatitis with buccal preparations, and rarely anaphylactoid reactions.

## 11- Overdose and Treatment

Numerous reports of ingestion of large doses of estrogen-containing oral contraceptives by young children indicate that serious ill effects do not occur. Overdosage of estrogen may cause nausea, and withdrawal bleeding may occur in females.

There have been no reports of acute overdosage with the androgens.

## 12- Packaging and storage condition

ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE TABLETS Half Strength in bottles of 100.

ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE TABLETS Half Strength green, oval (oblong(caplets)), oral tablets. Contains: 0.625 mg of Esterified Estrogens, USP and 1.25 mg of Methyltestosterone, USP.

Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F). [See USP Controlled Room Temperature.]

Rx only

## 13- References

- Herbst, A., et al.: Obstet. Gynecol. 40: 287-298, 1972.
- Stafl, A., et al.: Obstet. Gynecol. 43: 118-128, 1974.
- Janerich, D.T., et al.: N. Engl. J. Med. 291: 697-700, 1974.
- Estrogens for Oral or Parenteral Use: Federal Register 40: 8212, 1975.
- Hoover, R. et al.: N. Engl. J. Med. 295: 401-405, 1976.
- Boston Collaborative Drug Surveillance Program: Lancet 1: 1399-1404, 1973.
- The Veterans Administration Cooperative Urological Research Group: J. Urol, 98: 516-522, 1967.
- Inman, W.H.W., et al.: Br. Med. J. 2: 193-199, 1968.

Anzalp Pharmasolutions Pvt Ltd  
Healthcare with Ethics

**Dosage:** See package insert for full prescribing information.

**DISPENSER:** Include one patient information leaflet "WHAT YOU SHOULD KNOW ABOUT ESTROGENS" with each package dispensed.

**Distributed by:** Anzalp Pharmasolutions USA  
511 S Harbor Blvd Ste F, La Habra, CA 90631-9375.

## Packaging

NDC no. 84346-450-01



Anzalp Pharmasolutions Pvt Ltd  
Healthcare with Ethics

## Esterified Estrogens and Methyltestosterone Tablets

HALF STRENGTH

0.625mg/1.25mg

### Each tablet contains:

Esterified Estrogens, USP ..... 0.625mg  
Methyltestosterone, USP ..... 1.25mg

**Dosage:** See package insert for full prescribing information.

**DISPENSER:** Include one patient information leaflet "WHAT YOU SHOULD KNOW ABOUT ESTROGENS" with each package dispensed.

Store at 20° - 25°C (68° - 77°F); excursions permitted to 15° - 30°C (59° - 86°F). [See USP Controlled Room Temperature.]

Distributed by: Anzalp Pharmasolutions USA  
511 S Harbor Blvd Ste F, La Habra, CA 90631-9375.



## ESTERIFIED ESTROGENS AND METHYLTESTOSTERONE

esterified estrogens, methyltestosterone tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:84346-450
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIII

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ESTROGENS, ESTERIFIED</b> (UNII: 3ASP8Q3768) (ESTROGENS, ESTERIFIED - UNII:3ASP8Q3768)	ESTROGENS, ESTERIFIED	0.625 mg
<b>METHYLTESTOSTERONE</b> (UNII: V9EFU16ZIF) (METHYLTESTOSTERONE - UNII:V9EFU16ZIF)	METHYLTESTOSTERONE	1.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	

<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL (Oblong (Caplets))	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84346-450-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/20/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/20/2024	

**Labeler** - Anzalp Pharmasolutions USA LLC (119239230)

Revised: 5/2024

Anzalp Pharmasolutions USA LLC