

SODIUM BICARBONATE ANTACID- sodium bicarbonate tablet
Rugby Laboratories

4624 - Rugby

Drug Facts

Active ingredient (in each tablet)

Sodium bicarbonate 10 gr (650 mg)

Purpose

Antacid

Uses

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms

Warnings

Do not take more than 24 tablets for adults up to 60 years of age (or 12 tablets for adults 60 years of age and older) in a 24-hour period nor use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Ask a doctor before use if you have a sodium restricted diet.

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not use the maximum dosage for more than 2 weeks
- tablets may be swallowed whole or dissolved in water prior to use
- **adults 60 years of age and over:** 1-2 tablets every 4 hours, not more than 12 tablets in 24 hours

- **adults under 60 years of age:** 1-4 tablets every 4 hours, not more than 24 tablets in 24 hours

Other information

- **each tablet contains:** sodium 178 mg
- store in a dry place at 15° - 30°C (59° - 86°F).

Inactive ingredients

croscarmellose sodium, microcrystalline cellulose, silica, stearic acid, talc.

Questions or comments?

1-800-231-4670

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Distributed by:

RUGBY® LABORATORIES

Indianapolis, IN 46268

Questions or comments?

Call (800) 616-2471

www.majorpharmaceuticals.com

WARNING: THIS PRODUCT CAN EXPOSE YOU TO CHEMICALS INCLUDING LEAD, WHICH IS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER AND BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. FOR MORE INFORMATION GO TO WWW.P65WARNINGS.CA.GOV.

Rev. 10/22 R-29

Rugby

NDC 0536-1047-10

Sodium Bicarbonate 10 gr (650 mg)

ANTACID

1000 TABLETS

Rugby
NDC 0536-1047-10

Sodium Bicarbonate 10 gr (650 mg)

ANTACID

1000 TABLETS

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each tablet)

Sodium bicarbonate 10 gr (650 mg)

Purpose

Antacid

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach ■ upset stomach associated with these symptoms

Warnings

Do not take more than 24 tablets for adults up to 60 years of age (or 12 tablets for adults 60 years of age and older) in a 24-hour period nor use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Ask a doctor before use if you have a sodium restricted diet.

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Do not use the maximum dosage for more than 2 weeks ■ tablets may be swallowed whole or dissolved in water for use ■ adults 60 years of age and over: 1-2 tablets every 4 hours, not more than 12 tablets in 24 hours

■ adults under 60 years of age: 1-4 tablets every 4 hours, not more than 24 tablets in 24 hours

Other information ■ each tablet contains: sodium 176 mg ■ store in a dry place at 15° - 30°C (59° - 86°F).

Inactive ingredients croscarmellose sodium, microcrystalline cellulose, silica, stearic acid, TiO_2

Questions or comments? 1-800-231-4670


Distributed by:

RUSBY LABORATORIES
Indianapolis, IN 46286

Questions or comments?
call (800) 231-4671
www.majorpharmaceuticals.com

CA WARNING: THIS PRODUCT CAN EXPOSE YOU TO CHEMICALS INCLUDING LEAD, WHICH IS HARMFUL TO THE STATE OF CALIFORNIA TO CAUSE CANCER AND BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. FOR MORE INFORMATION GO TO WWW.P65WARNINGS.CA.GOV.

Rev. 10/22 R-23 Re-order No. 370387 R51365



SODIUM BICARBONATE ANTACID

sodium bicarbonate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1047
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (SODIUM CATION - UNII:L4YR4MONH37)	SODIUM BICARBONATE	650 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	CPC77
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1047-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/25/2015	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001		03/25/2015	

Labeler - Rugby Laboratories (079246066)

Revised: 1/2024

Rugby Laboratories