

**MAGNESIUM CITRATE SALINE LAXATIVE- magnesium citrate liquid
TOP CARE (Topco Associates LLC)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Magnesium citrate 1.745g per fl. oz.

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Do not use

laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet
- a sodium restricted diet
- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 1 week

Ask a doctor or pharmacist before use if you

are presently taking a prescription drug.

When using this product

do not exceed the maximum recommended daily dosage in a 24 hour period.

Stop use and ask a doctor if

you have rectal bleeding or no bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age	1/2 bottle to 1 bottle (10 fl oz)
children 6 to 11 years of age	1/3 bottle to 1/2 bottle
children under 6 years of age	consult a doctor

Other information

- **each fluid ounce contains:** magnesium 280 mg, sodium 37 mg
- store between 46-86°F (8-30°C). Do not refrigerate.
- protect from excessive heat and freezing
- discard unused product within 24 hours of opening bottle

Inactive ingredients

citric acid, flavors, purified water, FD&C red 40, saccharin sodium, sodium bicarbonate

Questions or comments?

Call toll free 1-888-423-0139

Principal Display Panel

Magnesium Citrate

SALINE LAXATIVE

ORAL SOLUTION

For Relief of Occasional Constipation

CHERRY FLAVOR

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007 © TOPCO

topcare@topco.com

www.topcarebrand.com

Package Label

NDC:36800-626-10



Magnesium Citrate

SALINE LAXATIVE
ORAL SOLUTION

For Relief of Occasional
Constipation

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

10 FL OZ (296 mL)



CHERRY
FLAVOR



Drug Facts

Active ingredient (in each fl oz) Magnesium citrate 1.745g.....**Purpose** Saline laxative

Uses ■ relieves occasional constipation (irregularity)
■ generally produces bowel movement in 1/2 to 6 hours

Warnings
Do not use laxative products for longer than 1 week unless directed by a doctor.

Ask a doctor before use if you have
 ■ kidney disease ■ a magnesium restricted diet
 ■ a sodium restricted diet ■ stomach pain, nausea, or vomiting
 ■ noticed a sudden change in bowel habits that lasts more than 1 week

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

When using this product, do not exceed the maximum recommended daily dosage in a 24 hour period.

Stop use and ask a doctor if you have rectal bleeding or no bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. ▶

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
 © TOPCO PLVA0318 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com

Drug Facts (continued)

Directions
 ■ drink a full glass (8 ounces) of liquid with each dose
 ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	1/2 bottle to 1 bottle (10 fl oz)
children 6 to 11 years of age	1/3 bottle to 1/2 bottle
children under 6 years of age	consult a doctor

Other information
 ■ each fluid ounce contains: magnesium 280 mg, sodium 37 mg
 ■ store between 46-86°F (8-30°C). Do not refrigerate.
 ■ protect from excessive heat and freezing
 ■ discard unused product within 24 hours of opening bottle

Inactive ingredients citric acid, FD&C red #40, flavor, purified water, saccharin sodium, sodium bicarbonate

Questions or comments?
 Call toll free 1-888-423-0139

PLD-1273D
LB005171





0 36800 09266 2

TOPCARE HEALTH Magnesium Citrate

MAGNESIUM CITRATE SALINE LAXATIVE			
magnesium citrate liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-626
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)		MAGNESIUM CITRATE	1.745 g in 29.6 mL
Inactive Ingredients			
	Ingredient Name		Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			

WATER (UNII: 059QF0KO0R)

SACCHARIN SODIUM (UNII: SB8ZUX40TY)

FD&C RED NO. 40 (UNII: WZB9127XOA)

SODIUM BICARBONATE (UNII: 8MDF5V39QO)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-626-10	296 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	07/31/2015	07/31/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	07/31/2015	07/31/2025

Labeler - TOP CARE (Topco Associates LLC) (006935977)

Revised: 12/2022

TOP CARE (Topco Associates LLC)