

TOPCARE CLOTRIMAZOLE 3- clotrimazole cream
Topco Associates LLC

TopCare®
Clotrimazole 3

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
Clotrimazole USP 2% (100 mg in each applicatorful)	Vaginal Antifungal
Clotrimazole USP 2% (external cream)	Vaginal Antifungal

Use

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

Warnings

For vaginal use only

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have

- **vaginal itching and discomfort for the first time**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

When using this product

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief ask a doctor before using another product

Stop use and ask a doctor if

- **symptoms do not get better in 3 days**
- **symptoms last more than 7 days**
- **you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- before using this product read the enclosed educational brochure for complete directions and information
- adults and children 12 years of age and over:
 - **vaginal cream:** insert one applicatorful of cream into the vagina at bedtime for 3 days in a row. Throw applicator away after use.
 - **external cream:** use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
- children under 12 years of age: ask a doctor

Other information

- To open: unscrew cap, use pointed end on cap to puncture seal
- do not use if carton is opened
- **safety sealed:** the tube opening should be sealed. If the seal has been punctured or is not visible, do not use the product.
- store between 20° to 25°C (68° to 77°F)
- see flap of carton or crimp of tube for lot number and expiration date

Inactive ingredients

benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate-60, purified water, and sorbitan monostearate

Questions or comments?

Call toll-free, **1-888-423-0139** for our 24 hour automated response system.

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007

PRINCIPAL DISPLAY PANEL - 21 g Tube Carton

TopCare®

NDC 36800-062-00

3-DAY TREATMENT

Clotrimazole 3

CLOTRIMAZOLE VAGINAL CREAM USP (2%)
VAGINAL ANTIFUNGAL

• **Cures most vaginal yeast infections**

QUALITY
GUARANTEED

COMPARE TO
GYNE-LOTRIMIN 3®
active ingredients*

One 21g (0.74 OZ) tube of three-day vaginal cream
with three disposable applicators

NET WT 0.74 OZ (21 g) TUBE

TopCare®



Vaginal Yeast Infection Relief

- Three-day treatment
- Cures most vaginal yeast infections
- Three disposable applicators



TopCare®

NDC 36800-062-00

3-DAY TREATMENT

Clotrimazole 3

CLOTRIMAZOLE VAGINAL CREAM USP (2%)
VAGINAL ANTIFUNGAL

- Cures most vaginal yeast infections

One 21g (0.74 OZ) tube of three-day vaginal cream with three disposable applicators

NET WT 0.74 OZ (21 g) TUBE



COMPARE TO
GYNE-LOTRIMIN 3®
active ingredients*



Drug Facts (continued)

Other information • To open: unscrew cap, use pointed end on cap to puncture seal
do not use if carton is opened
safety sealed: the tube opening should be sealed. If the seal has been punctured or is not visible, do not use the product.
store between 20° to 25° C (68° to 77° F)
see flap of carton or crimp of tube for lot number and expiration date

Inactive ingredients benzyl alcohol, cetyl esters wax, 2-octyldodecanol, polyorbata-60, purified water, and sorbitan monostearate

Questions or comments? Call toll-free, 1-888-423-0139 for our 24 hour automated response system.

*This product is not manufactured or distributed by MSD Consumer Care Inc., a subsidiary of Merck & Co., Inc.

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
1-888-423-0139 ©TOPCO TA8912
MADE IN CANADA

Directions • before using this product read the enclosed educational brochure for complete directions and information
• adults and children 12 years of age and over:
• vaginal cream: insert one applicatorful of cream into the vagina at bedtime for 3 days in a row. Throw applicator away after use.
• external cream: use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
• children under 12 years of age: ask a doctor

Stop use and ask a doctor if
• symptoms do not get better in 3 days • symptoms last more than 7 days
• you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

• do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
• mild increase in vaginal burning, itching or irritation may occur
• if you do not get complete relief ask a doctor before using another product

TopCare®

Drug Facts

Active ingredients
Clotrimazole USP 2% (100 mg in each applicator)(1)..... Vaginal Antifungal
Clotrimazole USP 2% (external cream)..... Vaginal Antifungal

Use • Treats vaginal yeast infections
• relieves external itching and irritation due to a vaginal yeast infection


Warnings For vaginal use only
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have
• vaginal itching and discomfort for the first time
• lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
• vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
• been exposed to the human immunodeficiency virus (HIV) that causes AIDS

When using this product
• Do not use because of severe allergic reactions or other medical conditions. Continue and discontinue use as directed and follow the treatment instructions.

This TOP CARE® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

Clotrimazole 3
CLOTRIMAZOLE VAGINAL CREAM USP (2%)
VAGINAL ANTIFUNGAL




TopCare®
Clotrimazole 3
CLOTRIMAZOLE VAGINAL
CREAM USP (2%)
VAGINAL ANTIFUNGAL

**NO COPY ON THIS FLAP
FOR LOT # AND EXPIRY
DATE PRINT**

LPK-3877-3
0912-3
M133

T1128B



TOPCARE CLOTRIMAZOLE 3

clotrimazole cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-062
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Clotrimazole (UNII: G07GZ97H65) (Clotrimazole - UNII:G07GZ97H65)	Clotrimazole	2 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
benzyl alcohol (UNII: LKG8494WBH)	
cetostearyl alcohol (UNII: 2DMT128M1S)	
cetyl esters wax (UNII: D072FFP9GU)	
octyldodecanol (UNII: 461N1O614Y)	
polysorbate 60 (UNII: CAL22UVI4M)	
water (UNII: 059QF0KO0R)	
sorbitan monostearate (UNII: NVZ4I0H58X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-062-00	1 in 1 CARTON		
1		21 g in 1 TUBE, WITH APPLICATOR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021143	04/12/2000	

Labeler - Topco Associates LLC (006935977)

Registrant - Taro Pharmaceuticals U.S.A., Inc. (145186370)

Establishment

Name	Address	ID/FEI	Business Operations
Taro Pharmaceuticals Inc.		206263295	MANUFACTURE(36800-062)

Revised: 1/2013

Topco Associates LLC