

**FAMILY CARE COLD AND FLU MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid**

**United Exchange Corp.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

Active ingredients (in each 30 mL dose cup)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- running nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** if sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers

- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL dose cup contains:
- potassium 3 mg
- sodium 7 mg
- store at room temperature

Inactive ingredients

acesulfame potassium, alcohol, cherry essence, citric acid hydrate, FD&C Blue No. 1, FD&C Red No. 40, high fructose corn syrup, L-menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate hydrate, sodium saccharin

Distributed by:

United Exchange Corp.

Cerritos, CA 90703 U.S.A.

Made in Korea

**TWIST TO OPEN**

**Drug Facts (continued)**

- trouble urinating due to enlarged prostate gland • a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

**When using this product** • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days, redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions** • take only as directed • use dose cup or tablespoon (TBS) • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBS) every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information** • each 30 mL dose cup contains: potassium 3 mg • sodium 7 mg • store at room temperature

**Inactive ingredients** • acesulfame potassium, alcohol, cherry essence, citric acid hydrate, FD&C Blue No. 1, FD&C Red No. 40, high fructose corn syrup, L-menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate hydrate, sodium saccharin

TAMPER EVIDENT: Do not use if printed shrink-band is missing or broken. Failure to follow these warnings could result in serious consequences.  
\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu Cherry.

\*Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu Cherry

**PARENTS:** Learn about teen medicine abuse. www.StopMedicineAbuse.org

ACETAMINOPHEN  
DEXTROMETHORPHAN HBr  
DOXYLAMINE SUCCINATE  
4 FL OZ (118 mL)

**Drug Facts**

**Active ingredients (in each 30 mL dose cup) Purpose**

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

**Uses** temporarily relieves common cold/flu symptoms: • cough due to minor throat and bronchial irritation • sore throat • headache • minor aches and pains • fever • runny nose and sneezing

**Warnings** Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks daily while using this product **Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, hausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleep

**Ask a doctor before use if you have** • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema

**SIDE**

**TWIST TO OPEN**

FAMILY CARE COLD AND FLU MULTI SYMPTOM			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:65923-630
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	
<b>Inactive Ingredients</b>			
Ingredient Name	Strength		
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)			
ALCOHOL (UNII: 3K9958V90M)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			

<b>FD&amp;C BLUE NO. 1</b> (UNII: HBR47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>METHYL PARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOLS</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL PARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0K00R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65923-630-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/30/2015	

**Labeler** - United Exchange Corp. (840130579)

Revised: 7/2015

United Exchange Corp.