

PAIN RELIEF CHILDRENS- acetaminophen tablet, chewable
Rite Aid Corporation

Rite Aid 44-450 Childrens

Active ingredient (in each chewable tablet)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - flu
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has
liver disease.

Ask a doctor or pharmacist before use if your child is
taking the blood thinning drug warfarin.

When using this product
do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

NDC 11822-4450-8

Compare to the active ingredient in **Children's Tylenol®****

FOR AGES 2-11 YEARS

**CHILDREN'S PAIN RELIEF
ACETAMINOPHEN**

ACETAMINOPHEN 160mg CHEWABLE TABLETS

PAIN RELIEVER/FEVER REDUCER

contains no aspirin or ibuprofen

chew or crush tablets completely
before swallowing.

ACTUAL SIZE

**BUBBLE GUM
FLAVOR**

**24
CHEWABLE TABLETS**

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by Johnson
& Johnson Corporation, owner of the registered trademark
Children's Tylenol®. 50844 REV0522C45008

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE
CAMP HILL, PA 17011

If you're not satisfied, we'll happily refund your money.

no print / no varnish area

lot no. & exp. date

0 11822 68619 8

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

NDC 11822-4450-8 Compare to the active ingredient in **Children's Tylenol®****

FOR AGES 2-11 YEARS

CHILDREN'S PAIN RELIEF ACETAMINOPHEN

ACETAMINOPHEN 160 mg CHEWABLE TABLETS

PAIN RELIEVER/FEVER REDUCER
contains no aspirin or ibuprofen
chew or crush tablets completely before swallowing.

BUBBLE GUM FLAVOR

24

CHEWABLE TABLETS

CHILDREN'S PAIN RELIEF ACETAMINOPHEN

ACETAMINOPHEN 160 mg CHEWABLE TABLETS

PAIN RELIEVER/FEVER REDUCER contains no aspirin or ibuprofen chew or crush tablets completely before swallowing

24

CHEWABLE TABLETS

Drug Facts (continued)

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

Inactive ingredients citric acid, croscopolone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrolates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments? 1-800-426-9391

**This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®. 50844 RE 05/22/45008

Drug Facts (continued)

■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use

- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

SATISFACTION GUARANTEE

If you're not satisfied, we'll happily refund your money.

Drug Facts (continued)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each chewable tablet)
Acetaminophen 160 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use

- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

B-1702-450-08-R2
REV05/22C45008

PAIN RELIEF CHILDRENS

acetaminophen tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-4450
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	pink	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	BUBBLE GUM	Imprint Code	44;450
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4450-8	1 in 1 CARTON	02/25/2005	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-4450-2	1 in 1 CARTON	02/25/2005	09/03/2021
2		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/25/2005	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-4450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-4450) , pack(11822-4450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-4450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-4450)

Revised: 5/2023

Rite Aid Corporation