

CVS MOTION SICKNESS FAST MELTING- meclizine hydrochloride tablet, orally disintegrating

CVS Pharmacy, Inc.

CVS Motion Sickness Fast Melting

Active ingredient (in each tablet)

Meclizine hydrochloride 25mg

Purpose

Meclizine hydrochloride Antiemetic

Uses

■ For the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use in children under 12 years of age unless directed by a physician

Ask a doctor before use if you have

- glaucoma ■ a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a physician or pharmacist before use if you are

- taking sedatives or tranquilizers

When using this product

- you may get drowsy ■ avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

To prevent motion sickness take it at least one hour before traveling

Adults and children 12 years of age and over: take 1 to 2 (25 to 50 mg) tablets once daily, or as directed by a physician.

Other information

- Store at 20-30°C (68-86°F)

Inactive ingredients

acesulfame potassium, erythritol, hydroxypropyl cellulose, colloidal silicon dioxide, mannitol, menthol, sodium stearyl fumarate, yellow ferric oxide.



FOR THE PREVENTION AND TREATMENT OF TRAVEL SICKNESS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN, OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts

Active ingredient (in each tablet) Purpose
Meclizine hydrochloride 25mg.....Antiemetic

Uses

For the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use in children under 12 years of age unless directed by a physician.

Ask a doctor before use if you have

glucose a breathing problem such as emphysema or chronic bronchitis
difficulty in urination due to enlargement of the prostate gland

Ask a physician or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

you may get drowsy avoid alcoholic beverages
be careful when driving a motor vehicle or operating machinery
alcohol, sedatives, and tranquilizers may increase the drowsiness effect

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

to prevent motion sickness, take it at least one hour before traveling
adults and children 12 years of age and over: take 1 to 2 (15 to 50 mg) tablets once daily, or as directed by a physician.

Other information

store at 20-30°C (68-86°F)

Inactive ingredients

acesulfame potassium, erythritol, hydroxypropyl cellulose, colloidal silicon dioxide, mannitol, menthol, sodium stearyl fumarate, yellow ferric oxide

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in Japan



US-A



#998892

Actual Size



12 Fast-Melting Tablets (25 mg EACH)

- Less drowsy formula
- Lasts up to 24 hours
- Prevents nausea, dizziness & vomiting

MECLIZINE HYDROCHLORIDE / ANTIEMETIC

MOTION SICKNESS
FAST MELTING

CVS
pharmacy

NEW!

MECLIZINE HYDROCHLORIDE / ANTIEMETIC

MOTION SICKNESS
FAST MELTING

CVS
pharmacy

CVS
pharmacy

MOTION SICKNESS
FAST MELTING

CVS
pharmacy
MOTION SICKNESS
FAST MELTING

CVS MOTION SICKNESS FAST MELTING

meclizine hydrochloride tablet, orally disintegrating

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-534
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ERYTHRITOL (UNII: RA96B954X6)	
MANNITOL (UNII: 3OWL53L36A)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
LEVOMENTHOL (UNII: BZ1R15MTK7)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	SJ
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-534-01	2 in 1 CARTON	08/27/2015	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	08/27/2015	

