

CVS MOTION SICKNESS FAST MELTING- meclizine hydrochloride tablet, orally disintegrating
CVS Pharmacy, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS BOX

Active ingredient (in each tablet)

Meclizine hydrochloride 25mg

Purpose

Meclizine hydrochloride Antiemetic

Uses

- For the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use in children under 12 years of age unless directed by a physician

Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a physician or pharmacist before use if you are

- taking sedatives or tranquilizers

When using this product

- you may get drowsy
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

To prevent motion sickness take it at least one hour before traveling

Adults and children 12 years of age and over: take 1 to 2 (25 to 50 mg) tablets once daily, or as directed by a physician.

Other information

- Store at 20-30°C (68-86°F)

Inactive ingredients

acesulfame potassium, erythritol, hydroxypropyl cellulose, colloidal silicon dioxide, mannitol, menthol, sodium stearyl fumarate, yellow ferric oxide.



FOR THE PREVENTION AND TREATMENT OF TRAVEL SICKNESS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN, OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts

Active ingredient (in each tablet) Purpose
 Meclizine hydrochloride 25mg.....Antiemetic

Uses
 ■ for the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness

Warnings
 Do not use in children under 12 years of age unless directed by a physician

Ask a doctor before use if you have
 ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis
 ■ difficulty in urination due to enlargement of the prostate gland

Ask a physician or pharmacist before use if you are
 ■ taking sedatives or tranquilizers

When using this product
 ■ you may get drowsy ■ avoid alcoholic beverages
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
 ■ to prevent motion sickness, take it at least one hour before traveling
 ■ adults and children 12 years of age and over: take 1 to 2 (15 to 50 mg) tablets once daily, or as directed by a physician

Other information
 ■ store at 20-30°C (68-86°F)

Inactive ingredients
 acesulfame potassium, erythritol, hydroxypropyl cellulose, colloidal silicon dioxide, mannitol, menthol, sodium stearyl fumarate, yellow ferric oxide

EXP: LOT:

Distributed by: **CVS Pharmacy, Inc.**
 One CVS Drive, Woonsocket, RI 02895
 © 2015 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS
 Made in Japan



US-A



#998892



Actual Size

12 Fast-Melting Tablets (25 mg EACH)

- Prevents nausea, dizziness & vomiting
- Lasts up to 24 hours
- Less drowsy formula

MECLIZINE HYDROCHLORIDE / ANTIEMETIC

MOTION SICKNESS FAST MELTING



NEW!

MECLIZINE HYDROCHLORIDE / ANTIEMETIC

MOTION SICKNESS FAST MELTING



MOTION SICKNESS FAST MELTING

MOTION SICKNESS FAST MELTING



NEW!

MOTION SICKNESS

FAST MELTING

MECLIZINE HYDROCHLORIDE / ANTIEMETIC

- Prevents nausea, dizziness & vomiting
- Lasts up to 24 hours
- Less drowsy formula

12 Fast-Melting Tablets (25 mg EACH)



Actual Size

CVS MOTION SICKNESS FAST MELTING

meclizine hydrochloride tablet, orally disintegrating

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-534
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ERYTHRITOL (UNII: RA96B954X6)	
MANNITOL (UNII: 3OWL53L36A)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
HYDROXYPROPYL CELLULOSE (TYPE H) (UNII: RFW2ET671P)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UJ)	
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LEVOMENTHOL (UNII: BZ1R15MTK7)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	SJ
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-534-01	1 in 1 CARTON		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	08/27/2015	

Labeler - CVS Pharmacy, Inc. (062312574)

Revised: 8/2015

CVS Pharmacy, Inc.