

**ITRACONAZOLE- itraconazole capsule**  
**Amneal Pharmaceuticals LLC**

-----

**Itraconazole Capsules, USP**  
**(100 mg)**  
**Rx Only**

## BOXED WARNING

### Congestive Heart Failure, Cardiac Effects and Drug Interactions

#### Congestive Heart Failure and Cardiac Effects:

- Itraconazole capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF. If signs or symptoms of congestive heart failure occur during administration of itraconazole capsules, discontinue administration.
- When itraconazole was administered intravenously to dogs and healthy human volunteers, negative inotropic effects were seen (see **CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations** for more information).


#### Drug Interactions:

- Co-administration of a number of CYP3A4 substrates are contraindicated with itraconazole capsules. Some examples of drugs that are contraindicated for co-administration with itraconazole capsules are: methadone, disopyramide, dofetilide, dronedarone, quinidine, isavuconazole, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam, pimozide, triazolam, felodipine, nisoldipine, ivabradine, ranolazine, eplerenone, cisapride, naloxegol, lomitapide, lovastatin, simvastatin, avanafil, ticagrelor, finerenone, voclosporin.
- Co-administration with colchicine, fesoterodine and solifenacin is contraindicated in subjects with varying degrees of renal or hepatic impairment.
- Co-administration with eliglustat is contraindicated in subjects that are poor or intermediate metabolizers of CYP2D6 and in subjects taking strong or moderate CYP2D6 inhibitors.
- Co-administration with venetoclax is contraindicated in patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) during the dose initiation and ramp-up phase of venetoclax (see **PRECAUTIONS: Drug Interactions Section** for specific examples).
- Co-administration with itraconazole can cause elevated plasma concentrations of these drugs and may increase or prolong both the pharmacologic effects and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of *torsades de pointes*, a potentially fatal arrhythmia (see **CONTRAINDICATIONS and WARNINGS Sections, and PRECAUTIONS: Drug Interactions Section** for specific examples).

## DESCRIPTION

Itraconazole, USP is an azole antifungal agent. Itraconazole, USP is a 1:1:1:1 racemic

mixture of four diastereomers (two enantiomeric pairs), each possessing three chiral centers. It may be represented by the following structural formula and nomenclature:



(±)-1-[(R<sup>\*</sup>)-sec-butyl]-4-[p-[4-[p-[[(2R<sup>\*</sup>,4S<sup>\*</sup>)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one mixture with (±)-1-[(R<sup>\*</sup>)-sec-butyl]-4-[p-[4-[p-[[(2S<sup>\*</sup>,4R<sup>\*</sup>)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one

or

(±)-1-[(RS)-sec-butyl]-4-[p-[4-[p-[[(2R,4S)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one

Itraconazole, USP has a molecular formula of C<sub>35</sub>H<sub>38</sub>Cl<sub>2</sub>N<sub>8</sub>O<sub>4</sub> and a molecular weight of 705.64. It is a white or almost white powder. It is insoluble in water, very slightly soluble in alcohols, and freely soluble in dichloromethane. It has a pKa of 3.70 (based on extrapolation of values obtained from methanolic solutions) and a log (n-octanol/water) partition coefficient of 5.66 at pH 8.1.

Itraconazole capsules, USP contain 100 mg of itraconazole, USP coated on sugar spheres. Inactive ingredients are black iron oxide, FD&C Blue No. 1, gelatin, hypromellose, liquid glucose, microcrystalline cellulose, polyethylene glycol, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, starch (of maize), strong ammonia solution, sucrose and titanium dioxide.

## CLINICAL PHARMACOLOGY

### Pharmacokinetics and Metabolism

#### General Pharmacokinetic Characteristics

Peak plasma concentrations of itraconazole are reached within 2 to 5 hours following oral administration. As a consequence of non-linear pharmacokinetics, itraconazole accumulates in plasma during multiple dosing. Steady-state concentrations are generally reached within about 15 days, with C<sub>max</sub> values of 0.5 mcg/mL, 1.1 mcg/mL and 2

mcg/mL after oral administration of 100 mg once daily, 200 mg once daily and 200 mg b.i.d., respectively. The terminal half-life of itraconazole generally ranges from 16 to 28 hours after single dose and increases to 34 to 42 hours with repeated dosing. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. Itraconazole mean total plasma clearance following intravenous administration is 278 mL/min. Itraconazole clearance decreases at higher doses due to saturable hepatic metabolism.

## **Absorption**

Itraconazole is rapidly absorbed after oral administration. Peak plasma concentrations of itraconazole are reached within 2 to 5 hours following an oral capsule dose. The observed absolute oral bioavailability of itraconazole is about 55%.

The oral bioavailability of itraconazole is maximal when itraconazole capsules are taken immediately after a full meal. Absorption of itraconazole capsules is reduced in subjects with reduced gastric acidity, such as subjects taking medications known as gastric acid secretion suppressors (e.g., H<sub>2</sub>-receptor antagonists, proton pump inhibitors) or subjects with achlorhydria caused by certain diseases (see **PRECAUTIONS: Drug Interactions**). Absorption of itraconazole under fasted conditions in these subjects is increased when itraconazole capsules are administered with an acidic beverage (such as a non-diet cola). When itraconazole capsules were administered as a single 200-mg dose under fasted conditions with non-diet cola after ranitidine pretreatment, a H<sub>2</sub>-receptor antagonist, itraconazole absorption was comparable to that observed when itraconazole capsules were administered alone (see **PRECAUTIONS: Drug Interactions**).

Itraconazole exposure is lower with the capsule formulation than with the oral solution when the same dose of drug is given (see **WARNINGS**).

## **Distribution**

Most of the itraconazole in plasma is bound to protein (99.8%), with albumin being the main binding component (99.6% for the hydroxy-metabolite). It has also a marked affinity for lipids. Only 0.2% of the itraconazole in plasma is present as free drug. Itraconazole is distributed in a large apparent volume in the body (> 700 L), suggesting extensive distribution into tissues. Concentrations in lung, kidney, liver, bone, stomach, spleen and muscle were found to be two to three times higher than corresponding concentrations in plasma, and the uptake into keratinous tissues, skin in particular, up to four times higher. Concentrations in the cerebrospinal fluid are much lower than in plasma.

## **Metabolism**

Itraconazole is extensively metabolized by the liver into a large number of metabolites. *In vitro* studies have shown that CYP3A4 is the major enzyme involved in the metabolism of itraconazole. The main metabolite is hydroxy-itraconazole, which has *in vitro* antifungal activity comparable to itraconazole; trough plasma concentrations of this metabolite are about twice those of itraconazole.

## **Excretion**

Itraconazole is excreted mainly as inactive metabolites in urine (35%) and in feces (54%) within one week of an oral solution dose. Renal excretion of itraconazole and the active metabolite hydroxy-itraconazole account for less than 1% of an intravenous dose.

Based on an oral radiolabeled dose, fecal excretion of unchanged drug ranges from 3% to 18% of the dose.

As re-distribution of itraconazole from keratinous tissues appears to be negligible, elimination of itraconazole from these tissues is related to epidermal regeneration. Contrary to plasma, the concentration in skin persists for 2 to 4 weeks after discontinuation of a 4-week treatment and in nail keratin – where itraconazole can be detected as early as 1 week after start of treatment – for at least six months after the end of a 3-month treatment period.

## **Special Populations**

### **Renal Impairment**

Limited data are available on the use of oral itraconazole in patients with renal impairment. A pharmacokinetic study using a single 200-mg oral dose of itraconazole was conducted in three groups of patients with renal impairment (uremia: n=7; hemodialysis: n=7; and continuous ambulatory peritoneal dialysis: n=5). In uremic subjects with a mean creatinine clearance of 13 mL/min.  $\times$  1.73 m<sup>2</sup>, the exposure, based on AUC, was slightly reduced compared with normal population parameters. This study did not demonstrate any significant effect of hemodialysis or continuous ambulatory peritoneal dialysis on the pharmacokinetics of itraconazole ( $T_{max}$ ,  $C_{max}$ , and  $AUC_{0-8h}$ ). Plasma concentration-versus-time profiles showed wide intersubject variation in all three groups. After a single intravenous dose, the mean terminal half-lives of itraconazole in patients with mild (defined in this study as CrCl 50 mL/min to 79 mL/min), moderate (defined in this study as CrCl 20 mL/min to 49 mL/min), and severe renal impairment (defined in this study as CrCl < 20 mL/min) were similar to that in healthy subjects (range of means 42 to 49 hours vs 48 hours in renally impaired patients and healthy subjects, respectively). Overall exposure to itraconazole, based on AUC, was decreased in patients with moderate and severe renal impairment by approximately 30% and 40%, respectively, as compared with subjects with normal renal function. Data are not available in renally impaired patients during long-term use of itraconazole. Dialysis has no effect on the half-life or clearance of itraconazole or hydroxy-itraconazole (see **PRECAUTIONS and DOSAGE AND ADMINISTRATION**).

### **Hepatic Impairment**

Itraconazole is predominantly metabolized in the liver. A pharmacokinetic study was conducted in 6 healthy and 12 cirrhotic subjects who were administered a single 100-mg dose of itraconazole as capsule. A statistically significant reduction in mean  $C_{max}$  (47%) and a twofold increase in the elimination half-life ( $37 \pm 17$  hours vs.  $16 \pm 5$  hours) of itraconazole were noted in cirrhotic subjects compared with healthy subjects. However, overall exposure to itraconazole, based on AUC, was similar in cirrhotic patients and in healthy subjects. Data are not available in cirrhotic patients during long-term use of itraconazole (see **CONTRAINDICATIONS, PRECAUTIONS: Drug Interactions and DOSAGE AND ADMINISTRATION**).

### **Decreased Cardiac Contractility**

When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later. If signs or symptoms of congestive heart failure appear during

administration of itraconazole capsules, itraconazole should be discontinued (see **BOXED WARNING, CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions and ADVERSE REACTIONS: Post-marketing Experience** for more information).

## **MICROBIOLOGY**

### **Mechanism of Action**

*In vitro* studies have demonstrated that itraconazole inhibits the cytochrome P450-dependent synthesis of ergosterol, which is a vital component of fungal cell membranes.

### **Antimicrobial Activity**

Itraconazole exhibits *in vitro* activity against *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Histoplasma duboisii*, *Aspergillus flavus*, *Aspergillus fumigatus*, and *Trichophyton* species (see **INDICATIONS AND USAGE: Description of Clinical Studies**).

### **Susceptibility Testing Methods**

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

### **Resistance**

Isolates from several fungal species with decreased susceptibility to itraconazole have been isolated *in vitro* and from patients receiving prolonged therapy.

Itraconazole is not active against *Zygomycetes* (e.g., *Rhizopus* spp., *Rhizomucor* spp., *Mucor* spp. and *Absidia* spp.), *Fusarium* spp., *Scedosporium* spp. and *Scopulariopsis* spp.

### **Cross-resistance**

Several *in vitro* studies have reported that some fungal clinical isolates with reduced susceptibility to one azole antifungal agent may also be less susceptible to other azole derivatives. The finding of cross-resistance is dependent on a number of factors, including the species evaluated, its clinical history, the particular azole compounds compared, and the type of susceptibility test that is performed.

Studies (both *in vitro* and *in vivo*) suggest that the activity of amphotericin B may be suppressed by prior azole antifungal therapy. As with other azoles, itraconazole inhibits the <sup>14</sup>C-demethylation step in the synthesis of ergosterol, a cell wall component of fungi. Ergosterol is the active site for amphotericin B. In one study the antifungal activity of amphotericin B against *Aspergillus fumigatus* infections in mice was inhibited by ketoconazole therapy. The clinical significance of test results obtained in this study is unknown.

## **INDICATIONS AND USAGE**

Itraconazole capsules are indicated for the treatment of the following fungal infections in immunocompromised and non-immunocompromised patients:

1. Blastomycosis, pulmonary and extrapulmonary
2. Histoplasmosis, including chronic cavitary pulmonary disease and disseminated, non-meningeal histoplasmosis, and
3. Aspergillosis, pulmonary and extrapulmonary, in patients who are intolerant of or who are refractory to amphotericin B therapy.

Specimens for fungal cultures and other relevant laboratory studies (wet mount, histopathology, serology) should be obtained before therapy to isolate and identify causative organisms. Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, antiinfective therapy should be adjusted accordingly.

Itraconazole capsules are also indicated for the treatment of the following fungal infections in non-immunocompromised patients:

1. Onychomycosis of the toenail, with or without fingernail involvement, due to dermatophytes (tinea unguium), and
2. Onychomycosis of the fingernail due to dermatophytes (tinea unguium).

Prior to initiating treatment, appropriate nail specimens for laboratory testing (KOH preparation, fungal culture, or nail biopsy) should be obtained to confirm the diagnosis of onychomycosis (see **CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, WARNINGS, and ADVERSE REACTIONS: Post-marketing Experience** for more information).

## **Description of Clinical Studies**

### **Blastomycosis**

Analyses were conducted on data from two open-label, non-concurrently controlled studies (N=73 combined) in patients with normal or abnormal immune status. The median dose was 200 mg/day. A response for most signs and symptoms was observed within the first 2 weeks, and all signs and symptoms cleared between 3 and 6 months. Results of these two studies demonstrated substantial evidence of the effectiveness of itraconazole for the treatment of blastomycosis compared with the natural history of untreated cases.

### **Histoplasmosis**

Analyses were conducted on data from two open-label, non-concurrently controlled studies (N=34 combined) in patients with normal or abnormal immune status (not including HIV-infected patients). The median dose was 200 mg/day. A response for most signs and symptoms was observed within the first 2 weeks, and all signs and symptoms cleared between 3 and 12 months. Results of these two studies demonstrated substantial evidence of the effectiveness of itraconazole for the treatment of histoplasmosis, compared with the natural history of untreated cases.

### **Histoplasmosis in HIV-infected patients**

Data from a small number of HIV-infected patients suggested that the response rate of histoplasmosis in HIV-infected patients is similar to that of non-HIV-infected patients. The clinical course of histoplasmosis in HIV-infected patients is more severe and usually requires maintenance therapy to prevent relapse.

### **Aspergillosis**

Analyses were conducted on data from an open-label, “single-patient-use” protocol designed to make itraconazole available in the U.S. for patients who either failed or were intolerant of amphotericin B therapy (N=190). The findings were corroborated by two smaller open-label studies (N=31 combined) in the same patient population. Most adult patients were treated with a daily dose of 200 mg to 400 mg, with a median duration of 3 months. Results of these studies demonstrated substantial evidence of effectiveness of itraconazole as a second-line therapy for the treatment of aspergillosis compared with the natural history of the disease in patients who either failed or were intolerant of amphotericin B therapy.

### **Onychomycosis of the toenail**

Analyses were conducted on data from three double-blind, placebo-controlled studies (N=214 total; 110 given itraconazole capsules) in which patients with onychomycosis of the toenails received 200 mg of itraconazole capsules once daily for 12 consecutive weeks. Results of these studies demonstrated mycologic cure, defined as simultaneous occurrence of negative KOH plus negative culture, in 54% of patients. Thirty-five percent (35%) of patients were considered an overall success (mycologic cure plus clear or minimal nail involvement with significantly decreased signs) and 14% of patients demonstrated mycologic cure plus clinical cure (clearance of all signs, with or without residual nail deformity). The mean time to overall success was approximately 10 months. Twenty-one percent (21%) of the overall success group had a relapse (worsening of the global score or conversion of KOH or culture from negative to positive).

### **Onychomycosis of the fingernail**

Analyses were conducted on data from a double-blind, placebo-controlled study (N=73 total; 37 given itraconazole capsules) in which patients with onychomycosis of the fingernails received a 1-week course (pulse) of 200 mg of itraconazole capsules b.i.d., followed by a 3-week period without itraconazole capsules, which was followed by a second 1-week pulse of 200 mg of itraconazole capsules b.i.d. Results demonstrated mycologic cure in 61% of patients. Fifty-six percent (56%) of patients were considered an overall success and 47% of patients demonstrated mycologic cure plus clinical cure. The mean time to overall success was approximately 5 months. None of the patients who achieved overall success relapsed.

## **CONTRAINDICATIONS**

### **Congestive Heart Failure**

Itraconazole capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF (see **BOXED WARNING, WARNINGS, PRECAUTIONS: Drug Interactions-Calcium Channel Blockers, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations**).

### **Drug Interactions**

Co-administration of a number of CYP3A4 substrates are contraindicated with itraconazole. Some examples of drugs for which plasma concentrations increase are: methadone, disopyramide, dofetilide, dronedarone, quinidine, isavuconazole, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam,



pimozide, triazolam, felodipine, nisoldipine, ivabradine, ranolazine, eplerenone, cisapride, naloxegol, lomitapide, lovastatin, simvastatin, avanafil, ticagrelor, finerenone, voclosporin. In addition, co-administration with colchicine, fesoterodine and solifenacin is contraindicated in subjects with varying degrees of renal or hepatic impairment, and co-administration with eliglustat is contraindicated in subjects that are poor or intermediate metabolizers of CYP2D6 and in subjects taking strong or moderate CYP2D6 inhibitors (see **PRECAUTIONS: Drug Interactions Section** for specific examples). This increase in drug concentrations caused by co-administration with itraconazole may increase or prolong both the pharmacologic effects and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of *torsade de pointes*, a potentially fatal arrhythmia. Some specific examples are listed in **PRECAUTIONS: Drug Interactions**.

Co-administration with venetoclax is contraindicated in patients with CLL/SLL during the dose initiation and ramp-up phase of venetoclax due to the potential for an increased risk of tumor lysis syndrome.

Itraconazole should not be administered for the treatment of onychomycosis to pregnant patients or to women contemplating pregnancy.

Itraconazole capsules are contraindicated for patients who have shown hypersensitivity to itraconazole. There is limited information regarding cross-hypersensitivity between itraconazole and other azole antifungal agents. Caution should be used when prescribing itraconazole capsules to patients with hypersensitivity to other azoles.

## **WARNINGS**

### **Hepatic Effects**

**Itraconazole has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition, and some of these cases developed within the first week of treatment. If clinical signs or symptoms develop that are consistent with liver disease, treatment should be discontinued and liver function testing performed. Continued itraconazole use or reinstatement of treatment with itraconazole is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk (see **PRECAUTIONS: Information for Patients and ADVERSE REACTIONS**).**

### **Cardiac Dysrhythmias**

Life-threatening cardiac dysrhythmias and/or sudden death have occurred in patients using drugs such as cisapride, pimozide, methadone, or quinidine concomitantly with itraconazole and/or other CYP3A4 inhibitors. Concomitant administration of these drugs with itraconazole is contraindicated (see **BOXED WARNING, CONTRAINDICATIONS, and PRECAUTIONS: Drug Interactions**).

### **Cardiac Disease**

**Itraconazole capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as**

**congestive heart failure (CHF) or a history of CHF.** Itraconazole capsules should not be used for other indications in patients with evidence of ventricular dysfunction unless the benefit clearly outweighs the risk.

For patients with risk factors for congestive heart failure, physicians should carefully review the risks and benefits of itraconazole therapy. These risk factors include cardiac disease such as ischemic and valvular disease; significant pulmonary disease such as chronic obstructive pulmonary disease; and renal failure and other edematous disorders. Such patients should be informed of the signs and symptoms of CHF, should be treated with caution, and should be monitored for signs and symptoms of CHF during treatment. If signs or symptoms of CHF appear during administration of itraconazole capsules, discontinue administration.

Itraconazole has been shown to have a negative inotropic effect. When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later.

Itraconazole has been associated with reports of congestive heart failure. In post-marketing experience, heart failure was more frequently reported in patients receiving a total daily dose of 400 mg although there were also cases reported among those receiving lower total daily doses.

Calcium channel blockers can have negative inotropic effects which may be additive to those of itraconazole. In addition, itraconazole can inhibit the metabolism of calcium channel blockers. Therefore, caution should be used when co-administering itraconazole and calcium channel blockers due to an increased risk of CHF. Concomitant administration of itraconazole and felodipine or nisoldipine is contraindicated.

Cases of CHF, peripheral edema, and pulmonary edema have been reported in the post-marketing period among patients being treated for onychomycosis and/or systemic fungal infections (see **CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, PRECAUTIONS: Drug Interactions, and ADVERSE REACTIONS: Post-marketing Experience** for more information).

### **Interaction potential**

Itraconazole has a potential for clinically important drug interactions. Co-administration of specific drugs with itraconazole may result in changes in efficacy of itraconazole and/or the co-administered drug, life-threatening effects and/or sudden death. Drugs that are contraindicated, not recommended or recommended for use with caution in combination with itraconazole are listed in **PRECAUTIONS: Drug Interactions**.

### **Interchangeability**

Itraconazole capsules and itraconazole oral solution should not be used interchangeably. This is because drug exposure is greater with the Oral Solution than with the Capsules when the same dose of drug is given. In addition, the topical effects of mucosal exposure may be different between the two formulations. Only the oral solution has been demonstrated effective for oral and/or esophageal candidiasis.

## PRECAUTIONS

### General

Itraconazole capsules should be administered after a full meal (see **CLINICAL PHARMACOLOGY: Pharmacokinetics and Metabolism**).

Under fasted conditions, itraconazole absorption was decreased in the presence of decreased gastric acidity. The absorption of itraconazole may be decreased with the concomitant administration of antacids or gastric acid secretion suppressors. Studies conducted under fasted conditions demonstrated that administration with 8 ounces of a non-diet cola beverage resulted in increased absorption of itraconazole in AIDS patients with relative or absolute achlorhydria. This increase relative to the effects of a full meal is unknown (see **CLINICAL PHARMACOLOGY: Pharmacokinetics and Metabolism**).

### Hepatotoxicity

Rare cases of serious hepatotoxicity have been observed with itraconazole treatment, including some cases within the first week. It is recommended that liver function monitoring be considered in all patients receiving itraconazole. Treatment should be stopped immediately and liver function testing should be conducted in patients who develop signs and symptoms suggestive of liver dysfunction.

### Neuropathy

If neuropathy occurs that may be attributable to itraconazole capsules, the treatment should be discontinued.

### Immunocompromised Patients

In some immunocompromised patients (e.g., neutropenic, AIDS or organ transplant patients), the oral bioavailability of itraconazole capsules may be decreased. Therefore, the dose should be adjusted based on the clinical response in these patients.

### Cystic Fibrosis

If a cystic fibrosis patient does not respond to itraconazole capsules, consideration should be given to switching to alternative therapy. For more information concerning the use of itraconazole in cystic fibrosis patients see the prescribing information for itraconazole oral solution.

### Hearing Loss

Transient or permanent hearing loss has been reported in patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (see **BOXED WARNING: Drug Interactions, CONTRAINDICATIONS: Drug Interactions and PRECAUTIONS: Drug Interactions**). The hearing loss usually resolves when treatment is stopped, but can persist in some patients.

### Information for Patients

- The topical effects of mucosal exposure may be different between the itraconazole capsules and oral solution. Only the oral solution has been demonstrated effective for oral and/or esophageal candidiasis. Itraconazole capsules should not be used interchangeably with itraconazole oral solution.
- Instruct patients to take itraconazole capsules with a full meal. Itraconazole capsules

must be swallowed whole.

- Instruct patients about the signs and symptoms of congestive heart failure, and if these signs or symptoms occur during itraconazole administration, they should discontinue itraconazole and contact their healthcare provider immediately.
- Instruct patients to stop itraconazole treatment immediately and contact their healthcare provider if any signs and symptoms suggestive of liver dysfunction develop. Such signs and symptoms may include unusual fatigue, anorexia, nausea and/or vomiting, jaundice, dark urine, or pale stools.
- Instruct patients to contact their physician before taking any concomitant medications with itraconazole to ensure there are no potential drug interactions.
- Instruct patients that hearing loss can occur with the use of itraconazole. The hearing loss usually resolves when treatment is stopped, but can persist in some patients. Advise patients to discontinue therapy and inform their physicians if any hearing loss symptoms occur.
- Instruct patients that dizziness or blurred/double vision can sometimes occur with itraconazole. Advise patients that if they experience these events, they should not drive or use machines.

## Drug Interactions

### Effect of Itraconazole on Other Drugs

Itraconazole and its major metabolite, hydroxy-itraconazole, are potent CYP3A4 inhibitors. Itraconazole is an inhibitor of the drug transporters P-glycoprotein and breast cancer resistance protein (BCRP). Consequently, itraconazole has the potential to interact with many concomitant drugs resulting in either increased or sometimes decreased concentrations of the concomitant drugs. Increased concentrations may increase the risk of adverse reactions associated with the concomitant drug which can be severe or life-threatening in some cases (e.g., QT prolongation, *Torsade de Pointes*, respiratory depression, hepatic adverse reactions, hypersensitivity reactions, myelosuppression, hypotension, seizures, angioedema, atrial fibrillation, bradycardia, priapism). Reduced concentrations of concomitant drugs may reduce their efficacy. Table 1 lists examples of drugs that may have their concentrations affected by itraconazole, but it is not a comprehensive list.

Refer to the approved product labeling to become familiar with the interaction pathways, risk potential, and specific actions to be taken with regards to each concomitant drug prior to initiating therapy with itraconazole.

Although many of the clinical drug interactions in Table 1 are based on information with a similar azole antifungal, ketoconazole, these interactions are expected to occur with itraconazole.

### Table 1: Drug Interactions with Itraconazole that Affect Concomitant Drug Concentrations

Examples of Concomitant Drugs Within Class	Prevention or Management
--	--------------------------

**Drug Interactions with Itraconazole that Increase Concomitant Drug Concentrations and May Increase Risk of Adverse Reactions Associated with the Concomitant Drug****Alpha Blockers**

Alfuzosin Silodosin Tamsulosin	Not recommended during and 2 weeks after itraconazole treatment.
--------------------------------------	--

**Analgesics**

Methadone	Contraindicated during and 2 weeks after itraconazole treatment.
Fentanyl	Not recommended during and 2 weeks after itraconazole treatment.
Alfentanil Buprenorphine (IV and sublingual) Oxycodone <sup>a</sup> Sufentanil	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.

**Antiarrhythmics**

Disopyramide Dofetilide Dronedarone Quinidine <sup>a</sup>	Contraindicated during and 2 weeks after itraconazole treatment.
Digoxin <sup>a</sup>	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.

**Antibacterials**

Bedaquiline <sup>b</sup>	Concomitant itraconazole not recommended for more than 2 weeks at any time during bedaquiline treatment.
Rifabutin	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment. See also Table 2.
Clarithromycin	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary. See also Table 2.
Trimetrexate	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.

**Anticoagulants and Antiplatelets**

Ticagrelor	Contraindicated during and 2 weeks after itraconazole treatment.
Apixaban Rivaroxaban Vorapaxar	Not recommended during and 2 weeks after itraconazole treatment.
Cilostazol Dabigatran Warfarin	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.

**Anticonvulsants**

Carbamazepine	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment. See also Table 2.
---------------	---

**Antidiabetic Drugs**

Repaglinide <sup>a</sup> Saxagliptin	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.	
<b>Anthelmintics, Antifungals and Antiprotozoals</b>		
Isavuconazonium	Contraindicated during and 2 weeks after itraconazole treatment.	
Praziquantel	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.	
Artemether-lumefantrine Quinine <sup>a</sup>	Monitor for adverse reactions.	
<b>Antimigraine Drugs</b>		
Ergot alkaloids (e.g., dihydroergotamine, ergotamine)	Contraindicated during and 2 weeks after itraconazole treatment.	
Eletriptan	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.	
<b>Antineoplastics</b>		
Irinotecan	Contraindicated during and 2 weeks after itraconazole treatment.	
Venetoclax	Contraindicated during the dose initiation and ramp-up phase in patients with CLL/SLL. Refer to the venetoclax prescribing information for dosing and safety monitoring instructions.	
Mobocertinib <sup>a</sup>	Avoid use during and 2 weeks after itraconazole treatment.	
Axitinib Bosutinib Cabazitaxel Cabozantinib Ceritinib Cobimetinib <sup>a</sup> Crizotinib Dabrafenib Dasatinib	Docetaxel Ibrutinib Lapatinib Nilotinib Olaparib <sup>a</sup> Pazopanib Sunitinib Trabectedin Trastuzumab- emtansine Vinca alkaloids	Avoid use during and 2 weeks after itraconazole treatment.
Entrectinib <sup>a</sup> Pemigatinib <sup>a</sup> Talazoparib <sup>a</sup>	Refer to the entrectinib, pemigatinib and talazoparib prescribing information for dosing instructions if concomitant use cannot be avoided.	
Glasdegib	Refer to the glasdegib prescribing information for safety monitoring if concomitant use cannot be avoided.	
Bortezomib Brentuximab- vedotin Busulfan <sup>a</sup> Erlotinib Gefitinib <sup>a</sup> Idelalisib Imatinib Ixabepilone	Nintedanib Panobinostat Ponatinib Ruxolitinib Sonidegib Tretinoin (oral) Vandetanib <sup>a</sup>	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary. For idelalisib, see also Table 2.

<b>Antipsychotics, Anxiolytics and Hypnotics</b>		
Alprazolam <sup>a</sup> Aripiprazole <sup>a</sup> Buspirone <sup>a</sup> Cariprazine Diazepam <sup>a</sup> Haloperidol <sup>a</sup>	Midazolam (IV) <sup>a</sup> Quetiapine Ramelteon Risperidone <sup>a</sup> Suvorexant	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
Zopiclone <sup>a</sup>		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
Lurasidone Midazolam (oral) <sup>a</sup> Pimozide Triazolam <sup>a</sup>		Contraindicated during and 2 weeks after itraconazole treatment.
<b>Antivirals</b>		
Daclatasvir Indinavir <sup>a</sup> Maraviroc		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary. For indinavir, see also Table 2.
Cobicistat Elvitegravir (ritonavir-boosted) Ombitasvir/Paritaprevir/Ritonavir with or without Dasabuvir Ritonavir Saquinavir (unboosted) <sup>a</sup>		Monitor for adverse reactions. See also Table 2.
Elbasvir/grazoprevir Glecaprevir/pibrentasvir Tenofovir disoproxil fumarate		Not recommended during and 2 weeks after itraconazole treatment. Monitor for adverse reactions. Monitor for adverse reactions.
<b>Beta Blockers</b>		
Nadolol <sup>a</sup>		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
<b>Calcium Channel Blockers</b>		
Felodipine <sup>a</sup> Nisoldipine		Contraindicated during and 2 weeks after itraconazole treatment.
Diltiazem Other dihydropyridines Verapamil		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary. For diltiazem, see also Table 2.
<b>Cardiovascular Drugs, Miscellaneous</b>		
Ivabradine Ranolazine		Contraindicated during and 2 weeks after itraconazole treatment.
Aliskiren <sup>a</sup> Riociguat Sildenafil (for pulmonary hypertension) Tadalafil (for pulmonary hypertension)		Not recommended during and 2 weeks after itraconazole treatment. For sildenafil and tadalafil, see also Urologic Drugs below.
Bosentan Guanfacine		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
<b>Contraceptives*</b>		

Dienogest Ulipristal		Monitor for adverse reactions.
<b>Diuretics</b>		
Eplerenone Finerenone		Contraindicated during and 2 weeks after itraconazole treatment.
<b>Gastrointestinal Drugs</b>		
Cisapride Naloxegol		Contraindicated during and 2 weeks after itraconazole treatment.
Aprepitant Loperamide <sup>a</sup>		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
Netupitant		Monitor for adverse reactions.
<b>Immunosuppressants</b>		
Voclosporin		Contraindicated during and for 2 weeks after itraconazole treatment.
Everolimus Sirolimus Temsilimus (IV)		Not recommended during and 2 weeks after itraconazole treatment.
Budesonide (inhalation) <sup>a</sup> Budesonide (non-inhalation) Ciclesonide (inhalation) Cyclosporine (IV) <sup>a</sup> Cyclosporine (non-IV) Dexamethasone <sup>a</sup>	Fluticasone (inhalation) <sup>a</sup> Fluticasone (nasal) Methylprednisolone <sup>a</sup> Tacrolimus (IV) <sup>a</sup> Tacrolimus (oral)	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
<b>Lipid-Lowering Drugs</b>		
Lomitapide Lovastatin <sup>a</sup> Simvastatin <sup>a</sup>		Contraindicated during and 2 weeks after itraconazole treatment.
Atorvastatin <sup>a</sup>		Monitor for drug adverse reactions. Concomitant drug dose reduction may be necessary.
<b>Respiratory Drugs</b>		
Salmeterol		Not recommended during and 2 weeks after itraconazole treatment.
<b>SSRIs, Tricyclics and Related Antidepressants</b>		
Venlafaxine		Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
<b>Urologic Drugs</b>		
Avanafil		Contraindicated during and 2 weeks after itraconazole treatment.
Fesoterodine		<i>Patients with moderate to severe renal or hepatic impairment:</i> Contraindicated during and 2 weeks after itraconazole treatment. <i>Other patients:</i> Monitor for adverse reactions. Concomitant drug dose reduction may be



	necessary.
Solifenacin	<i>Patients with severe renal or moderate to severe hepatic impairment:</i> Contraindicated during and 2 weeks after itraconazole treatment. <i>Other patients:</i> Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
Darifenacin Vardenafil	Not recommended during and 2 weeks after itraconazole treatment.
Dutasteride Oxybutynin <sup>a</sup> Sildenafil (for erectile dysfunction) Tadalafil (for erectile dysfunction and benign prostatic hyperplasia) Tolterodine	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary. For sildenafil and tadalafil, see also Cardiovascular Drugs above.
<b>Miscellaneous Drugs and Other Substances</b>	
Colchicine	<i>Patients with renal or hepatic impairment:</i> Contraindicated during and 2 weeks after itraconazole treatment. <i>Other patients:</i> Not recommended during and 2 weeks after itraconazole treatment.
Eliglustat	<i>CYP2D6 EMs<sup>c</sup> taking a strong or moderate CYP2D6 inhibitor, CYP2D6 IMs<sup>c</sup>, or CYP2D6 PMs<sup>c</sup>:</i> Contraindicated during and 2 weeks after itraconazole treatment. <i>CYP2D6 EMs<sup>c</sup> not taking a strong or moderate CYP2D6 inhibitor:</i> Monitor for adverse reactions. Eliglustat dose reduction may be necessary.
Lumacaftor/Ivacaftor	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment.
Alitretinoin (oral) Cabergoline Cannabinoids Cinacalcet Galantamine Ivacaftor	Monitor for adverse reactions. Concomitant drug dose reduction may be necessary.
Valbenazine	Concomitant drug dose reduction is necessary. Refer to the valbenazine prescribing information for dosing instructions.
<b>Vasopressin Receptor Antagonists</b>	
Conivaptan Tolvaptan	Not recommended during and 2 weeks after itraconazole treatment.
<b>Drug Interactions with Itraconazole that Decrease Concomitant Drug Concentrations and May Reduce Efficacy of the Concomitant Drug</b>	
<b>Antineoplastics</b>	
Regorafenib	Not recommended during and 2 weeks after itraconazole treatment.
<b>Gastrointestinal Drugs</b>	
<i>Saccharomyces boulardii</i>	Not recommended during and 2 weeks after

<i>Saccharomyces boulardii</i>	itraconazole treatment.
<b>Nonsteroidal Anti-Inflammatory Drugs</b>	
Meloxicam <sup>a</sup>	Concomitant drug dose increase may be necessary.
*CYP3A4 inhibitors (including itraconazole) may increase systemic contraceptive hormone concentrations.	
<sup>a</sup> Based on clinical drug interaction information with itraconazole.	
<sup>b</sup> Based on 400 mg Bedaquiline once daily for 2 weeks.	
<sup>c</sup> EMs: extensive metabolizers; IMs: intermediate metabolizers, PMs: poor metabolizers	

## Effect of Other Drugs on Itraconazole

Itraconazole is mainly metabolized through CYP3A4. Other substances that either share this metabolic pathway or modify CYP3A4 activity may influence the pharmacokinetics of itraconazole. Some concomitant drugs have the potential to interact with itraconazole resulting in either increased or sometimes decreased concentrations of itraconazole. Increased concentrations may increase the risk of adverse reactions associated with itraconazole. Decreased concentrations may reduce itraconazole efficacy.

Table 2 lists examples of drugs that may affect itraconazole concentrations, but is not a comprehensive list. Refer to the approved product labeling to become familiar with the interaction pathways, risk potential and specific actions to be taken with regards to each concomitant drug prior to initiating therapy with itraconazole.

Although many of the clinical drug interactions in Table 2 are based on information with a similar azole antifungal, ketoconazole, these interactions are expected to occur with itraconazole.

**Table 2: Drug Interactions with Other Drugs that Affect Itraconazole Concentrations**

<b>Examples of Concomitant Drugs Within Class</b>	<b>Prevention or Management</b>
<b>Drug Interactions with Other Drugs that Increase Itraconazole Concentrations and May Increase Risk of Adverse Reactions Associated with Itraconazole</b>	
<b>Antibacterials</b>	
Ciprofloxacin <sup>a</sup> Erythromycin <sup>a</sup> Clarithromycin <sup>a</sup>	Monitor for adverse reactions. Itraconazole dose reduction may be necessary.
<b>Antineoplastics</b>	
Idelalisib	Monitor for adverse reactions. Itraconazole dose reduction may be necessary. See also Table 1.
<b>Antivirals</b>	
Cobicistat Darunavir (ritonavir-boosted) Elvitegravir (ritonavir-boosted) Fosamprenavir (ritonavir-	Monitor for adverse reactions. Itraconazole dose

boosted) Indinavir <sup>a</sup> Ombitasvir/Paritaprevir/Ritonavir with or without Dasabuvir Ritonavir Saqinavir	reduction may be necessary. For, cobicistat, elvitegravir, indinavir, ombitasvir/paritaprevir/ritonavir with or without dasabuvir, ritonavir, and saquinavir, see also Table 1.
<b>Calcium Channel Blockers</b>	
Diltiazem	Monitor for adverse reactions. Itraconazole dose reduction may be necessary. See also Table 1.
<b>Drug Interactions with Other Drugs that Decrease Itraconazole Concentrations and May Reduce Efficacy of Itraconazole</b>	
<b>Antibacterials</b>	
Isoniazid Rifampicin <sup>a</sup>	Not recommended 2 weeks before and during itraconazole treatment.
Rifabutin <sup>a</sup>	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment. See also Table 1.
<b>Anticonvulsants</b>	
Phenobarbital Phenytoin <sup>a</sup>	Not recommended 2 weeks before and during itraconazole treatment.
Carbamazepine	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment. See also Table 1.
<b>Antivirals</b>	
Efavirenz <sup>a</sup> Nevirapine <sup>a</sup>	Not recommended 2 weeks before and during itraconazole treatment.
<b>Gastrointestinal Drugs</b>	
Drugs that reduce gastric acidity e.g., acid neutralizing medicines such as aluminum hydroxide, or acid secretion suppressors such as H <sub>2</sub> - receptor antagonists and proton pump inhibitors.	Use with caution. Administer acid neutralizing medicines at least 2 hours before or 2 hours after the intake of itraconazole capsules
<b>Miscellaneous Drugs and Other Substances</b>	
Lumacaftor/Ivacaftor	Not recommended 2 weeks before, during, and 2 weeks after itraconazole treatment.
<sup>a</sup> Based on clinical drug interaction information with itraconazole.	

## Pediatric Population

Interaction studies have only been performed in adults.

## Carcinogenesis, Mutagenesis, and Impairment of Fertility

Itraconazole showed no evidence of carcinogenicity potential in mice treated orally for 23 months at dosage levels up to 80 mg/kg/day (approximately 1 time the maximum recommended human dose [MRHD] of 400 mg/day based on body surface area comparisons). Male rats treated with 25 mg/kg/day (0.6 times the MRHD based on body surface area comparisons) had a slightly increased incidence of soft tissue sarcoma. These sarcomas may have been a consequence of hypercholesterolemia, which is a

response of rats, but not dogs or humans, to chronic itraconazole administration. Female rats treated with 50 mg/kg/day (1.2 times the MRHD based on body surface area comparisons) had an increased incidence of squamous cell carcinoma of the lung (2/50) as compared to the untreated group. Although the occurrence of squamous cell carcinoma in the lung is extremely uncommon in untreated rats, the increase in this study was not statistically significant.

Itraconazole produced no mutagenic effects when assayed in DNA repair test (unscheduled DNA synthesis) in primary rat hepatocytes, in Ames tests with *Salmonella typhimurium* (6 strains) and *Escherichia coli*, in the mouse lymphoma gene mutation tests, in a sex-linked recessive lethal mutation (*Drosophila melanogaster*) test, in chromosome aberration tests in human lymphocytes, in a cell transformation test with C3H/10T $\frac{1}{2}$ ; C18 mouse embryo fibroblasts cells, in a dominant lethal mutation test in male and female mice, and in micronucleus tests in mice and rats.

Itraconazole did not affect the fertility of male or female rats treated orally with dosage levels of up to 40 mg/kg/day (1 time the MRHD based on body surface area comparisons), even though parental toxicity was present at this dosage level. More severe signs of parental toxicity, including death, were present in the next higher dosage level, 160 mg/kg/day (4 times the MRHD based on body surface area comparisons).

### **Pregnancy: Teratogenic effects**

Itraconazole was found to cause a dose-related increase in maternal toxicity, embryotoxicity, and teratogenicity in rats at dosage levels of approximately 40 mg/kg/day to 160 mg/kg/day (1 to 4 times the MRHD based on body surface area comparisons), and in mice at dosage levels of approximately 80 mg/kg/day (1 time the MRHD based on body surface area comparisons). Itraconazole has been shown to cross the placenta in a rat model. In rats, the teratogenicity consisted of major skeletal defects; in mice, it consisted of encephaloceles and/or macroglossia.

There are no studies in pregnant women. Itraconazole should be used for the treatment of systemic fungal infections in pregnancy only if the benefit outweighs the potential risk.

Itraconazole should not be administered for the treatment of onychomycosis to pregnant patients or to women contemplating pregnancy. Itraconazole should not be administered to women of childbearing potential for the treatment of onychomycosis unless they are using effective measures to prevent pregnancy and they begin therapy on the second or third day following the onset of menses. Highly effective contraception should be continued throughout itraconazole therapy and for 2 months following the end of treatment.

During post-marketing experience, cases of congenital abnormalities have been reported (see **ADVERSE REACTIONS: Post-marketing Experience**).

### **Nursing Mothers**

Itraconazole is excreted in human milk; therefore, the expected benefits of itraconazole therapy for the mother should be weighed against the potential risk from exposure of itraconazole to the infant. The U.S. Public Health Service Centers for Disease Control and Prevention advises HIV-infected women not to breast-feed to avoid potential transmission of HIV to uninfected infants.

## Pediatric Use

The efficacy and safety of itraconazole have not been established in pediatric patients.

The long-term effects of itraconazole on bone growth in children are unknown. In three toxicology studies using rats, itraconazole induced bone defects at dosage levels as low as 20 mg/kg/day (0.5 times the MRHD of 400 mg based on body surface area comparisons). The induced defects included reduced bone plate activity, thinning of the zona compacta of the large bones, and increased bone fragility. At a dosage level of 80 mg/kg/day (2 times the MRHD based on body surface area comparisons) over 1 year or 160 mg/kg/day (4 times the MRHD based on body surface area comparisons) for 6 months, itraconazole induced small tooth pulp with hypocellular appearance in some rats.

## Geriatric Use

Clinical studies of itraconazole capsules did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. It is advised to use itraconazole capsules in these patients only if it is determined that the potential benefit outweighs the potential risks. In general, it is recommended that the dose selection for an elderly patient should be taken into consideration, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Transient or permanent hearing loss has been reported in elderly patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (see **BOXED WARNING: Drug Interactions, CONTRAINDICATIONS: Drug Interactions and PRECAUTIONS: Drug Interactions**).

## HIV-Infected Patients

Because hypochlorhydria has been reported in HIV-infected individuals, the absorption of itraconazole in these patients may be decreased.

## Renal Impairment

Limited data are available on the use of oral itraconazole in patients with renal impairment. The exposure of itraconazole may be lower in some patients with renal impairment. Caution should be exercised when itraconazole is administered in this patient population and dose adjustment may be needed (see **CLINICAL PHARMACOLOGY: Special Populations and DOSAGE AND ADMINISTRATION**).

## Hepatic Impairment

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population. It is recommended that patients with impaired hepatic function be carefully monitored when taking itraconazole. It is recommended that the prolonged elimination half-life of itraconazole observed in the single oral dose clinical trial with itraconazole capsules in cirrhotic patients be considered when deciding to initiate therapy with other medications metabolized by CYP3A4.

In patients with elevated or abnormal liver enzymes or active liver disease, or who have experienced liver toxicity with other drugs, treatment with itraconazole is strongly discouraged unless there is a serious or life-threatening situation where the expected

benefit exceeds the risk. It is recommended that liver function monitoring be done in patients with pre-existing hepatic function abnormalities or those who have experienced liver toxicity with other medications (see **CLINICAL PHARMACOLOGY: Special Populations and DOSAGE AND ADMINISTRATION**).

## ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Itraconazole has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition. If clinical signs or symptoms develop that are consistent with liver disease, treatment should be discontinued and liver function testing performed. The risks and benefits of itraconazole use should be reassessed (see **WARNINGS: Hepatic Effects** and **PRECAUTIONS: Hepatotoxicity and Information for Patients**).

### Adverse Events in the Treatment of Systemic Fungal Infections

Adverse event data were derived from 602 patients treated for systemic fungal disease in U.S. clinical trials who were immunocompromised or receiving multiple concomitant medications. Treatment was discontinued in 10.5% of patients due to adverse events. The median duration before discontinuation of therapy was 81 days (range: 2 to 776 days). The table lists adverse events reported by at least 1% of patients.

**Table 3: Clinical Trials of Systemic Fungal Infections: Adverse Events Occurring with an Incidence of Greater than or Equal to 1%**

Body System/Adverse Event	Incidence(%) (N=602)
<b>Gastrointestinal</b>	
Nausea	11
Vomiting	5
Diarrhea	3
Abdominal Pain	2
Anorexia	1
<b>Body as a Whole</b>	
Edema	4
Fatigue	3
Fever	3
Malaise	1
<b>Skin and Appendages</b>	
Rash*	9
Pruritus	3
<b>Central/Peripheral Nervous System</b>	
Headache	4
Dizziness	2
<b>Psychiatric</b>	

Libido Decreased	1
Somnolence	1
<b>Cardiovascular</b>	
Hypertension	3
<b>Metabolic/Nutritional</b>	
Hypokalemia	2
<b>Urinary System</b>	
Albuminuria	1
<b>Liver and Biliary System</b>	
Hepatic Function Abnormal	3
<b>Reproductive System, Male</b>	
Impotence	1
* Rash tends to occur more frequently in immunocompromised patients receiving immunosuppressive medications.	

Adverse events infrequently reported in all studies included constipation, gastritis, depression, insomnia, tinnitus, menstrual disorder, adrenal insufficiency, gynecomastia, and male breast pain.

#### **Adverse Events Reported in Toenail Onychomycosis Clinical Trials**

Patients in these trials were on a continuous dosing regimen of 200 mg once daily for 12 consecutive weeks.

The following adverse events led to temporary or permanent discontinuation of therapy.

**Table 4: Clinical Trials of Onychomycosis of the Toenail: Adverse Events Leading to Temporary or Permanent Discontinuation of Therapy**

<b>Adverse Event</b>	<b>Incidence (%) Itraconazole (N=112)</b>
Elevated Liver Enzymes (greater than twice the upper limit of normal)	4
Gastrointestinal Disorders	4
Rash	3
Hypertension	2
Orthostatic Hypotension	1
Headache	1
Malaise	1
Myalgia	1
Vasculitis	1
Vertigo	1

The following adverse events occurred with an incidence of greater than or equal to 1% (N=112): headache: 10%; rhinitis: 9%; upper respiratory tract infection: 8%; sinusitis, injury: 7%; diarrhea, dyspepsia, flatulence, abdominal pain, dizziness, rash: 4%; cystitis, urinary tract infection, liver function abnormality, myalgia, nausea: 3%; appetite increased, constipation, gastritis, gastroenteritis, pharyngitis, asthenia, fever, pain, tremor, herpes zoster, abnormal dreaming: 2%.

## Adverse Events Reported in Fingernail Onychomycosis Clinical Trials

Patients in these trials were on a pulse regimen consisting of two 1-week treatment periods of 200 mg twice daily, separated by a 3-week period without drug.

The following adverse events led to temporary or permanent discontinuation of therapy.

**Table 5: Clinical Trials of Onychomycosis of the Fingernail: Adverse Events Leading to Temporary or Permanent Discontinuation of Therapy**

Adverse Event	Incidence (%) Itraconazole (N=37)
Rash/Pruritus	3
Hypertriglyceridemia	3

The following adverse events occurred with an incidence of greater than or equal to 1% (N=37): headache: 8%; pruritus, nausea, rhinitis: 5%; rash, bursitis, anxiety, depression, constipation, abdominal pain, dyspepsia, ulcerative stomatitis, gingivitis, hypertriglyceridemia, sinusitis, fatigue, malaise, pain, injury: 3%.

## Adverse Events Reported from Other Clinical Trials

In addition, the following adverse drug reaction was reported in patients who participated in itraconazole capsules clinical trials: *Hepatobiliary Disorders*: hyperbilirubinemia.

The following is a list of additional adverse drug reactions associated with itraconazole that have been reported in clinical trials of itraconazole oral solution and itraconazole IV excluding the adverse reaction term "Injection site inflammation" which is specific to the injection route of administration:

*Cardiac Disorders*: cardiac failure, left ventricular failure, tachycardia;

*General Disorders and Administration Site Conditions*: face edema, chest pain, chills;

*Hepatobiliary Disorders*: hepatic failure, jaundice;

*Investigations*: alanine aminotransferase increased, aspartate aminotransferase increased, blood alkaline phosphatase increased, blood lactate dehydrogenase increased, blood urea increased, gamma-glutamyltransferase increased, urine analysis abnormal;

*Metabolism and Nutrition Disorders*: hyperglycemia, hyperkalemia, hypomagnesemia;

*Psychiatric Disorders*: confusional state;

*Renal and Urinary Disorders*: renal impairment;

*Respiratory, Thoracic and Mediastinal Disorders*: dysphonia, cough;

*Skin and Subcutaneous Tissue Disorders*: rash erythematous, hyperhidrosis;

*Vascular Disorders*: hypotension

## Post-marketing Experience

Adverse drug reactions that have been first identified during post-marketing experience with itraconazole (all formulations) are listed in the table below. Because these reactions



are reported voluntarily from a population of uncertain size, reliably estimating their frequency or establishing a causal relationship to drug exposure is not always possible.

**Table 6: Post-marketing Reports of Adverse Drug Reactions**

<b>Blood and Lymphatic System Disorders:</b>	Leukopenia, neutropenia, thrombocytopenia
<b>Immune System Disorders:</b>	Anaphylaxis; anaphylactic, anaphylactoid and allergic reactions; serum sickness; angioneurotic edema
<b>Nervous System Disorders:</b>	Peripheral neuropathy, paresthesia, hypoesthesia, tremor
<b>Eye Disorders:</b>	Visual disturbances, including vision blurred and diplopia
<b>Ear and Labyrinth Disorders:</b>	Transient or permanent hearing loss
<b>Cardiac Disorders:</b>	Congestive heart failure
<b>Respiratory, Thoracic and Mediastinal Disorders:</b>	Pulmonary edema, dyspnea
<b>Gastrointestinal Disorders:</b>	Pancreatitis, dysgeusia
<b>Hepatobiliary Disorders:</b>	Serious hepatotoxicity (including some cases of fatal acute liver failure), hepatitis
<b>Skin and Subcutaneous Tissue Disorders:</b>	Toxic epidermal necrolysis, Stevens-Johnson syndrome, acute generalized exanthematous pustulosis, erythema multiforme, exfoliative dermatitis, leukocytoclastic vasculitis, alopecia, photosensitivity, urticaria
<b>Musculoskeletal and Connective Tissue Disorders:</b>	Arthralgia
<b>Renal and Urinary Disorders:</b>	Urinary incontinence, pollakiuria
<b>Reproductive System and Breast Disorders:</b>	Erectile dysfunction
<b>General Disorders and Administration Site Conditions:</b>	Peripheral edema
<b>Investigations:</b>	Blood creatine phosphokinase increased

There is limited information on the use of itraconazole during pregnancy. Cases of congenital abnormalities including skeletal, genitourinary tract, cardiovascular and ophthalmic malformations as well as chromosomal and multiple malformations have been reported during post-marketing experience. A causal relationship with itraconazole has not been established (see **CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS: Drug Interactions** for more information).

**To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **OVERDOSAGE**

Itraconazole is not removed by dialysis. In the event of accidental overdose, supportive measures should be employed. Contact a certified poison control center for the most up-to-date information on the management of itraconazole capsules overdose (1-800-222-1222 or [www.poison.org](http://www.poison.org)).

In general, adverse events reported with overdose have been consistent with adverse drug reactions already listed in this package insert for itraconazole (see **ADVERSE REACTIONS**).

## **DOSAGE AND ADMINISTRATION**

Itraconazole capsules should be taken with a full meal to ensure maximal absorption. Itraconazole capsules must be swallowed whole.

Itraconazole capsules are a different preparation than itraconazole oral solution and should not be used interchangeably.

### **Treatment of Blastomycosis and Histoplasmosis**

The recommended dose is 200 mg once daily (2 capsules). If there is no obvious improvement, or there is evidence of progressive fungal disease, the dose should be increased in 100-mg increments to a maximum of 400 mg daily. Doses above 200 mg/day should be given in two divided doses.

### **Treatment of Aspergillosis**

A daily dose of 200 to 400 mg is recommended.

### **Treatment in Life-Threatening Situations**

In life-threatening situations, a loading dose should be used.

Although clinical studies did not provide for a loading dose, it is recommended, based on pharmacokinetic data, that a loading dose of 200 mg (2 capsules) three times daily (600 mg/day) be given for the first 3 days of treatment.

Treatment should be continued for a minimum of three months and until clinical parameters and laboratory tests indicate that the active fungal infection has subsided. An inadequate period of treatment may lead to recurrence of active infection.

Itraconazole capsules and itraconazole oral solution should not be used interchangeably. Only the oral solution has been demonstrated effective for oral and/or esophageal candidiasis.

### **Treatment of Onychomycosis**

Toenails with or without fingernail involvement: The recommended dose is 200 mg (2 capsules) once daily for 12 consecutive weeks.

### **Treatment of Onychomycosis**

Fingernails only: The recommended dosing regimen is 2 treatment pulses, each consisting of 200 mg (2 capsules) b.i.d. (400 mg/day) for 1 week. The pulses are separated by a 3-week period without itraconazole capsules.

### **Use in Patients with Renal Impairment**

Limited data are available on the use of oral itraconazole in patients with renal

impairment. Caution should be exercised when this drug is administered in this patient population (see **CLINICAL PHARMACOLOGY: Special Populations and PRECAUTIONS**).

### **Use in Patients with Hepatic Impairment**

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population (see **CLINICAL PHARMACOLOGY: Special Populations, WARNINGS, and PRECAUTIONS**).

### **HOW SUPPLIED**

Itraconazole capsules USP, **100 mg**, are supplied as a hard gelatin capsule filled with white to off-white pellets having “AMNEAL” printed on blue opaque cap and “630” on blue transparent body in black ink.

They are available as follows:

Bottles of 30: NDC 65162-630-03

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture.

Keep out of reach of children.

Manufactured by:

**Amneal Pharmaceuticals Pvt. Ltd.**  
Ahmedabad 382220, INDIA

Distributed by:

**Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Rev. 05-2024-07

### **PATIENT INFORMATION**

#### **PATIENT INFORMATION**

#### **Itraconazole (it"ra kon'a zole) Capsules, USP**

Read this Patient Information that comes with itraconazole capsules before you start taking it and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

#### **What is the most important information I should know about itraconazole capsules?**

**Itraconazole capsules can cause serious side effects, including:**

**1. Heart failure.** Do not take itraconazole capsules if you have had heart failure, including congestive heart failure.

**Stop taking itraconazole capsules and call your healthcare provider right away if you have any of these symptoms of congestive heart failure:**

- shortness of

breath <ul style="list-style-type: none"> <li>• swelling of your feet, ankles or legs</li> <li>• sudden weight gain</li> <li>• increased tiredness</li> </ul>	<ul style="list-style-type: none"> <li>• coughing up white or pink mucus (phlegm)</li> <li>• fast heartbeat</li> <li>• waking up at night more than normal for you</li> </ul>
--	---

**2. Heart problems and other serious medical problems.** Serious medical problems that affect the heart and other parts of your body can happen if you take itraconazole capsules with certain other medicines. **Do not take itraconazole capsules if you also take the following medicines:**

<ul style="list-style-type: none"> <li>• methadone</li> <li>• disopyramide</li> <li>• dofetilide</li> <li>• dronedarone</li> <li>• quinidine</li> <li>• isavuconazole</li> <li>• ergot alkaloids (such as dihydroergotamine, ergometrine ergonovine)</li> <li>• ergotamine</li> </ul>	<ul style="list-style-type: none"> <li>• methylergometrine (methylergonovine)</li> <li>• irinotecan</li> <li>• lurasidone</li> <li>• oral midazolam</li> <li>• pimozide</li> <li>• triazolam</li> <li>• felodipine</li> <li>• nisoldipine</li> <li>• ivabradine</li> </ul>	<ul style="list-style-type: none"> <li>• ranolazine</li> <li>• eplerenone</li> <li>• cisapride</li> <li>• naloxegol</li> <li>• lomitapide</li> <li>• lovastatin</li> <li>• simvastatin</li> <li>• avanafil</li> <li>• ticagrelor</li> <li>• venetoclax (see below)</li> <li>• finerenone</li> <li>• voclosporin</li> </ul>
---	--	--

**Do not** take itraconazole capsules with venetoclax for chronic lymphocytic leukemia/small lymphocytic lymphoma when you first start treatment with venetoclax or with increasing doses of venetoclax.

This is not a complete list of medicines that can interact with itraconazole capsules. Itraconazole capsules may affect the way other medicines work, and other medicines may affect how itraconazole capsules works. You can ask your pharmacist for a list of medicines that interact with itraconazole capsules.

Before you start taking itraconazole capsules, tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Before you start any new medicine, ask your healthcare provider or pharmacist if it is safe to take it with itraconazole capsules.

**3. Liver problems.** Itraconazole capsules can cause serious liver problems which may be severe and lead to death. **Stop taking itraconazole capsules and call your healthcare provider right away if you have any of these symptoms of liver problems:**

For more information about side effects, see **“What are the possible side effects of itraconazole capsules?”**

<ul style="list-style-type: none"> <li>• tiredness</li> <li>• loss of appetite for several days or longer</li> <li>• nausea or vomiting</li> <li>• dark or “tea-colored” urine</li> </ul>	<ul style="list-style-type: none"> <li>• your skin or the white part of your eyes turn yellow (jaundice)</li> <li>• light-colored stools (bowel movement)</li> </ul>
---	--

**What are itraconazole capsules?**

- Itraconazole capsules are a prescription medicine used to treat the following fungal infections of the toenails, fingernails and other parts of the body: blastomycosis, histoplasmosis, aspergillosis, and onychomycosis.
- It is not known if itraconazole capsules are safe and effective in children.

**Do not take itraconazole capsules if you:**

- have or have had heart failure, including congestive heart failure.
- take certain medicines. See “**What is the most important information I should know about itraconazole capsules?**”
- are pregnant or plan to become pregnant. Itraconazole capsules can harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking itraconazole capsules. Females who are able to become pregnant must use effective forms of birth control during treatment and for 2 months after stopping treatment with itraconazole capsules.
- are allergic to itraconazole or any of the ingredients in itraconazole capsules. See the end of this Patient Information leaflet for a complete list of ingredients in itraconazole capsules.

**Before taking itraconazole capsules, tell your healthcare provider about all of your medical conditions, including if you:**

- have heart problems.
- have liver problems.
- have kidney problems.
- have a weakened immune system (immunocompromised).
- have lung problems including cystic fibrosis.
- are breastfeeding or plan to breastfeed. Itraconazole can pass into your breast milk. You and your healthcare provider should decide if you will take itraconazole capsules or breastfeed.

Taking itraconazole capsules with certain medicines may affect each other. Taking itraconazole capsules with other medicines can cause serious side effects.

**How should I take itraconazole capsules?**

- Take itraconazole capsules exactly as prescribed by your healthcare provider. Your healthcare provider will tell you how much itraconazole capsules to take and when to take it.
- You will receive itraconazole capsules in a bottle. Your healthcare provider will decide the type of itraconazole capsules that is right for you.
- Take itraconazole capsules with a full meal.
- Swallow itraconazole capsules whole.
- You should not take itraconazole capsules oral solution instead of itraconazole capsules, because they will not work the same way.
- If you take too much itraconazole capsules, call your healthcare provider or go to the nearest hospital emergency room right away.

**What should I avoid while taking itraconazole capsules?**

Itraconazole capsules can cause dizziness and vision problems. Do not drive or operate machinery until you know how itraconazole capsules affect you.

**What are the possible side effects of itraconazole capsules?**

**Itraconazole capsules may cause serious side effects, including:**

- See “**What is the most important information I should know about itraconazole capsules?**”
- **Nerve problems (neuropathy).** Call your healthcare provider right away if you have tingling or numbness in your hands or feet. Your healthcare provider may stop your treatment with itraconazole capsules if you have nerve problems.
- **Hearing loss.** Hearing loss can happen for a short time or permanently in some people who take itraconazole capsules. Stop taking itraconazole capsules and call your healthcare provider right away if you have any changes in your hearing.

**The most common side effects of itraconazole capsules include:** headache, rash, digestive system problems (such as nausea and vomiting), and edema. Additional possible side effects include upset stomach, constipation, fever, inflammation of the pancreas, menstrual disorder, erectile dysfunction, dizziness, muscle pain, painful joints, unpleasant taste, or hair loss. These are not all the possible side effects of itraconazole capsules. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store itraconazole capsules?**

- Store itraconazole capsules at room temperature between 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F).
- Keep itraconazole capsules dry and away from light.

**Keep itraconazole capsules and all medicines out of the reach of children.**

#### **General information about the safe and effective use of itraconazole capsules**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use itraconazole capsules for a condition for which it was not prescribed. Do not give itraconazole capsules to other people, even if they have the same symptoms that you have. It may harm them. You can ask your doctor or pharmacist for information about itraconazole capsules that is written for health professionals.

#### **What are the ingredients in itraconazole capsules?**

**Active ingredients:** itraconazole, USP

**Inactive ingredients:** black iron oxide, FD&C Blue No. 1, gelatin, hypromellose, liquid glucose, microcrystalline cellulose, polyethylene glycol, potassium hydroxide, propylene glycol, shellac, sodium lauryl sulfate, starch (of maize), strong ammonia solution, sucrose and titanium dioxide.

This Patient Information has been approved by the U.S. Food and Drug Administration.

\*All trademarks are property of their respective owner.

Manufactured by:

**Amneal Pharmaceuticals Pvt. Ltd.**

Ahmedabad 382220, INDIA

Distributed by:

**Amneal Pharmaceuticals LLC**

Bridgewater, NJ 08807

Rev. 05-2024-06

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 65162-630-03

Itraconazole Capsules, 100 mg

Rx only

30 Capsules

Amneal Pharmaceuticals LLC

NDC 65162-630-03

# Itraconazole Capsules, USP

## 100 mg



Rx only  
30 Capsules



**Each capsule contains:** itraconazole, USP 100 mg  
**Dosage:** See accompanying product literature.  
This is a bulk container. Not intended for household use. Dispense in tight container, as defined in the USP.  
Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture.  
Keep out of reach of children.  
Mfg. Lic. No. G/25/1941  
Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
Ahmedabad 382220, INDIA  
Distributed by: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807  
Rev. 08-2019-02



Non-Vanish Area  
(Size : 24 x 38 mm)  
For Lot No. and Exp. Date

## ITRACONAZOLE

itraconazole capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:65162-630
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ITRACONAZOLE (UNII: 304NUG5GF4) (ITRACONAZOLE - UNII:304NUG5GF4)	ITRACONAZOLE	100 mg

### Inactive Ingredients

Ingredient Name	Strength
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>AMMONIA</b> (UNII: 5138Q19F1X)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	

### Product Characteristics

<b>Color</b>	blue (transparent) , blue (opaque)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	22mm
<b>Flavor</b>		<b>Imprint Code</b>	AMNEAL;630
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65162-630-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	09/27/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205080	09/27/2016	

**Labeler** - Amneal Pharmaceuticals LLC (123797875)

### Establishment

Name	Address	ID/FEI	Business Operations
Amneal Pharmaceuticals Private Limited		915076126	analysis(65162-630) , label(65162-630) , manufacture(65162-630) , pack(65162-630)

Revised: 6/2024

Amneal Pharmaceuticals LLC