

AMPROMED-P FOR POULTRY- amprolium solution
Bimeda Inc., Division of Cross Vetpharm Group

AmproMed™- P FOR POULTRY

(amprolium)

9.6% Oral Solution

Coccidiostat

Water soluble treatment for coccidiosis

Treats 800 Gallons at 0.012% level.

Restricted Drug (California) – Use Only as Directed

ANADA 200-496, Approved by FDA

Net Contents:

1Gallon (128 fl oz) (3.785 L)

ACTIVE INGREDIENT: amprolium.....9.6%

FOR ANIMAL USE ONLY

To prepare 50 Gallon of Medicated Water

DOSAGE LEVEL MIXING DIRECTIONS

0.024 %	Add 1 pint (16 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012 %	Follow same directions as above but use 1/2 pint (8 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution.
0.006 %	Follow same directions as above but use 4 fluid ounces of AmproMed™ P FOR POULTRY 9.6% Oral Solution.

PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:

For use in automatic water proportioners that meter 1 fluid ounce of stock solution for per gallon of drinking water.

TO PREPARE DOSAGE LEVELS OF	AmproMed™ P FOR POULTRY 9.6 % Oral Solution	PER GALLON OF STOCK SOLUTION
0.024 %		41 fl oz
0.012 %		20.5 fl oz
0.006 %		10.25 fl oz

NOTE: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

Benzoic acid as Sodium Benzoate 0.1% added as preservative.

STORAGE: Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F) with brief excursions to 40°C.

To obtain an MSDS or for assistance contact Bimeda, Inc., at 1-888-524-6332.

INDICATIONS: AmproMed™ P FOR POULTRY 9.6 % Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

USE DIRECTIONS: Give amprolium at the 0.012% level (8fl oz AmproMed™ P FOR POULTRY 9.6% Oral Solution per 50 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024 % level). Continue with 0.006 % amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

WARNING: Keep this and all drugs out of reach of children.

NOT FOR HUMAN USE

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

FOR ANIMAL USE ONLY
To prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use ½ pint (8 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of AmproMed™ P FOR POULTRY 9.6% Oral Solution.

PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:
For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.


TO PREPARE DOSAGE LEVELS OF	AmproMed™ P FOR POULTRY 9.6% Oral Solution PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz


Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

Benzoic acid as Sodium Benzoate 0.1% added as preservative.

STORAGE: Store container in an upright position. Store between 5°C-25°C (41°F-77°F) with brief excursions to 40° C.

To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

AmproMed is a Trademark of Bimeda, Inc. 



INDICATIONS: AmproMed™ P FOR POULTRY 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

USE DIRECTIONS: Give amprolium at the 0.012% level (8 fl oz AmproMed™ P FOR POULTRY 9.6% Oral Solution per 50 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

WARNING: Keep this and all drugs out of reach of children. NOT FOR HUMAN USE.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

Product No. 1AMP024 8AMP016 Rev. 06/12



Lot:
Exp:

AMPROMED-P FOR POULTRY

amprolium solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:61133-8290
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM ION	96 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-8290-1	3785 mL in 1 JUG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200496	07/30/2012	

Labeler - Bimeda Inc., Division of Cross Vetpharm Group (060492923)

Registrant - Bimeda Inc., Division of Cross Vetpharm Group (060492623)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda Inc., division of Cross Vetpharm Group		060492923	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Zhejiang Jiuzhou		654682061	api manufacture

Revised: 8/2013

Bimeda Inc., Division of Cross Vetpharm Group