LEADER LICE TREATMENT- permethrin lotion Cardinal Health 110, LLC. dba Leader

Cardinal Health Lice Treatment Creme Rinse Drug Facts

Active ingredient (in each fluid ounce)

Permethrin 280 mg (1%)

Purpose

Lice treatment

Use

treats head lice

Warnings

For external use only

Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes.

See your doctor.

Ask a doctor before use if you are

• allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and see a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- · use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with the creme rinse

Treat

- wash hair with a shampoo without conditioner. Do not use a shampoo that
 contains a conditioner or a conditioner alone since this may decrease the activity of
 the creme rinse. Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle well
- completely saturate the hair and scalp with the creme rinse. Begin to apply the creme rinse behind the ears and at the back of the neck.
- keep the creme rinse out of the eyes. Protect the eyes with a washcloth or towel.
- leave the creme rinse on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

Remove Lice/Nits

- remove nits by combing the hair with the special small tooth comb provided.
 Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close
 to the scalp as possible and comb with a firm, even motion away from the scalp to
 the end of the hair.

- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

Other information

- read all the directions and warnings in the Consumer Information Insert before use. Keep the carton. It contains important information.
- store at 20 to 25°C (68 to 77°F)

Inactive ingredients

abies balsamea (balsam canada) extract, anhydrous citric acid, butylparaben, ceteth-10, cetyl alcohol, ethylparaben, FD&C yellow no. 6, fragrance, hydrolyzed animal protein, hydroxyethyl cellulose, isopropyl alcohol, methylparaben, phenoxyethanol, propylene glycol, propylparaben, purified water, stearalkonium chloride

Questions or comments?

1-800-719-9260

Consumer information

Permethrin Lotion, 1%

Lice Treatment Creme Rinse

THE FACTS OF LICE

Please read all information before using Lice Treatment Creme Rinse.

If you or a member of your household has head lice, don't panic. Keep calm.

Anyone can get head lice - it has nothing to do with cleanliness or poor hygiene. Head lice are treatable and very rarely cause serious health problems.

What are head lice?

- head lice are small insects that live and feed on the human scalp
- they may get as big as sesame seeds and are brown
- lice do not fly or jump from one person to another. They can only crawl.

What are nits?

- nits are lice eggs laid on the hair near the scalp. One louse may lay over 100 nits.
- nits are oval shaped, tiny and greyish-white
- they are attached to hair with a waterproof glue-like substance that cannot be washed out or blown away

Because nits (eggs) hatch in about one week, all nits must be removed to stop lice.

How do head lice spread?

- head lice spread easily from close head-to-head or hand-to-head contact with persons who have lice
- they may also spread by sharing hats, helmets, scarves, headphones, brushes, combs, bedding and clothing

How do you know if you have lice?

- itching is the most common symptom
- scratching behind the ears or the back of the neck is a sign that you should check for head lice
- scabs or blood spots may be seen

How do you prevent the spread of lice?

- do not share or borrow personal items like combs, brushes, or headbands
- do not share or borrow hats, sweaters, coats, scarves, helmets, headphones, pillows or stuffed animals
- store hats in coat sleeves and hang coats separately, so that they do not touch other people's hats and coats
- wash under fingernails when you first see lice or nits (eggs)
- check your child's head daily when you hear of a lice outbreak in day care or school

How to use the creme rinse

I. Inspect

If you suspect head lice - check. Head lice may be hard to locate because they avoid light. Nits are easier to see.



- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair

- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with the creme rinse



II. Treat

- wash hair with a shampoo without conditioner. Do not use a shampoo that
 contains a conditioner or a conditioner alone since this may decrease the activity of
 the creme rinse. Rinse with water.
- · towel dry hair so it is damp but not wet
- shake the bottle well
- completely saturate the hair and scalp with the creme rinse. Begin to apply the creme rinse behind the ears and at the back of the neck.
- · keep the creme rinse out of the eyes. Protect the eyes with a washcloth or towel.
- leave the creme rinse on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given



III. Remove Lice/Nits

Treatment kills lice but does not remove the nits.

remove nits by combing the hair with the special small tooth comb provided.
 Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.

- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close
 to the scalp as possible and comb with a firm, even motion away from the scalp to
 the end of the hair.
- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed



IV. Clean home and personal items

After using the creme rinse, it is very important to clean personal items and your home to prevent the spread of lice. Nits may live away from the human head for about 7-12 days. Lice may live away from the human head for about 1 day.

- all personal head gear (hats, hair ribbons, etc.), scarves, coats, towels and bed linens should be washed in hot water (above 130°F), then dried in the dryer using the hottest cycle for at least 20 minutes
- personal combs, including nit combs, and brushes should be soaked in hot water (above 130°F) for at least 10 minutes
- personal articles such as clothing, bedspreads, blankets, pillows or stuffed animals that cannot be washed should be dry-cleaned or sealed tightly in a plastic bag for a period of at least 4 weeks. Items should be taken out of the plastic bag outdoors and shaken out very hard before using again.
- for anything you cannot wash, dry-clean or store in a plastic bag, spray with lice control spray.
- vacuum everywhere. Vacuum all carpets, mattresses, upholstered furniture and car seats that may have been used by infested household members.



Warnings

For external use only

Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. See your doctor.

Ask a doctor before use if you are

• allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and see a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Questions or comments? 1-800-719-9260

Distributed By

Perrigo ®

Allegan, MI 49010 :95500 00 J2

Principal Display Panel

Permethrin Lotion 1%

Lice Treatment Creme Rinse

With a Single Application

Kills Lice & Their Eggs

Prevents Reinfestation for 14 Days

COMPARE TO NIX $^{\circledR}$ active ingredient


SAFE

Approved for Use on Children over 2 Months of Age

100% Money Back Guarantee

2 FL OZ (59 mL)

Contains: 1 Bottle Crème Rinse and 1 Nit Removal Comb



LEADER LICE TREATMENT

permethrin lotion

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:70000-0041 Route of Administration TOPICAL

| Active Ingredient/Active Moiety | | | | |
|--|--------------------------|---------------|--|--|
| Ingredient Name | Basis of Strength | Strength | | |
| PERMETHRIN (UNII: 509F88P9SZ) (PERMETHRIN - UNII:509F88P9SZ) | PERMETHRIN | 1 g in 100 mL | | |

| Inactive Ingredients | | |
|--|----------|--|
| Ingredient Name | Strength | |
| BALSAM FIR LEAFY TWIG (UNII: H1TFV1454Z) | | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | | |
| BUTYLPARABEN (UNII: 3QPI1U3FV8) | | |

| 1 | |
|---|--|
| CETETH-10 (UNII: LF9X1PN3XJ) | |
| CETYL ALCOHOL (UNII: 936JST6JCN) | |
| ETHYLPARABEN (UNII: 14255EXE39) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| ISOPROPYL ALCOHOL (UNII: ND2M416302) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| PROPYLPARABEN (UNII: Z8IX2SC10H) | |
| WATER (UNII: 059QF0KO0R) | |
| STEARALKONIUM CHLORIDE (UNII: 00U026BB88) | |
| f | |

| F | Packaging | | | | |
|---|----------------------|--|-------------------------|-----------------------|--|
| # | tem Code | Package Description | Marketing Start Date | Marketing End Date | |
| 1 | NDC:70000- 0041-1 | 1 in 1 CARTON | 03/19/2020 | | |
| 1 | | 59 mL in 1 BOTTLE; Type 0: Not a Combination Product | | | |

| Marketing Information | | | |
|-----------------------|---|-------------------------|-----------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA076090 | 03/19/2020 | |
| | | | |

Labeler - Cardinal Health 110, LLC. dba Leader (063997360)

Revised: 6/2021 Cardinal Health 110, LLC. dba Leader