

**MEIJER STOMACH RELIEF REGULAR STRENGTH- bismuth
subsalicylate suspension
MEIJER DISTRIBUTION INC**

**Meijer Stomach Relief Regular Strength Bismuth Subsalicylate 525 mg Drug
Facts**

Active ingredient (in each 30 mL dose cup)

Bismuth subsalicylate 525 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- only use for dose cup provided
- **adults and children 12 years and over:** (30 mL) 1 dose every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** sodium 10 mg
- **salicylate 261 mg**
- low sodium
- sugar free

- store at room temperature
- protect from freezing
- avoid excessive heat (over 104°F 40°C)

Inactive ingredients

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Do not use if imprinted shrink band is missing or broken.

Principal Display Panel

meijer®

NDC# **41250-670-12**

Compare To Pepto-Bismol® Regular Strength active ingredient

Stomach Relief

Bismuth Subsalicylate 525 mg

Regular Strength

Original Flavor

Naturally and Artificially Flavored

Provides soothing relief for:

Nausea

Heartburn

Indigestion

Upset Stomach

Diarrhea

12 FL. OZ. (354 mL)

OUR QUALITY GURANTEE

WWW.MEIJER.COM/SATISFACTION

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

***This product is not manufactured or distributed by The Procter & Gamble**

Co., owner of the registered trademark Pepto-Bismol®.

Drug Facts (continued)

Directions ■ shake well before use ■ only use dose cup provided
 ■ adults and children 12 years and over:
 ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)

Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments? 1-866-467-2748



Compare to Pepto-Bismol®
Regular Strength
active ingredient*

stomach relief

Bismuth Subsalicylate 525 mg

Regular Strength

Original Flavor
Naturally and Artificially Flavored

Provides soothing relief for:
Nausea, Heartburn, Indigestion
Upset Stomach, Diarrhea

12 FL. OZ. (354 mL)

Do not use if imprinted shrink band is missing or broken

Drug Facts

Active ingredient **Purposes**
(in each 30 mL dose)
Bismuth subsalicylate 525 mgUpset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:
 ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. ▶

*This product is not manufactured or distributed by The Procter & Gamble Co., owner of the registered trademark Pepto-Bismol®.

OUR QUALITY GUARANTEE

70612MJBLR PID 315000

LOT: UNVARNISHED AREA

EXP: 7 08820 06597 0

Do not print

MEIJER STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-670
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM HYDROXIDE (UNII: WZ H3C48M4T)	
POTASSIUM SORBATE (UNII: 1VPU26JZ4)	
WATER (UNII: 059QF0KO0R)	

SALICYLIC ACID (UNII: O414PZ4LPZ)

DIMETHICONE (UNII: 92RU3N3Y1O)

SODIUM BENZOATE (UNII: OJ245FE5EU)

SUCRALOSE (UNII: 96K6UQ3ZD4)

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	PINK (viscous)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-670-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/07/2017	

Labeler - MEIJER DISTRIBUTION INC (006959555)

Revised: 10/2023

MEIJER DISTRIBUTION INC