RECTICARE- lidocaine and glycerin cloth Ferndale Laboratories, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

RectiCare Medicated Anorectal Wipes

Active Ingredients

Lidocaine 5% w/w

Glycerin 20% w/w

Purpose

Lidocaine 5% w/w.....Local anesthetic

Glycerin 20% w/w.....Protectant

Uses

- helps relieve the pain, itching and burning associated with anorectal disorders
- temporarily provides a coating for relief of anorectal discomforts
- temporarily protects irritated and inflamed perianal skin

Warnings

For external use only

Do not

- exceed the recommended daily dosage unless directed by a doctor
- put this product into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor if

- an allergic reaction develops
- condition worsens or does not improve witin 7 days
- redness, irriation, swelling, pain or other symptoms begin or increase
- bleeding occurs

Keep out of reach of children. If swallowed, seek medical attention or contact a Poison Control Center right away.

Directions

- when practical, cleanse the affected area with mild soap and warm water and rinse thoroughly
- gently apply wipe to the affected area by patting and then discard
- adults and children 12 years & older: apply up to 6 times daily or after each bowel movement
- children under 12 years: consult a doctor

Inactive Ingredients

benzyl alcohol, citric acid, purified water

Package Label



Drug Facts	
Active Ingredients Lidocaine 5% w/w	
Uses a helps relieve the pain, itch ancrecial disorders a temporarily provid discomforts a temporarily protects in	les a coating for relief of anorecta
Warnings For external use only	
Do not mexceed the recommended of a doctor me put this product into the remechanical device or applicator	
Stop use and ask a dector if man a dition worsens or does not improve w swelling, pain or other symptoms begin	thin 7 days a redness, irritation
Keep out of reach of children. If sw or contact a Poison Control Center rigi	
Directions when practical, cle soap and warm water and rinse thoro affected area by patting and then disca and older: apply up to 6 times daily - children under 12 years: consult a c	ughly u gently apply wipe to the rd u adults and children 12 years or after each bowel movement
inactive Ingredients benzyl	alcohol, citric acid, purified water

RECTICARE

lidocaine and glycerin cloth

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0496-0980
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LIDO CAINE (UNII: 98 PI200987) (LIDO CAINE - UNII:98 PI200987)	LIDOCAINE	50 mg in 1 g	
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	200 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
WATER (UNII: 059QF0KO0R)		

ı	Packaging			
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:0496-0980-65	12 in 1 CARTON	0 1/0 1/20 16	
ı	1 NDC:0496-0980-76	6 g in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part346	0 1/0 1/20 16	

Labeler - Ferndale Laboratories, Inc. (005320536)

Registrant - Ferndale Laboratories, Inc. (005320536)

Establishment			
Name	Address	ID/FEI	Business Operations
Ferndale Laboratories, Inc.		005320536	manufacture(0496-0980)

Revised: 11/2019 Ferndale Laboratories, Inc.