

**PAIN RELIEF IBUPROFEN- ibuprofen tablet, film coated**  
**Rite Aid Corporation**

-----  
**Ibuprofen**

***Active ingredient (in each orange tablet)***

Ibuprofen USP, 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - menstrual cramps
  - toothache
  - the common cold
  - backache
  - headache
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - leg swelling
  - trouble breathing
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

***Principal Display Panel***

Compare to the active ingredient in **Motrin® IB Tablets\*\***

NDC 11822-0392-4

PAIN RELIEF

**IBUPROFEN**

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER

FEVER REDUCER

**(NSAID)**

ACTUAL SIZE

**500**

TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Tablets.  
50844 ORG122139214

**DISTRIBUTED BY:**  
RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011  
[www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE:**  
If you're not satisfied, we'll happily refund your money.

Compare to the active ingredient in Motrin® IB Tablets\*\*

NDC 11822-0392-4

**PAIN RELIEF**

**IBUPROFEN**

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER  
FEVER REDUCER  
(NSAID)

ACTUAL SIZE

**500**  
TABLETS



no print / no varnish area  
lot no. & exp. date

<b>Drug Facts</b>	<p><b>Active ingredient (in each orange tablet)</b> Ibuprofen USP 200 mg (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b> ■ temporarily relieves minor aches and pains due to:                  ■ backache ■ menstrual cramps ■ toothache                  ■ headache ■ the common cold ■ muscular aches                  ■ minor pain of arthritis ■ temporarily reduces fever</p> <p><b>Warnings</b>                  Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:                  ■ facial swelling ■ shock ■ asthma (wheezing)                  ■ skin reddening ■ hives ■ blisters ■ rash                  If an allergic reaction occurs, stop use and seek medical help right away.                  Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you                  ■ take more or for a longer time than directed                  ■ take a blood thinning (anticoagulant) or steroid drug                  ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older                  ■ have had stomach ulcers or bleeding problems                  ■ have 3 or more alcoholic drinks every day while using this product  <b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b>                  ■ if you have had an allergic reaction to any other pain reliever/fever reducer                  ■ right before or after heart surgery</p> <p><b>** This product is not manufactured or distributed by Johnson &amp; Johnson Corporation, owner of the registered trademark Motrin® IB Tablets.</b></p>
-------------------	---

**DISTRIBUTED BY:**  
RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011  
[www.riteaid.com](http://www.riteaid.com)



0 11822 1 42096 9

PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

<b>Questions or comments?</b>	<p>Call 1-800-426-5391 8:30 AM-4:00 PM ET, Monday-Friday</p>
<b>Inactive ingredients</b>	<p>camellia wax, colloidal silicon dioxide, corn starch, FD&amp;C yellow #5 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide</p>
<b>Other information</b>	<p>■ store between 20°-25°C (68°-77°F)                  ■ avoid excessive heat 40°C (104°F)                  ■ use by expiration date on package</p>
<b>Directions</b>	<p>■ do not take more than directed                  ■ the smallest effective dose should be used                  ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist                  ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used                  ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor                  ■ children under 12 years: ask a doctor</p>
<b>Drug Facts (continued)</b>	<p><b>Ask a doctor before use if</b>                  ■ stomach bleeding warning applies to you                  ■ you have a history of stomach problems, such as heartburn                  ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic                  ■ you have problems or serious side effects from taking pain relievers or fever reducers</p> <p><b>Ask a doctor or pharmacist before use if you are</b>                  ■ under a doctor's care for any serious condition                  ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug</p> <p><b>When using this product</b>                  ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b>                  ■ you experience any of the following signs of stomach bleeding:                  ■ feel faint ■ have bloody or black stools ■ vomit blood                  ■ have stomach pain that does not get better                  ■ you have symptoms of heart problems or stroke:                  ■ chest pain ■ slurred speech                  ■ leg swelling ■ trouble breathing                  ■ weakness in one part or side of body                  ■ pain gets worse or lasts more than 10 days                  ■ fever gets worse or lasts more than 3 days                  ■ redness or swelling is present in the painful area                  ■ any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.                  Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>

# PAIN RELIEF IBUPROFEN

ibuprofen tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0392
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;392
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-0392-2	1 in 1 CARTON	03/30/2002	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11822-0392-4	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/30/2002	
3	NDC:11822-0392-5	1 in 1 CARTON	03/30/2002	03/05/2022
3		50 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/30/2002	

**Labeler** - Rite Aid Corporation (014578892)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0392)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0392)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-0392)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11822-0392)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-0392)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0392)

Revised: 10/2023

Rite Aid Corporation