

**CHILDRENS COUGH MUCUS RELIEF- dextromethorphan hbr,
guaifenesin liquid
Discount Drug Mart**

Children's Cough Mucus Relief 512

Drug Facts

**Active ingredients
(in each 5 mL teaspoonful)**

Dextromethorphan HBr, USP 5 mg
Guaifenesin, USP 100 mg

Purpose

Cough suppressant
Expectorant

Uses

- temporarily relieves: cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain the bronchial tubes

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions , or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. if you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache.

These could be signs of a serious condition.

Keep out of reach of children.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- do not take more than 6 doses in any 24-hour period

age	dose
Children 6 years to 12 years hours	1 to 2 teaspoonfuls (tsps) every 4
Children 4 years to under 6 years hours	1/2 to 1 teaspoonful (tsps) every 4
Children under 4 years	do not use

Other information

- each teaspoonful contains sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided
- **Keep carton for full direction for use**

Inactive ingredients

anhydrous citric acid, dextrose, D&C red #33, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

Children's Cough Mucus Relief product label

DISCOUNT

drug

mart

FOOD FAIR

DO NOT USE IF PRINTED SEAL, UNDER CAP IS TORN OR MISSING

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX® COUGH*

Children's Cough Mucus Relief

Cough Suppressant

Expectorant

Dextromethorphan HBr 5 mg

Guaifensin 100 mg

Relieves Chest Congestion

Thins & Loosens Mucus

Alcohol free

for ages 4 to 12

CHERRY FLAVOR

4 FL OZ (118 ml)

* this product is not manufactured or distributed by Reckitt Benckiser Inc., the distributor of Childrens Mucinex® Cough.

SATISFACTION GUARANTEED

IF DISSATISFIED, RETURN THE UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS, AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

BX-052

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients *Purpose*
(in each 5 mL teaspoonful)
Dextromethorphan HBr, USP 5 mg.....Cough suppressant
Guaifenesin, USP 100 mg.....Expectorant

Uses

- temporarily relieves: cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain the bronchial tubes

Warnings
Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children
 In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 do not take more than 6 doses in any 24-hour period

age	dose
Children 6 years to under 12 years	1 to 2 teaspoonfuls (tsps) every 4 hours
Children 4 years to under 6 years	1/2 to 1 teaspoonful (tsp) every 4 hours
Children under 4 years	do not use

Drug Facts (continued)

Other information

- each teaspoon contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided
- Keep carton for full direction for use

Inactive ingredients
 anhydrous citric acid, dextrose, D&C red #33, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Children's Mucinex® Cough.

SATISFACTION GUARANTEED
 IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



BX-052



COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX® COUGH*

Children's Cough Mucus Relief

Cough Suppressant Expectorant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg

Relieves Chest Congestion
Thins & Loosens Mucus

Alcohol free
For ages 4 to 12

CHERRY FLAVOR



4 FL OZ (118 mL)



COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX® COUGH

Children's Cough Mucus Relief

Cough Suppressant Expectorant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg
Relieves Chest Congestion
Thins & Loosens Mucus
Alcohol free
For ages 4 to 12
CHERRY FLAVOR



DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING
Active ingredients (in each 5 mL): Cough suppressant Dextromethorphan HBr 5 mg, Expectorant Guaifenesin 100 mg.

Uses: Temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold. Helps loosen phlegm (mucus) and thin bronchial secretions to drain the bronchial tubes.

Warnings: Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has: cough that occurs with too much phlegm (mucus); persistent or chronic cough such as occurs with asthma.

When using this product: do not use more than directed.

Stop use and ask a doctor if: cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: do not take more than 6 doses in any 24-hour period.

Age	Dose
Children 6 years to under 12 years	1 to 2 teaspoonfuls (tsps) every 4 hours
Children 4 years to under 6 years	1/2 to 1 teaspoonful (tsp) every 4 hours
Children under 4 years	Do not use

Other information: each teaspoon contains sodium 3 mg. Store between 20-25°C (68-77°F). Do not refrigerate. Dosing cup provided. Keep carton for full direction for use.

Inactive ingredients: anhydrous citric acid, dextrose, D&C red #33, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944. **LOT:** **EXP:**

res

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients Purpose (in each 5 mL teaspoonful)

Dextromethorphan HBr, USP 5 mg.....Cough suppressant
Guaifenesin, USP 100 mg.....Expectorant

Uses

- temporarily relieves: cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain the bronchial tubes

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period

age	dose
Children 6 years to under 12 years	1 to 2 teaspoonfuls (tsps) every 4 hours
Children 4 years to under 6 years	1/2 to 1 teaspoonful (tsp) every 4 hours
Children under 4 years	do not use

Drug Facts (continued)

Other information

- each teaspoon contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided
- Keep carton for full direction for use

Inactive ingredients

anhydrous citric acid, dextrose, D&C red #33, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Children's Mucinex® Cough.

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



BX-052



COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX® COUGH*

Children's Cough Mucus Relief

Cough Suppressant Expectorant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg

Relieves Chest Congestion
Thins & Loosens Mucus

Alcohol free
For ages 4 to 12

CHERRY FLAVOR



4 FL OZ (118 mL)



COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX® COUGH

Children's Cough Mucus Relief

Cough Suppressant Expectorant

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg

Relieves Chest Congestion
Thins & Loosens Mucus

Alcohol free
For ages 4 to 12



CHERRY FLAVOR

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Active ingredients (in each teaspoonful, 5 mL):
Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 100 mg.....Expectorant

Uses: temporarily relieves: cough due to minor throat and bronchial irritation as may occur with a cold

Warnings: Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has: cough that occurs with too much phlegm (mucus)

When using this product: do not use more than directed

Stop use and ask a doctor if: cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: do not take more than 6 doses in any 24-hour period

Age	Dose
Children 6 years to under 12 years	1 to 2 teaspoonfuls every 4 hours
Children 4 years to under 6 years	1/2 to 1 teaspoonful every 4 hours
Children under 4 years	Do not use

Other information: each teaspoon contains sodium 3 mg

Inactive ingredients: anhydrous citric acid, dextrose, D&C red #33, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

SATISFACTION GUARANTEED IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LOT: EXP:

CHILDRENS COUGH MUCUS RELIEF

dextromethorphan hbr, guaifenesin liquid

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:53943-512

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DEXTROSE (UNII: IY9XDZ35W2)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALTITOL (UNII: D65DG142WK)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-512-24	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/04/2016	

Labeler - Discount Drug Mart (047741335)

Registrant - AptaPharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(53943-512)

Revised: 12/2023

Discount Drug Mart