

**COLD AND FLU RELIEF NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid
ASSURED / DOLLAR TREE (Greenbrier International, Inc.)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
 - headache
 - minor aches and pains
 - sore throat
 - runny nose and sneezing
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 doses of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause

liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **When using other Day Time or Night Time products, carefully read each label to ensure correct dosing**

Other information

- each 30 mL contains: potassium 5 mg
- each 30 mL contains: Sodium 19 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6,, flavor,high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

COMPARE TO ACTIVE INGREDIENTS IN VICKS® NYQUIL® COLD & FLU*

MULTI-SYMPTOM

Nitetime

Cold & Flu Relief

Acetaminophen 650 mg - Pain Reliever / Fever Reducer

Dextromethorphan HBr 30 mg - Cough Suppressant

Doxylamine succinate 12.5 mg - Antihistamine

- Alcohol 10%

Relieves Fever, Headache, Sore Throat, Cough, Minor Aches & Pains, Runny Nose & Sneezing

AGES 12 +

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY

GREENBRIER INTERNATIONAL, INC,

500 VOLVO PARKWAY

CHESAPEAKE, VA 23320

Product Label

COMPARE TO ACTIVE INGREDIENTS IN VICKS® NYQUIL® COLD & FLU*

ASSURED NDC 33992-0344-4

MULTI-SYMPTOM

Nitetime

Cold & Flu Relief

Acetaminophen 650 mg
Pain Reliever / Fever Reducer
Dextromethorphan HBr 30 mg - Cough Suppressant
Doxylamine succinate 12.5 mg - Antihistamine

- Alcohol 10%

Relieves Fever, Headache, Sore Throat, Cough, Minor Aches & Pains, Runny Nose & Sneezing

AGES 12+ 4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses ■ temporarily relieves common cold/flu symptoms ■ headache ■ minor aches and pains ■ sore throat ■ runny nose and sneezing ■ fever ■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY
GREENBRIER INTERNATIONAL, INC.
500 VOLVO PARKWAY
CHESAPEAKE, VA 23320

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

PLD-8407B
LB004213

Drug Facts (continued)

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers

Drug Facts (continued under label)

PEEL HERE →



Drug Facts (continued)

■ taking the blood thinning drug warfarin

When using this product ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see **Overdose warning**) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.

■ mL = milliliter ■ keep dosing cup with product ■ adults and children 12 years and over: 30 mL every 6 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information ■ each 30 mL contains: potassium 5 mg ■ each 30 mL contains: sodium 19 mg ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

ASSURED^N NiteTime Cold & Flu Relief Original Liquid

COLD AND FLU RELIEF NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:33992-0344
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33992-0344-4	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/31/2017	10/31/2024

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/31/2017	10/31/2024

Labeler - ASSURED / DOLLAR TREE (Greenbrier International, Inc.) (610322518)