

**PHENOXYBENZAMINE HYDROCHLORIDE - phenoxybenzamine
hydrochloride capsule
Novitium Pharma LLC**

**Phenoxybenzamine Hydrochloride Capsules, USP 10 mg
adrenergic, *alpha*-receptor-blocking agent
Rx only**


DESCRIPTION

Each Phenoxybenzamine Hydrochloride Capsule, USP 10 mg is a size #3 capsule with red transparent cap and body, imprinted "365" on cap and "novitium 10 mg" on body with grey ink, filled with white to off white powder containing 10 mg of Phenoxybenzamine Hydrochloride USP and the following inactive ingredients: lactose and talc.

Each of the empty hard gelatin capsule contains gelatin, purified water, sodium lauryl sulfate, D&C Red 33, FD&C Yellow 6 and FD&C Red 3.

The imprinting ink contains the following: shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, strong ammonia solution, potassium hydroxide, titanium dioxide, black iron oxide and purified water.

Phenoxybenzamine Hydrochloride is N-(2-Chloroethyl)-N-(1-methyl-2-phenoxyethyl) benzylamine hydrochloride:



Phenoxybenzamine hydrochloride is a white to off-white powder with a molecular weight of 340.3, which melts between 136° and 141°C. It is soluble in water, alcohol and chloroform; insoluble in ether.

CLINICAL PHARMACOLOGY

Phenoxybenzamine hydrochloride is a long-acting, adrenergic, *alpha*-receptor-blocking agent, which can produce and maintain "chemical sympathectomy" by oral administration. It increases blood flow to the skin, mucosa and abdominal viscera, and

lowers both supine and erect blood pressures. It has no effect on the parasympathetic system.

Twenty to 30 percent of orally administered phenoxybenzamine appears to be absorbed in the active form.¹

The half-life of orally administered phenoxybenzamine hydrochloride is not known; however, the half-life of intravenously administered drug is approximately 24 hours. Demonstrable effects with intravenous administration persist for at least 3 to 4 days, and the effects of daily administration are cumulative for nearly a week.¹

INDICATIONS & USAGE

Phenoxybenzamine Hydrochloride Capsules are indicated in the treatment of pheochromocytoma, to control episodes of hypertension and sweating. If tachycardia is excessive, it may be necessary to use a *beta*-blocking agent concomitantly.

CONTRAINDICATIONS

Conditions where a fall in blood pressure may be undesirable; hypersensitivity to the drug or any of its components.

WARNINGS

Phenoxybenzamine hydrochloride-induced *alpha*-adrenergic blockade leaves *beta*-adrenergic receptors unopposed. Compounds that stimulate both types of receptors may, therefore, produce an exaggerated hypotensive response and tachycardia.

PRECAUTIONS

GENERAL PRECAUTIONS

Administer with caution in patients with marked cerebral or coronary arteriosclerosis or renal damage. Adrenergic blocking effect may aggravate symptoms of respiratory infections.

DRUG INTERACTIONS

Phenoxybenzamine hydrochloride may interact with compounds that stimulate both *alpha*- and *beta*-adrenergic receptors (i.e., epinephrine) to produce an exaggerated hypotensive response and tachycardia. (See WARNINGS.)

Phenoxybenzamine hydrochloride blocks hyperthermia production by levarterenol, and blocks hypothermia production by reserpine.

CARCINOGENESIS & MUTAGENESIS & IMPAIRMENT OF FERTILITY

Case reports of carcinoma in humans after long-term treatment with

phenoxybenzamine have been reported. Hence long-term use of phenoxybenzamine is not recommended.^{3,4} Carefully weigh the benefits and risks before prescribing this drug.

Phenoxybenzamine hydrochloride showed *in vitro* mutagenic activity in the Ames test and mouse lymphoma assay; it did not show mutagenic activity *in vivo* in the micronucleus test in mice. In rats and mice, repeated intraperitoneal administration of phenoxybenzamine hydrochloride (three times per week for up to 52 weeks) resulted in peritoneal sarcomas. Chronic oral dosing in rats (for up to 2 years) produced malignant tumors of the small intestine and non-glandular stomach, as well as ulcerative and/or erosive gastritis of the glandular stomach. Whereas squamous cell carcinomas of the non-glandular stomach were observed at all tested doses of phenoxybenzamine hydrochloride, there was a no-observed-effect-level of 10 mg/kg for tumors (carcinomas and sarcomas) of the small intestine. This dose is, on a body surface area basis, about twice the maximum recommended human dosage of 20 mg b.i.d.

PREGNANCY

Teratogenic Effects

Adequate reproductive studies in animals have not been performed with phenoxybenzamine hydrochloride. It is also not known whether phenoxybenzamine hydrochloride can cause fetal harm when administered to a pregnant woman. Phenoxybenzamine hydrochloride should be given to a pregnant woman only if clearly needed.

NURSING MOTHERS

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, and because of the potential for serious adverse reactions from phenoxybenzamine hydrochloride, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

PEDIATRIC USE

Safety and effectiveness in pediatric patients have not been established.

ADVERSE REACTIONS

The following adverse reactions have been observed, but there are insufficient data to support an estimate of their frequency.

Autonomic Nervous System*: Postural hypotension, tachycardia, inhibition of ejaculation, nasal congestion, miosis.

*These so-called "side effects" are actually evidence of adrenergic blockade and vary according to the degree of blockade.

Miscellaneous: Gastrointestinal irritation, drowsiness, fatigue.

To report SUSPECTED ADVERSE REACTIONS, contact Novitium Pharma, LLC at 1-855-204-1431 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

SYMPTOMS – These are largely the result of blocking of the sympathetic nervous system and of the circulating epinephrine. They may include postural hypotension, resulting in dizziness or fainting; tachycardia, particularly postural; vomiting; lethargy; shock.

TREATMENT

When symptoms and signs of overdosage exist, discontinue the drug. Treatment of circulatory failure, if present, is a prime consideration. In cases of mild overdosage, recumbent position with legs elevated usually restores cerebral circulation. In the more severe cases, the usual measures to combat shock should be instituted. Usual pressor agents are not effective. Epinephrine is contraindicated because it stimulates both *alpha*- and *beta*-receptors; since *alpha*-receptors are blocked, the net effect of epinephrine administration is vasodilation and a further drop in blood pressure (epinephrine reversal).

The patient may have to be kept flat for 24 hours or more in the case of overdose, as the effect of the drug is prolonged. Leg bandages and an abdominal binder may shorten the period of disability.

I.V. Infusion of levarterenol bitartrate** may be used to combat severe hypotensive reactions, because it stimulates *alpha*-receptors primarily. Although phenoxybenzamine hydrochloride is an *alpha*-adrenergic blocking agent, a sufficient dose of levarterenol bitartrate will overcome this effect.

The oral LD₅₀ for phenoxybenzamine hydrochloride is approximately 2000 mg/kg in rats and approximately 500 mg/kg in guinea pigs.

DOSAGE & ADMINISTRATION

The dosage should be adjusted to fit the needs of each patient. Small initial doses should be slowly increased until the desired effect is obtained or the side effects from blockade become troublesome. After each increase, the patient should be observed on that level before instituting another increase. The dosage should be carried to a point where symptomatic relief and/or objective improvement are obtained, but not so high that the side effects from blockade become troublesome.

Initially, 10 mg of phenoxybenzamine hydrochloride twice a day. Dosage should be increased every other day, usually to 20 to 40 mg 2 or 3 times a day, until an optimal dosage is obtained, as judged by blood pressure control.

Long-term use of phenoxybenzamine is not recommended (see PRECAUTIONS Carcinogenesis and Mutagenesis).

STORAGE

Store at 25°C (77°F); excursions permitted to 15°- 30°C (59°- 86°F) [See USP Controlled Room Temperature]. Dispense in a tight container.

HOW SUPPLIED

Phenoxybenzamine Hydrochloride Capsules, USP 10 mg are supplied as size #3 capsule with red transparent cap and body, imprinted "365" on cap and "novitium 10 mg" on body with grey ink, filled with white to off white powder in bottles of 100 (NDC 70954-365-10).

REFERENCES

1. Weiner, N.: Drugs That Inhibit Adrenergic Nerves and Block Adrenergic Receptors, in Goodman, L., and Gilman, A., *The Pharmacological Basis of Therapeutics*, ed. 6, New York, Macmillan Publishing Co., 1980, p. 179; p. 182.
2. Martin, E.W.: *Drug Interactions Index 1978/1979*, Philadelphia, J.B. Lippincott Co., 1978, pp. 209- 210.
3. Nettesheim O, Hoffken G, Gahr M, Breidert M: Haematemesis and dysphagia in a 20-year-old woman with congenital spine malformation and situs inversus partialis [German]. *Zeitschrift fur Gastroenterologie*. 2003; 41(4):319-24.
4. Vaidyanathan S, Mansour P, Soni BM, Hughes PL, Singh G: Chronic lymphocytic leukaemia, synchronous small cell carcinoma and squamous neoplasia of the urinary bladder in a paraplegic man following long-term phenoxybenzamine therapy. *Spinal Cord*. 2006;44(3):188-91.

**Available as Levophed® (brand of norepinephrine bitartrate) from Hospira Inc.

All trademarks are the property of their respective owners.

Manufactured by:

Novitium Pharma LLC

70 Lake Drive, East Windsor

New Jersey 08520

Issued: 06/2022

LB4512-00

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Phenoxybenzamine Hydrochloride Capsules, USP 10 mg
100 capsules per bottle

NDC 70954- 365 -10 100 Capsules		Each capsule contains phenoxybenzamine hydrochloride, USP 10 mg. Dosage: See accompanying prescribing information. Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Dispense in a tight container. Keep this and all medications out of the reach of children.	
Phenoxybenzamine Hydrochloride Capsules, USP 10 mg 		Manufactured by: Novitium Pharma LLC 70 Lake Drive, East Windsor New Jersey 08520 Rx only Issued: 06/2022 LB4513-00	

PHENOXYBENZAMINE HYDROCHLORIDE

phenoxybenzamine hydrochloride capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70954-365
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOXYBENZAMINE HYDROCHLORIDE (UNII: X1IEG24OHL) (PHENOXYBENZAMINE - UNII:0TTZ664R7Z)	PHENOXYBENZAMINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE (UNII: J2B2A4N98G)	
TALC (UNII: 7SEV7J4R1U)	
GELATIN (UNII: 2G86QN327L)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
SHELLAC (UNII: MB5IUD6JUA)	
ALCOHOL (UNII: 3K9958V90M)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
AMMONIA (UNII: 5138Q19F1X)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	RED	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	365;novitium10mg
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70954-365-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/19/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215042	07/19/2022	

Labeler - Novitium Pharma LLC (080301870)

Establishment

Name	Address	ID/FEI	Business Operations
Novitium Pharma LLC		080301870	ANALYSIS(70954-365) , LABEL(70954-365) , MANUFACTURE(70954-365) , PACK(70954-365)

Revised: 10/2022

Novitium Pharma LLC