

BENZO JEL- benzocaine spray, metered
Henry Schein Inc.

Active Ingredient

Benzocaine 20%

Purpose

Oral Anesthetic

Uses

- For the temporary relief of occasional minor irritation and pain associated with canker sores and minor dental procedures.
- For Professional Use Only.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Flammable. Do not use near fire or flame. Avoid smoking during use.

Do not use:

- for teething
- in children under 2 years of age

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Avoid spraying in eyes.

Dosage

A single metered spray dispenses 50 mg of Benzo-Jel Topical Anesthetic. Use only as

directed by a dentist.

Other information

Store at room temperature. Do not store above 120°F (49°C). Dispose of in a manner consistent with local regulations.

Inactive ingredients

Ethyl Alcohol, Flavor, Polyethelyn Glycol, Sodium Saccharin, Hydrocarbon Propellant (HFC-134a).

Questions or comments?

1-800-472-4346

HENRY SCHEIN® REF 570-0191

BENZO-JEL™ SPRAY

Metered Dose

- (EN) Topical Anesthetic
- (ES) Anestésico de uso tópico
- (FR) Analgésique local



Dosage: Single Spray 50 mg
Important: Read directions for proper use.

Net Wt.
2 oz. (57g)

Distributed by (in US only):
HENRY SCHEIN INC.
135 DURYEA ROAD
MELVILLE, NY 11747 USA

Drug Facts

Active ingredient	Purpose
Benzocaine 20%	Oral Anesthetic

Uses

- For the temporary relief of occasional minor irritation and pain associated with canker sores and minor dental procedures.
- For Professional Use Only.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Flammable. Do not use near fire or flame. Avoid smoking during use.

Do not use

- for teething
- in children under 2 years of age

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Avoid spraying in eyes.

Directions

Apply only amount needed to the oral mucosa to prevent or relieve pain.

- Children under 2 years of age: do not use

Other information

Store at room temperature. Do not store above 120°F (49°C). Dispose of in a manner consistent with local regulations.

Inactive ingredients

Ethyl Alcohol, Flavor, Polyethylene Glycol, Sodium Saccharin, Hydrocarbon Propellant (HFC-134a).

Questions or comments?

1-800-472-4346



9039550

Made in USA from
Global Materials
Rev 2022/01

HENRY SCHEIN® REF 570-0191

BENZO-JEL™ SPRAY

Metered Dose

- (EN) Topical Anesthetic
- (ES) Anestésico de uso tópico
- (FR) Analgésique local



Dosage: Single Spray 50 mg
Important: Read directions for proper use.

Net Wt.
2 oz. (57g)

Distributed by (in US only):
HENRY SCHEIN INC.
135 DURYEA ROAD
MELVILLE, NY 11747 USA




BENZO-JEL™ SPRAY

Metered Dose

- EN Topical Anesthetic
- ES Anestésico de uso tópico
- FR Analgésique local

Net Wt.
2 oz. (57g)



Drug Facts	
Active ingredient	Purpose
Benzocaine 20%	Oral Anesthetic
Uses	
<ul style="list-style-type: none"> • For the temporary relief of occasional minor irritation and pain associated with canker sores and minor dental procedures. • For Professional Use Only. 	
Warnings	
<p>Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy 	
<p>Allergy Alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.</p> <p>Flammable. Do not use near fire or flame. Avoid smoking during use.</p>	

Drug Facts (continued)
<p>Do not use</p> <ul style="list-style-type: none"> • for teething • in children under 2 years of age <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Avoid spraying in eyes.</p>
<p>Directions</p> <p>Apply only amount needed to the oral mucosa to prevent or relieve pain.</p> <ul style="list-style-type: none"> • Children under 2 years of age: do not use
<p>Other information</p> <p>Store at room temperature. Do not store above 120°F (49°C). Dispose of in a manner consistent with local regulations.</p>
<p>Inactive ingredients</p> <p>Ethyl Alcohol, Flavor, Polyethylene Glycol, Sodium Saccharin, Hydrocarbon Propellant (HFC-134a).</p>
<p>Questions or comments?</p> <p>1-800-472-4346</p>

<p>Dosage:</p> <p>Single Spray 50 mg</p> <p>Important: Read directions for proper use.</p>	<p><i>Distributed by (In US only):</i></p> <p>HENRY SCHEIN INC. 135 DURYEA ROAD MELVILLE, NY 11747 USA</p> <p>Made in USA from Global Materials Rev 2022/01 99088510</p>
--	--

BENZO JEL

benzocaine spray, metered

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0404-0018
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	11.4 g in 57 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	white (CLEAR)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0404-0018-35	57 g in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)	02/01/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M006	02/01/2016	

Labeler - Henry Schein Inc. (012430880)

Revised: 2/2024

Henry Schein Inc.