

**GOOD SENSE ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet  
L. Perrigo Company**

-----  
**Perrigo Anti-Diarrheal/Anti-Gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse

- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20° -25° C (68° -77° F). Protect from light.
- **do not use if printed foil under cap is broken or missing**

**Inactive ingredients**

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

Multi-Symptom Relief Caplet

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

Actual Size

Relieves symptoms of diarrhea plus Cramps & Pressure - Bloating - Gas

Compare to active ingredients of Imodium<sup>®</sup> Multi-Symptom Relief

100% SATISFACTION GUARANTEED

24 Caplets\*

\*Capsule-Shaped Tablets

GOODSENSE.

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

# Anti-Diarrheal/Anti-Gas

GOODSENSE.

NDC 0113-0087-62

Multi-Symptom Relief Caplet

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

# Anti-Diarrheal/Anti-Gas



Actual Size

Relieves symptoms of diarrhea **plus**  
• Cramps & Pressure • Bloating • Gas

24 Caplets\*

\*Capsule-Shaped Tablets

Compare to active ingredients of Imodium® Multi-Symptom Relief



OR162 C2 C1

### Drug Facts

**Active ingredients (in each caplet)** Purpose  
Loperamide HCl 2 mg.....Anti-diarrheal  
Simethicone 125 mg.....Anti-gas

**Uses** Relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

### Warnings

Allergic alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

### Do not use

- if you have blood or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have  
■ fever ■ mucus in the stool ■ a history of liver disease  
■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (1-800-222-1222).

### Directions

- drink plenty of clear fluids to help prevent dehydration on case of diarrhea
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz) glass of water
- find right dose on chart below. If possible, use weight to do so; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- do not use if printed blister unit is broken or torn

### Drug Facts (continued)

**Inactive ingredients** acetaminophen, polyethylene glycol, croscarmellose sodium, dibasic calcium phosphate, methylcellulose, microcrystalline cellulose, natural and artificial flavors, propylene glycol, stearic acid



Distributed By: Perrigo® Allegan, MI 49010

Questions or comments? 1-800-719-0260

Gluten Free



## GOOD SENSE ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-0087
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	L0R1;2;125
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0087-53	12 in 1 CARTON	10/11/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0113-0087-65	1 in 1 CARTON	10/16/2018	
2		30 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0113-0087-62	24 in 1 CARTON	06/10/2019	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	10/11/2018	

**Labeler** - L. Perrigo Company (006013346)

Revised: 6/2019

L. Perrigo Company