

**EQUALINE ANTI DIARRHEAL PLUS ANTI GAS- loperamide hydrochloride,  
simethicone tablet  
United Natural Foods, Inc. dba UNFI**

-----  
**SuperValu Inc. Anti-Diarrheal + Anti-Gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

|                                       |   |
|---------------------------------------|---|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 9-11 years (60-95 lbs)       | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours  |
| children 6-8 years (48-59 lbs)        | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours  |
| children 2-5 years (34 to 47 lbs)     | ask a doctor  |
| children under 2 years (up to 33 lbs) | do not use  |

## Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **Do not use if printed blister unit is broken or torn**

## Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

## Questions or comments?

**1-877-932-7948**

### **Package/Label Principal Display Panel**

compare to Imodium<sup>®</sup> Multi-Symptom Relief

active ingredients

anti-diarrheal + anti-gas

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg (anti-diarrheal + anti-gas)

multi-symptom relief caplet

relieves symptoms of diarrhea plus

cramps & pressure

bloating

gas

actual size

18 caplets\*

\*capsule-shaped tablets

# EQUALINE®

## anti-diarrheal + anti-gas

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg  
(anti-diarrheal + anti-gas)

# EQUALINE®

## anti-diarrheal + anti-gas

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg  
(anti-diarrheal + anti-gas)

multi-symptom relief caplet

relieves symptoms of diarrhea *plus*

- cramps & pressure
- bloating
- gas

compare to  
Imodium®  
Multi-Symptom Relief  
active ingredients\*\*  
NDC 41163-553-89



actual size

**18 caplets\*** \*capsule-shaped tablets

OR189 EL C1

### Drug Facts

**Active Ingredients (in each caplet)** **Purposes**  
 Loperamide HCl 2 mg..... Anti-diarrheal  
 Simethicone 125 mg..... Anti-gas

**Uses** relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

### Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
 Heart alert: Taking more than directed can cause serious heart problems or death

### Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

### Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

### Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

### Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

|                                       |   |
|---------------------------------------|---|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 9-11 years (60-65 lbs)       | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours  |
| children 6-8 years (48-50 lbs)        | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours  |
| children 2-5 years (34 to 47 lbs)     | ask a doctor  |
| children under 2 years (up to 33 lbs) | do not use  |

### Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- do not use if printed blister unit is broken or torn

\*\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Imodium®.

Distributed by  
SUPERVALU INC.  
Eden Prairie,  
MN 55344 USA

**100% Quality  
GUARANTEED**

Like it or let us make it right.  
That's our quality promise.  
877-933-7948  
supervalu private brands.com

### Drug Facts (continued)

**Inactive ingredients** acetaminophen, potassium



croscarmellose sodium, dibasic calcium phosphate,  
maltodextrin, microcrystalline cellulose, natural and artificial  
flavor, propylene glycol, stearic acid

Questions or comments? 1-877-932-7948

GLUTEN FREE



## EQUALINE ANTI DIARRHEAL PLUS ANTI GAS

loperamide hydrochloride, simethicone tablet

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:41163-553 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength        | Strength |
|---|--------------------------|----------|
| <b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II) | LOPERAMIDE HYDROCHLORIDE | 2 mg     |
| <b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)             | DIMETHICONE              | 125 mg   |

### Inactive Ingredients

| Ingredient Name   | Strength |
|---|----------|
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                     |          |
| <b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)                |          |
| <b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)               |          |
| <b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J) |          |
| <b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)                        |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)          |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                    |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                        |          |

### Product Characteristics

|                 |       |                     |            |
|-----------------|-------|---------------------|------------|
| <b>Color</b>    | WHITE | <b>Score</b>        | 2 pieces   |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 17mm       |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | L0R1;2;125 |
| <b>Contains</b> |       |                     |            |

### Packaging

| # | Item Code        | Package Description                                    | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:41163-553-89 | 18 in 1 CARTON   | 12/04/2019           |                    |
| 1 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA209837                               | 12/04/2019           |                    |

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 8/2023

United Natural Foods, Inc. dba UNFI