


ECONOPRED- prednisolone acetate suspension Alcon Laboratories, Inc.

1/8% Econopred®
1% Econopred® Plus
(prednisolone acetate)
Ophthalmic Suspension

DESCRIPTION

ECONOPRED® and ECONOPRED® Plus (Prednisolone Acetate) are adrenocortical steroid products prepared as sterile ophthalmic suspensions. The active ingredient is represented by the chemical structure:



Established name:
Prednisolone Acetate

Chemical name:
Pregna-1,4-diene-3,20-dione,
21-(acetyloxy)-11,17-
dihydroxy-, (11 β)-.

Each mL contains: Active: Prednisolone Acetate 1.0% or 0.125%. **Preservative:** Benzalkonium Chloride 0.01%. **Vehicle:** Hydroxypropyl Methylcellulose. **Inactive:** Dibasic Sodium Phosphate, Polysorbate 80, Edetate Disodium, Glycerin, Citric Acid and/or Sodium Hydroxide (to adjust pH), Purified Water. DM-02, DM-03

CLINICAL PHARMACOLOGY

Corticosteroids inhibit the inflammatory response to a variety of inciting agents and probably delay or slow healing. They inhibit the edema, fibrin deposition, capillary dilation, leukocyte migration, capillary proliferation, fibroblast proliferation, deposition of collagen, and scar formation associated with inflammation.

There is no generally accepted explanation for the mechanism of action of ocular corticosteroids. However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

Corticosteroids are capable of producing a rise in intraocular pressure.

INDICATIONS AND USAGE

Steroid responsive inflammatory conditions of the palpebral and bulbar conjunctiva, cornea, and anterior segment of the globe such as allergic conjunctivitis, acne rosacea, superficial punctate keratitis, herpes zoster keratitis, iritis, cyclitis, selected infective conjunctivides, when the inherent hazard of steroid use is accepted to obtain an advisable diminution in edema and inflammation; corneal injury from chemical, radiation, or thermal burns, or penetration of foreign bodies.

CONTRAINDICATIONS

ECONOPRED and ECONOPRED Plus are contraindicated in most viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures. ECONOPRED and ECONOPRED Plus are also contraindicated in individuals with known or suspected hypersensitivity to any of the ingredients of this preparation and to other corticosteroids.

WARNINGS

Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and in posterior subcapsular cataract formation. Prolonged use may also suppress the host immune response and thus increase the hazard of secondary ocular infections.

Various ocular diseases and long-term use of topical corticosteroids have been known to cause corneal and scleral thinning. Use of topical corticosteroids in the presence of thin corneal or scleral tissue may lead to perforation.

Acute purulent infections of the eye may be masked or activity enhanced by the presence of corticosteroid medication.

If this product is used for 10 days or longer, intraocular pressure should be routinely monitored even though it may be difficult in children and uncooperative patients. Steroids should be used with caution in the presence of glaucoma. Intraocular pressure should be checked frequently.

The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation.

Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex). Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution; frequent slit lamp microscopy is recommended.

Corticosteroids are not effective in mustard gas keratitis and Sjögren's keratoconjunctivitis.

PRECAUTIONS

General

The initial prescription and renewal of the medication order should be made by a physician only after examination of the patient with the aid of magnification, such as slit lamp biomicroscopy and, where appropriate, fluorescein staining. If signs and symptoms fail to improve after two days, the patient should be re-evaluated.

As fungal infections of the cornea are particularly prone to develop coincidentally with long-term local corticosteroid applications, fungal invasion should be suspected in any persistent corneal ulceration where a corticosteroid has been used or is in use. Fungal cultures should be taken when appropriate.

If this product is used for 10 days or longer, intraocular pressure should be monitored (SEE WARNINGS).

Information for Patients

If inflammation or pain persists longer than 48 hours or becomes aggravated, the patient should be advised to discontinue use of the medication and consult a physician.

This product is sterile when packaged. To prevent contamination, care should be taken to avoid touching the bottle tip to eyelids or to any other surface. The use of this bottle by more than one person

may spread infection. Keep bottle tightly closed when not in use. Keep out of the reach of children.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No studies have been conducted in animals or in humans to evaluate the potential of these effects.

Pregnancy

Teratogenic effects.

Pregnancy Category C. Prednisolone has been shown to be teratogenic in mice when given in doses 1-10 times the human dose. Dexamethasone, hydrocortisone and prednisolone were ocularly applied to both eyes of pregnant mice five times per day on days 10 through 13 of gestation. A significant increase in the incidence of cleft palate was observed in the fetuses of the treated mice. There are no adequate and well controlled studies in pregnant women. ECONOPRED® and ECONOPRED® Plus should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. Because of the potential for serious adverse reactions in nursing infants from prednisolone acetate, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

ADVERSE REACTIONS

Adverse reactions include, in decreasing order of frequency, elevation of intraocular pressure (IOP) with possible development of glaucoma and infrequent optic nerve damage, posterior subcapsular cataract formation, and delayed wound healing.

Although systemic effects are extremely uncommon, there have been rare occurrences of systemic hypercorticism after use of topical steroids.

Corticosteroid-containing preparations have also been reported to cause acute anterior uveitis and perforation of the globe. Keratitis, conjunctivitis, corneal ulcers, mydriasis, conjunctival hyperemia, loss of accommodation and ptosis have occasionally been reported following local use of corticosteroids.

The development of secondary ocular infection (bacterial, fungal and viral) has occurred. Fungal and viral infections of the cornea are particularly prone to develop coincidentally with long-term applications of steroid. The possibility of fungal invasion should be considered in any persistent corneal ulceration where steroid treatment has been used (SEE WARNINGS).

DOSAGE AND ADMINISTRATION

SHAKE WELL BEFORE USING. Two drops topically in the eye(s) four times daily. In cases of bacterial infections, concomitant use of anti-infective agents is mandatory. Care should be taken not to discontinue therapy prematurely.

If signs and symptoms fail to improve after two days, the patient should be re-evaluated (SEE PRECAUTIONS).

The dosing of ECONOPRED and ECONOPRED Plus may be reduced, but care should be taken not to

discontinue therapy prematurely. In chronic conditions, withdrawal of treatment should be carried out by gradually decreasing the frequency of applications.

HOW SUPPLIED

5mL and 10mL in plastic DROP-TAINER® dispensers.

1/8% ECONOPRED: 1% ECONOPRED Plus:

5 mL NDC 0998-0635-05 5 mL NDC 0998-0637-05

10 mL NDC 0998-0635-10 10 mL NDC 0998-0637-10

STORAGE: STORE at 8° - 24°C (46° - 75°F) in an UPRIGHT position.

Rx Only

Alcon Laboratories, Inc.

Fort Worth, Texas 76134 USA

Printed in USA

236055-0598

ECONOPRED				
prednisolone acetate suspension				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0998-0635	
Route of Administration	OPHTHALMIC			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
prednisolone acetate (UNII: 8B2807733D) (prednisolone - UNII:9PHQ9Y1OLM)			1.25 mg in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
benzalkonium chloride ()			0.1 mg in 1 mL	
hydroxypropyl methylcellulose ()				
dibasic sodium phosphate (UNII: GR686LBA74)				
polysorbate 80 ()				
edetate disodium (UNII: 7FLD91C86K)				
glycerin (UNII: PDC6A3C0OX)				
citric acid and/or sodium hydroxide ()				
water (UNII: 059QF0K00R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0998-0635-05	5 mL in 1 BOTTLE, PLASTIC		
2	NDC:0998-0635-10	10 mL in 1 BOTTLE, PLASTIC		

ECONOPRED

prednisolone acetate suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0998-0637
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
prednisolone acetate (UNII: 8B2807733D) (prednisolone - UNII:9PHQ9Y1OLM)		10 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
benzalkonium chloride ()	0.1 mg in 10 mL
hydroxypropyl methylcellulose ()	
dibasic sodium phosphate (UNII: GR686LBA74)	
polysorbate 80 ()	
edetate disodium (UNII: 7FLD91C86K)	
glycerin (UNII: PDC6A3C0OX)	
citric acid and/or sodium hydroxide ()	
water (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0998-0637-05	5 mL in 1 BOTTLE, PLASTIC		
2	NDC:0998-0637-10	10 mL in 1 BOTTLE, PLASTIC		

Labeler - Alcon Laboratories, Inc.

Revised: 9/2006

Alcon Laboratories, Inc.