

**DAYTIME COLD FLU MULTI SYMPTOM- acetaminophen, dextromethorphan hbr, phenylephrine hci capsule, liquid filled
P & L Development, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver

damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

butylated hydroxyanisole, butylated hydroxytoluene, FD&C red #40*, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide*, white ink

*may contain this ingredient

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

daytime multi-symptom

cold & flu relief

acetaminophen (pain reliever / fever reducer)

dextromethorphan HBr (cough suppressant)

phenylephrine HCl (nasal decongestant)

- non-drowsy
- alcohol-free
- antihistamine-free

softgels**

(**liquid-Filled capsules)

†Compare to the active ingredients in Vicks® Dayquil® Cold & Flu LiquiCap®

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by:

PL Developments

200 Hicks Street

Westbury, NY 11590

Product Label





daytime multi-symptom cold & flu relief

acetaminophen

(pain reliever/fever reducer)

dextromethorphan HBr

(cough suppressant)

phenylephrine HCl

(nasal decongestant)

- non-drowsy
- alcohol-free
- antihistamine-free

24 softgels**

(**liquid-filled capsules)



IF PER EVIDENCE: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
 KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

it No.:
 p. Date:

Drug Facts	
Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant	Purposes temporarily relieves common cold and flu symptoms minor aches and pains nasal congestion cough due to minor throat and bronchial irritation

Drug Facts (continued)	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not exceed recommended dosage.	Drug Facts (continued) Stop use and ask a doctor if ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleepiness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use Label to ensure correct dosing ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat Inactive ingredients butylated hydroxyanisole, butylated hydroxytoluene, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide, white ink Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



†Compare to the active ingredients in
Vicks® DayQuil® Cold & Flu LiquiCaps®





Lu


Ex

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

Distributed by: **PL Developments**
200 Hicks Street, Westbury, NY 11590

PLD-B40R FC005188

Actual Size





3 59726 46825 1

WELLNESS BASICS DayTime Multi-Symptom Cold & Flu Relief

DAYTIME COLD FLU MULTI SYMPTOM

acetaminophen, dextromethorphan hbr, phenylephrine hci capsule, liquid filled

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:59726-842
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)			PHENYLEPHRINE HYDROCHLORIDE	5 mg
Inactive Ingredients				
Ingredient Name				Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)				
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GELATIN (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SORBITAN (UNII: 6O92ICV9RU)				
SORBITOL (UNII: 506T60A25R)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	orange	Score	no score	
Shape	CAPSULE	Size	20mm	

Flavor			Imprint Code		P19;95A;512;AP016	
Contains						
Packaging						
#	Item Code	Package Description		Marketing Start Date	Marketing End Date	
1	NDC:59726-842-24	24 in 1 CARTON		05/31/2018	05/31/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				
2	NDC:59726-842-48	48 in 1 CARTON		05/31/2018	05/31/2024	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				
Marketing Information						
Marketing Category		Application Number or Monograph Citation		Marketing Start Date	Marketing End Date	
OTC monograph final		part341		05/31/2018	05/31/2024	

Labeler - P & L Development, LLC (800014821)

Revised: 9/2022

P & L Development, LLC