IBUPROFEN 200MG- ibuprofen 200mg tablet, coated Athlete's Needs, Inc.

Ibuprofen 200mg

Active ingredient (in each tablet)

Ibuprofen USP, 200mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding.

The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

· take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20-25 °C (68-77° F)
- read all warnings and directions before use
- do not use if seal under bottle cap is broken or missing

Inactive ingredients

colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

Call 1-888-952-0050

Monday through Friday 9AM - 5PM EST

Principal Display Panel

0800-536-886- FIBS) T23 MGa - MA6 yabiri dguorit ya brooM Questions or Comments?

RD&Cyellow #e∑hyprome to es, polydedro es, polyethylene glycch, povidone, pregletafinized sterch, purfilied wetter, so dum sterch dyvodets, streen is sich titten fund diodde Inactive ingredients coloidal silicon do xide, co n starch

(A 97.48) 2° 3° das mesuela de pote **a modistrandrai hartido** Il seu bin ob **a** esuelore berorian directions de los de la bin in seu bin objective de la bin de la bin lesse.

m and is and children 12 years and ower; sket 1 tabele way 4

Feet and profess of the run of a series of the run of a series of series makes to series of the run of a series of series o

Directions

do not takem one than directed in the small est effective doos should be used

SNRS,

Description of the second control of the special of the spe

Coccesses and selve a chacket in a you experience any of the blowing signs of spins of bleeding: a feel faint a womith blood blowing signs of spins of bleeding: a feel faint a womith of the office of the properties of the proper

When using this product take with tood or milk if stornach upset

Ask a dodor or pharmadat before use if you are in urder a actor or arto Aspassous or and aspassous actific berefit of actor or arto Aspassous or actification in the art actor or arto Aspassous or actification or actificati

 you have a history of stoms choroble ms, such as heartburn
 you have high boo do ressure, heart class as, liver circh cais,
 iddery disease, ashma, or had a stocke myou are taking a duetic srectibes rever in early and a single of the street reducers

Drug Facts (c ontinued)



Drug Facts

Active ingredient (in each tablet)

.....Pain reliever/fever reducer

lbuprofen USP, 200mg (NSAID)*...... *nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:

| headache | toothache | backache | menstrual cramps | the
| common cold | museular aches | minor pain of arthritis |
| temporarily achies | family |

Warnings
Allergy aler's tourrofen may cause a severe allergic reaction,
especially in people allergic to aspirin. Symptoms may include: withes
a facial swelling within (wheezing) wishock wiskin reddening
a rash wishsers
If an allergic reaction occurs, stop use and seek medical help right

Stomach bleeding warning: This product contains a nonsteroidal arti-inflammatory drug (IKSAID), which may cause severe stomach bleeding. The chance is higher if you mare age 80 or older marker had stomach ulcers or bleeding problems marke a blood blinning (antiboagularit) or steroid drug = take other drugs containing prescription or nonprescription NSAIDs (asprin, buproten, naproxen, or others) = have 3 or more alcoholic drinks every day white using this product = take more or for a longer time than directed

Heart attack and stroke warning: NSADs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer ≡ right before or after heart surgery

Ask a doctor before use if stomach bleeding warning applies to you you have problems or Drug Facts (continued on inside)

**This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advit®

Exp. Date:

Manufactured for: A.N.I. Renaissance PI

Toledo, OH 43623 I B1469

模



Clear Choice (logo)

IBUPROFEN

MODERN PAIN MEDICINE

PAIN & FEVER RELIEF

CONTAINS NO ASPIRIN

CAFFEINE FREE

Compare to Advil®**

SEE NEW WARNINGS INFORMATION

100 Tablets

200 mg Each

IBUPROFEN 200MG

ibuprofen 200mg tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53185-381
---------------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
POLYDEXTROSE (UNII: VH2XOU12IE)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POVIDONE (UNII: FZ 989GH94E)			
WATER (UNII: 059QF0KO0R)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics			
Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G2
Contains			

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:53185-381- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2016		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202312	10/07/2016	

Labeler - Athlete's Needs, Inc. (619703945)

Revised: 11/2017 Athlete's Needs, Inc.