LIQUID GLYCERIN LAXATIVES- glycerin liquid
Shanghai Yijing Medical Technology Co., Ltd.

Liquid Glycerin Laxatives

Liquid Glycerin Laxatives

GLYCERIN 5.3g

Hyperosmotic laxative

USe for relief of occasional constipation. This product usually produces bowel movement in 15 minutes to 1 hour.

May cause rectal discomfort or burning sensation.
For rectal use only

Do not use when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before using any laxative if you
have a sudden change in bowel habits lasting more than 2 weeks
already used a laxative for more than 1 week

Stop using this product and consult a doctor if you have
rectal bleeding
no bowel movement within 1 hour of using this product
These symptoms may indicate a serious condition

If swallowed, get medical help or contact a Poison Control Center right away.

Single Daily Dosage (per 24 hours)
Adults and children 12 years and over: 1 suppository or as directed by a doctore

Left-side position: Lie on left side with right knee bent and arms resting comfortably. Knee-chest position: Kneel, then lower head and chest forward until leftside of face is resting on surface with left arm folded comfortably. Administering the laxative: CAUTION: REMOVE ORANGE PROTECTIVESHIELD BEFORE INSERTING. Hold unit upright, grasping bulb of unit withfingers. Grasp orange protective shield with other hand, pull gently to remove.With steady pressure, insert tip into rectum with a slight side-to-sidemovement, with tip pointing toward navel. Do not force the applicator tip intothe rectum, as this can cause injury. Insert ion may be easier if person receivingsuppository bears down as if having a bowel movement. This helps relax themuscles around the anus. Squeeze the bulb until nearly all liquid is expelled.While continuing to squeeze the bulb, remove tip from rectum and discard unit.
it is not necessary to empty unit completely.

WATER



LIQUID GLYCERIN LAXATIVES

glycerin liquid

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:85157-002
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	7.2 g in 10 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85157-002-01	30 in 1 CARTON	02/11/2025	
1		10 mL in 1 APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/11/2025	

Labeler - Shanghai Yijing Medical Technology Co., Ltd. (567720668)

Establishment

Name	Address	ID/FEI	Business Operations
Shanghai Yijing Medical Technology Co., Ltd.		567720668	manufacture(85157-002)

Revised: 2/2025 Shanghai Yijing Medical Technology Co., Ltd.