

**TYLENOL COLD PLUS FLU SEVERE DAY/NIGHT- acetaminophen,  
chlorpheniramine maleate, dextromethorphan hydrobromide, guaifenesin,  
and phenylephrine hydrochloride  
Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**TYLENOL COLD PLUS FLU SEVERE DAY/NIGHT**

**TYLENOL<sup>®</sup> COLD + FLU SEVERE NIGHT**

**Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- for the temporary relief of the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
  - sinus congestion and pressure
  - sneezing and runny nose
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks

- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>▪ take 2 caplets every 4 hours</li> <li>▪ swallow whole; do not crush, chew or dissolve</li> <li>▪ do not take more than 10 caplets in 24 hours</li> </ul>
children under 12 years	ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

**Inactive ingredients**

anhydrous citric acid, carnauba wax, FD&C blue no. 1 aluminum lake, flavor, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polysorbate 80, potassium sorbate, powdered cellulose, pregelatinized starch, sodium benzoate, sodium citrate dihydrate, sodium starch glycolate, sucralose, titanium dioxide

**Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

## **TYLENOL® COLD + FLU SEVERE DAY**

### **Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

### **Uses**

- for the temporary relief of the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

### **Warnings**

#### **Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not exceed recommended dose**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 4 hours</li><li>▪ swallow whole; do not crush, chew or dissolve</li><li>▪ do not take more than 10 caplets in 24 hours</li></ul>
children under 12 years	ask a doctor

### **Other information**

- each caplet contains: **sodium 3 mg**
- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

### **Inactive ingredients**

carnauba wax, croscarmellose sodium, D&C yellow no. 10 aluminum lake, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sucralose, titanium dioxide

### **Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

### **PRINCIPAL DISPLAY PANEL**

NDC 50580-414-04

TYLENOL®  
FOR ADULTS

COLD + FLU  
SEVERE

Acetaminophen,  
Dextromethorphan HBr,  
Phenylephrine HCl, Guaifenesin  
Pain Reliever-Fever Reducer, Cough Suppressant,  
Nasal Decongestant, Expectorant

DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION

Actual  
Size

16 CAPLETS

Acetaminophen,  
Chlorpheniramine Maleate,  
Dextromethorphan HBr, Phenylephrine HCl  
Pain Reliever-Fever Reducer, Antihistamine,  
Cough Suppressant, Nasal Decongestant


NIGHT

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE

Actual  
Size

8 CAPLETS

TOTAL 24 CAPLETS



123665

NDC 50580-414-04

# TYLENOL®

FOR ADULTS

## COLD + FLU SEVERE

### Acetaminophen,

Dextromethorphan HBr,  
Phenylephrine HCl, Guaifenesin  
Pain Reliever–Fever Reducer, Cough Suppressant,  
Nasal Decongestant, Expectorant

### Acetaminophen,

Chlorpheniramine Maleate,  
Dextromethorphan HBr, Phenylephrine HCl  
Pain Reliever–Fever Reducer, Antihistamine,  
Cough Suppressant, Nasal Decongestant

TYLENOL®  
FOR ADULTS  
COLD + FLU  
SEVERE

DO NOT USE IF BLISTER  
UNIT IS TORN OR BROKEN

#### DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION



Actual Size

16 CAPLETS

#### NIGHT

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



Actual Size

8 CAPLETS


TOTAL 24 CAPLETS

### PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

30047050/  
123665



123665

How can we help?  
1-877-895-3665



#### Drug Facts (continued)

Save throat warning: if sore throat is severe or follows by fever, headache, rash, nausea

Do not use

- with any other drug containing acetaminophen
- if you are not sure whether a drug contains acetaminophen
- if you are now taking a prescription or over-the-counter drug for depression, psychiatric or emotional problems
- if you are taking the MAOI drug, it contains an MAOI, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen
- if you have liver disease
- if you have kidney disease
- if you have diabetes
- if you have high blood pressure
- if you are taking a blood thinning drug or warfarin

Ask a doctor or pharmacist before use if you have

- liver disease
- heart disease
- diabetes
- trouble urinating
- persistent or chronic cough such as occurs with asthma
- a breathing problem such as emphysema

When using this product

- do not exceed recommended dose
- eczema may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle

#### TYLENOL® COLD + FLU

#### Drug Facts (continued)

- Stop use and ask a doctor if you have:
  - nervousness, dizziness, or sleeplessness
  - pain, nasal congestion or coughing that worsens
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - cough comes on suddenly

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care provider about use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a poison control center. (1-800-272-1222) Ask a pharmacist for information about safe use for children even if you do not notice any symptoms.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.

Directions: See back of box for complete directions.



ing persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

other (prescription or nonprescription). If you are taking phenylephrine, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

High blood pressure ■ thyroid disease  
le to an enlarged prostate gland  
urs with smoking, asthma or empty sella  
1 (mucus) ■ glaucoma  
arbitronic bronchitis  
if you are  
■ taking sedatives or tranquilizers

hen  
increase drowsiness  
or operating machinery

**Drug Facts (continued)**

See the red warning: If severe illness severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are unsure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have  
■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease  
■ diabetes ■ trouble urinating due to an enlarged prostate gland  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema  
■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache at least  
These could be signs of a serious condition.

**FLU SEVERE NIGHT**

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

**TYLENOL® COLD + FLU SEVERE DAY**

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

1. medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.



Tylenol® Cold + Flu Severe Day Made in Italy  
Distributed by: JOHNSON & JOHNSON CONSUMER INC.  
McNeil Consumer Healthcare Division  
Fort Worth, TX, PA 19034 USA  
©J&JCI 2020

**Important: Read all product information before using. Keep this box for important information.**  
Do not take the day and night caplets at the same time. Do not take more than a total of 10 caplets in a 24-hour period. Take only as directed.

**TYLENOL® COLD + FLU SEVERE NIGHT**

**Drug Facts**

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** ■ for the temporary relief of the following cold/flu symptoms:  
 ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion  
 ■ cough ■ sinus congestion and pressure ■ sneezing and runny nose  
 ■ helps clear nasal passages ■ relieves cough to help you sleep  
 ■ temporarily reduces fever

**Warnings**  
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**TYLENOL® COLD + FLU SEVERE DAY**

**Drug Facts**

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** ■ for the temporary relief of the following cold/flu symptoms:  
 ■ minor aches and pains ■ headache ■ sore throat  
 ■ nasal congestion ■ cough  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive ■ temporarily reduces fever

**Warnings**  
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

LIFT HERE

**TYLENOL COLD PLUS FLU SEVERE DAY/NIGHT**  
 acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride kit

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-414
---------------------	----------------	---------------------------	---------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-414-03	3 in 1 CELLO PACK	07/18/2016	
1	NDC:50580-414-01	1 in 1 CARTON		
2	NDC:50580-414-04	1 in 1 CARTON	12/21/2018	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	8
Part 2	2 BLISTER PACK	16

## Part 1 of 2

### TYLENOL COLD PLUS FLU SEVERE NIGHT

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, film coated

## Product Information

<b>Route of Administration</b>	ORAL
--------------------------------	------

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)

### Product Characteristics

<b>Color</b>	blue (Light blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	TYLENOL;1075
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

## Part 2 of 2

### TYLENOL COLD PLUS FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

### Product Information

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
---	-----------------------------	------

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	TYLENOL;SEVERE
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/01/2011	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/18/2016	

**Labeler** - Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division (878046358)