

**BACTIMICINA COUGH AND COLD- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution**

**DLC Laboratories, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**BACTIMICINA® COUGH AND COLD**

***Drug Facts***

<b><i>Active ingredients (in each 5 mL = 1 teaspoon)</i></b>	<b><i>Purposes</i></b>
Dextromethorphan HBr, USP 10 mg	Cough Suppressant
Guaifenesin, USP 100 mg	Expectorant
Phenylephrine HCl, USP 5 mg	Nasal Decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are** taking any other oral nasal decongestant or stimulant.

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- this adult product is not intended for children under 12 years old

<b>Age (years)</b>	<b>Dose</b>
12 and over	2 teaspoons (10 mL) every 4 hours
Under 12	do not use

**Other information**

- **each teaspoon (5 mL) contains:** sodium 2 mg
- store at 15-30°C (59-86°F)
- dosage cup provided

**Inactive ingredients**

citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

**Questions**

1-800-858-3889

Manufactured by  
DLC Laboratories, Inc.  
Paramount, CA 90723 USA

**PRINCIPAL DISPLAY PANEL - 118 mL Bottle Box**

**LIQUID**

**BACTIMICINA®**

**Dextromethorphan HBr / Cough Suppressant**

**Guaifenesin / Expectorant**

**Phenylephrine HCl / Nasal Decongestant**

**Cough  
& Cold**

**Relieves:**

- *COUGHS*
- *STUFFY NOSE*
- *CHEST CONGESTION*

For Ages 12 & Over

**Non-Drowsy Formula**

**Alcohol Free**

**Strawberry Flavor**

**4 FL OZ (118 mL)**

Do not use if printed bottle wrap is missing or broken.  
No usar si el sello impreso alrededor de la botella  
falta o está roto.

**B**ACTIMICINA<sup>®</sup> LIQUID

**Cough  
& Cold**



**Drug Facts**

Active ingredients (in each 5 mL = 1 teaspoon)	Purposes
Dextromethorphan HBr, USP 10 mg	Cough Suppressant
Guaifenesin, USP 100 mg	Expectorant
Phenylephrine HCl, USP 5 mg	Nasal Decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

- Do not use**
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- this adult product is not intended for children under 12 years old

**B**ACTIMICINA<sup>®</sup> LIQUID

Dextromethorphan HBr / Cough Suppressant  
Guaifenesin / Expectorant  
Phenylephrine HCl / Nasal Decongestant

**Cough  
& Cold**

**Relieves:**

- COUGHS
- STUFFY NOSE
- CHEST CONGESTION

For Ages 12 & Over  
Non-Drowsy Formula  
Alcohol Free



4 FL OZ (118 mL)

**Drug Facts** (continued)

Age (years)	Dose
12 and over	2 teaspoons (10 mL) every 4 hours
Under 12	do not use

**Other information**

- each teaspoon (5 mL) contains: sodium 2 mg
- store at 15-30°C (59-86°F)
- dosage cup provided

**Inactive ingredients**

citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

**Datos Medicinales**

Ingredientes activos (en cada 5 mL = 1 cucharadita)	Propósitos
Dextrometorfano, USP 10 mg	Antitusivo
Guaifenesina, USP 100 mg	Expectorante
Fenilefrina HCl, USP 5 mg	Descongestionante Nasal

**Usos**

- ayuda a soltar flema (mucoosidad) y a diluir las secreciones bronquiales para drenarlas de los bronquios
- alivia temporalmente estos síntomas ocurridos por el resfriado:
  - congestión nasal
  - tos debido a irritaciones menores de los bronquios y la garganta

**Advertencias**

**No usar**

- si está tomando un medicamento recetado inhibidor de la monoaminoxidasa (MAOI) (ciertos medicamentos para la depresión, afecciones psiquiátricas o emocionales, o enfermedad de Parkinson), o en un periodo de 2 semanas después de haber suspendido el uso del medicamento MAOI. Si no sabe si su receta contiene MAOI, pónguele a un médico o farmacéutico antes de usar este producto.

Manufactured by / Fabricado por:  
DLC Laboratories, Inc.  
Paramount, CA 90723 USA  
Questions / Preguntas: 1-800-858-3889  
[www.dlclabs.com](http://www.dlclabs.com)

© 2014 DLC Laboratories, Inc.

Peel Here Levante Aquí

### Datos Medicinales (continuación)

Pregúntele a un médico antes de usar si usted tiene

- alguna enfermedad cardíaca
- presión arterial alta
- enfermedad de la tiroides
- diabetes
- problemas al orinar debido al agrandamiento de la glándula prostática
- tos que ocurre con demasiada flema (mucosidad)
- tos persistente o crónica como ocurre por fumar, asma, bronquitis crónica o enfisema

Consulte con un médico o farmacéutico antes de usar si usted está tomando otro descongestionante nasal o estimulante.

Cuando use este producto no exceda la dosis recomendada

Deje de usar y consulte con un médico si

- se pone nervioso, se mareo o no puede dormir
- los síntomas no mejoran en 7 días o son acompañados por fiebre
- la tos dura más de 7 días, regresa o es acompañada por fiebre, sarpullido o dolor de cabeza persistente. Estos podrían ser síntomas de una afección grave.

Si está embarazada o lactando, pregúntele a un profesional de la salud antes de usar. Manténgase fuera del alcance de los niños. En caso de sobredosis, pida ayuda médica o contáctese con un Centro de Control de Envenenamientos inmediatamente.

### Instrucciones

- no tomar más de 6 dosis en un periodo de 24 horas
- este producto para adultos no es para niños menores de 12 años de edad

Edad	Dosis
12 años y mayores	2 cucharaditas (10 mL) cada 4 horas
Menor de 12	no tomar

### Otra información

- cada cucharadita (5 mL) contiene: 2 mg de sodio
- consérvese a 15-30°C (59-86°F)
- incluye vaso dosificador

### Ingredientes inactivos

ácido cítrico, FD&C rojo no. 40, saborizante, glicerina, glicol de propileno, agua purificada, sacarina de sodio, benzoato de sodio, sorbitol

# BACTIMICINA<sup>®</sup>

JARABE

Dextrometorfano HBr / Antitusivo  
Guafenesina / Expectorante  
Fenil efrina HCl / Descongestionante Nasal

## Tos y Resfrío

### Alivia:

- TOS
- CONGESTIÓN NASAL
- CONGESTIÓN DE PECHO

Para 12 años y mayores  
No Causa Somnolencia  
Sin Alcohol

Sabor a Fresa



4 FL OZ (118 mL)



LOT

EXP





## BACTIMICINA COUGH AND COLD

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:24286-1546
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	STRAWBERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24286-1546-4	1 in 1 BOX	11/06/2013	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	11/06/2013	

**Labeler** - DLC Laboratories, Inc. (093351930)

## Establishment

Name	Address	ID/FEI	Business Operations
DLC Laboratories, Inc.		093351930	MANUFACTURE(24286-1546) , LABEL(24286-1546)

Revised: 11/2018

DLC Laboratories, Inc.