

UP AND UP ANTACID ANTI GAS- aluminum hydroxide, magnesium hydroxide, simethicone suspension

Target Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Target Corporation Antacid Anti-Gas Drug Facts

Active ingredients (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- acid indigestion
- sour stomach
- upset stomach due to these symptoms
- pressure and bloating commonly referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- measure with dosing cup provided
- adults and children 12 years and over: 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
- do not take more than 60 mL in any 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor
- mL = milliliter

Other information

- each 10 mL contains: magnesium 350 mg, potassium 20 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F), do not freeze

Inactive ingredients

butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution

Questions?

Call 1-888-547-7400

Principal Display Panel

Compare to active ingredients in Mylanta® Maximum Strength

antacid

anti-gas

maximum strength

fast acting

soothing relief of heartburn, acid indigestion, sour stomach

12 FL OZ (355 mL)

NDC 11673-637-40

Compare to active ingredients in
Mylanta® Maximum Strength*

antacid anti-gas

maximum strength
fast acting

soothing relief of heartburn,
acid indigestion, sour stomach



12 FL OZ (355 mL)

: 34040 UW F1

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Aluminum hydroxide (equiv. to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg.....	Antigas

Uses relieves ■ heartburn ■ acid indigestion ■ sour stomach
■ upset stomach due to these symptoms
■ pressure and bloating commonly referred to as gas

Warnings
Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ shake well before use ■ measure with dosing cup provided
■ **adults and children 12 years and over:** 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
■ do not take more than 60 mL in any 24 hour period
■ do not use the maximum dosage for more than 2 weeks
■ **children under 12 years:** ask a doctor ■ mL = milliliter

Other information
■ each 10 mL contains: magnesium 350 mg, potassium 20 mg and sodium 3 mg
■ **does not meet USP requirements for preservative effectiveness**
■ store at 20-25°C (68-77°F), do not freeze

Inactive ingredients butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution ▶

Drug Facts (continued)

**Questions? Call
1-888-547-7400**

245 05 0322
R02 ID285492
Dist. by Target Corp.
Mpls., MN 55403
©2016 Target Brands, Inc.
Shop Target.com

: 34040 UW B1

**DO NOT USE
IF PRINTED
NECKBAND
IS BROKEN
OR MISSING**

*This product is not
manufactured or
distributed by Infirst
Healthcare Inc.,
distributor of Mylanta®
Maximum Strength.



3 70030 62909 1

UP AND UP ANTACID ANTI GAS

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-637
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BUTYL PARABEN (UNII: 3QP1IU3FV8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-637-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/31/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	08/31/2017	

