SEVERE CONGESTION AND COUGH RELIEF- dextromethorphan hbr, guaifenesin, phenylephrine hcl solution L.N.K. International, Inc.

Sound Body 44-004

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg Guaifenesin 400 mg Phenylephrine HCl 10 mg

Purpose

Cough suppressant Expectorant Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

Warnings

Do not use

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 10 mg
- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments?

1-800-426-9391

Principal Display Panel

SOUND**BODY™**

Compare to the active ingredients in Mucinex[®] FAST-MAX[™] Severe Congestion & Cough NDC 50844-400-45

MAXIMUM STRENGTH Severe Congestion and Cough Relief

Dextromethorphan HBr Guaifenesin Phenylephrine HCl

Cough Suppressant Expectorant Nasal Decongestant

Berry Flavor

Controls Cough Relieves Nasal & Chest Congestion Thins & Loosens Mucus

6 FL OZ (177 mL)

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex® FAST-MAX™ Severe Congestion & Cough.

50844 ORG012100445

PARENTS:

Learn about teen medicine abuse www.StopMedicineAbuse.org

Manufactured for Big Lots Stores, Inc. by **LNK INTERNATIONAL, INC** 60 Arkay Drive, Hauppauge, NY 11788 USA V#733000 ITEM#022700445





- bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves: cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing ■ the impulse to cough to help you get to sleep
- nasal congestion due to a cold

Warnings Do not use

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions,

Drug Facts (continued)

or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
 diabetes
- high blood pressure thyroid disease
- difficulty in urination due to enlargement of the prostate gland persistent or chronic cough such as
- occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage

- Stop use and ask a doctor if
- nervousness, dizziness, or sleeplessness occur symptoms do not get better
- within 7 days or occur with fever cough persists more than 7 days. tends to recur, or is accompanied by a fever, rash, or persistent

headache. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided ■ adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use


Other information

- each 20 mL contains: sodium 10
- use by expiration date on package store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments? 1-800-426-9391



Sound Body 44-004

SEVERE CONGESTION AND COUGH RELIEF

dextromethorphan hbr, quaifenesin, phenylephrine hcl solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-400
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)			
SODIUM METABISULFITE (UNII: 4VON5FNS3C)			
SORBITOL (UNII: 506T60A25R)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Product Characteristics			
Color	blue	Score	
Shape		Size	
Flavor	BERRY (MIXED)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:50844- 400-45	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/15/2021	01/09/2025

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/15/2021	01/09/2025

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(50844-400) , pack(50844-400)

Revised: 6/2024 L.N.K. International, Inc.