SMOOTH LAX- polyethylene glycol 3350 powder, for solution Walgreen Company

Walgreen Co. Smooth Lax[®] Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- use once a day
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-719-9260

Principal Display Panel

VALUE SIZE

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredient in MiraLAX®

SmoothLAX®

POLYETHYLENE GLYCOL 3350

POWDER FOR SOLUTION, OSMOTIC LAXATIVE

Unflavored Powder

Grit Free

- Relieves occasional constipation (irregularity)
- Softens stool

45 ONCE-DAILY DOSES

NET WT 26.9 OZ (765 g)

| VALUE SIZE | | | | |
|--|---|-----------------|--|--|
| Topalgreens Compare to the active ingredient in MiraLAX ^{s#} | N | DC 0363-0306-09 | | |
| SmoothLAX® POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION, OSMOTIC LAXATIVE Unflavored Powder Grit Free | | | | |
| • Relieves occasional constipation (irregularity) • Softens stool • Softens stool | | | | |
| Drug Facts | Drug Facts (continued) | | | |
| Active ingredient (in each do.se) Purpose Polyethylene Glycol 3350, 17 g (cap filled to line)Osmotic Lax ative Use Intervention (irregularity) generally produces a bowel movement in 1 to 3 days Marrings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have Inausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks | adults and children 17 years of age and older: use once a day fill to top of white section in cap which is marked to inc dose (17 g) stir and dissolve in any 4 to 8 ounces of beverage (cold temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor | | | |
| ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. | Other information store at 20°-25°C (68°-77°F) tamper-evident: do not use if printed foil seal under cap is missing, open or broken Inactive ingredients none Output ingredients 24, ppp. 740, ppp. | 378294 W1 | | |
| you get diarrhea you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). | Questions or comments?1-800-719-9260 ¹ Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. ¹¹ This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX [®] . DISTRIBUTED BY: WALGREEN CO. | 0107-0822-F | | |
| Directions do not take more than directed unless advised by your doctor the battle tas is a mercuring are marked to contain 17 arems of | 200 WILMOT RD., DEERFIELD, IL 60015 100% SATISFACTION GUARANITEED walgreens.com ©2021 Walgreen Co. | | | |



SMOOTH LAX polyethylene glycol 3350 powder, for solution **Product Information Product Type** HUMAN OTC DRUG NDC:0363-0306 Item Code (Source) **Route of Administration** ORAL **Active Ingredient/Active Moiety Ingredient Name Basis of Strength Strength** POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL POLYETHYLENE 17 g 3350 - UNII:G2M7P15E5P) in 17 g GLYCOL 3350 **Product Characteristics** Color WHITE Score Shape Size Flavor Imprint Code Contains Packaging **Marketing End Marketing Start** # **Item Code** Package Description Date Date NDC:0363-0306- 238 g in 1 BOTTLE; Type 0: Not a Combination 1 10/09/2009 02 Product NDC:0363-0306- 510 g in 1 BOTTLE; Type 0: Not a Combination 2 10/09/2009 03 Product NDC:0363-0306- 850 g in 1 BOTTLE; Type 0: Not a Combination 09/28/2016 3 03/11/2011 04 Product NDC:0363-0306- 765 g in 1 BOTTLE; Type 0: Not a Combination 4 04/10/2015 09 Product NDC:0363-0306- 119 g in 1 BOTTLE; Type 0: Not a Combination 5 04/10/2015 Product 01 NDC:0363-0306- 612 g in 1 BOTTLE; Type 0: Not a Combination 04/16/2015 09/30/2021 6 10 Product NDC:0363-0306- 10 in 1 CARTON 7 04/27/2010 52 17 g in 1 POUCH; Type 0: Not a Combination 7 Product NDC:0363-0306-8 20 in 1 CARTON 05/28/2019 60 17 g in 1 POUCH; Type 0: Not a Combination 8 Product

04/27/2010

NDC:0363-0306- 17 g in 1 POUCH; Type 0: Not a Combination

Product

9

00

| Marketing Information | | | | |
|-----------------------|---|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| ANDA | ANDA090685 | 10/09/2009 | | |
| | | | | |

Labeler - Walgreen Company (008965063)

Revised: 5/2023

Walgreen Company