RUGBY CHILDRENS ACETAMINOPHEN- acetaminophen suspension Rugby Laboratories

Rugby Laboratories Children's Acetaminophen Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

	· -	
72-95	11 years	15 mL

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

Rugby®

Compare to the active ingredient in Children's Tylenol® Oral Suspension

Children's

Acetaminophen 160 mg per 5 mL

Oral Suspension

Pain Reliever/Fever Reducer

Fast, effective relief of children's pain and fever

Ibuprofen Free

Alcohol Free

Aspirin Free

Grape Flavor

For Ages 2 to 11 Years

4 FL OZ (118 mL)





NDC 0536-1321-97

Children's

Acetaminophen 160 mg per 5 mL **Oral Suspension**

Pain Reliever/Fever Reducer

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Acetaminophen 160 mg.........Pain reliever/fever reduce

- Uses temporarily:

 reduces fever

 releves minor aches and pains due to:

 the common cold

 fit to headache

 sore throat

 toothache

Warnings

ng: This product contains acetaminophen. Liver warning: This product contains acetaminopher Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maxin

with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin

reactions. Symptoms may include:

It skin reddening I blisters III rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists

for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- III with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver diseas Asik a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- in gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- mew symptoms occur
- redness or swelling is present

 These could be signs of a serious condition.

Keep out of reach of children.

Neep out or reach or children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Genter right away (1–800-222-1222), Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

- this product does not contain directions or
- complete warnings for adult use

 do not give more than directed (see overdose
- shake well before using

 If ml. = milliliter
- find right dose on chart below. If possible, use weight
- Into high course of nath below, in polisities, use well to does; otherwise, use age.

 remove the child protective cap and squeeze your child's does into the desing cup

 repeat does every 4 hours while symptoms last

 do not give more than 5 times in 24 hours.

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or
- missing

Inactive ingredients

mactive ingretierits any district and anylydrus dirtic acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructibe com syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose solium, proylene glycci, purified water, sod ium benzoate, sorbital solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

Distributed by: RUGBY®LABORATORIES Indianapolis, IN 46268 (800) 616-2471 aiorpharmaceuticals.com

Rugby

NDC 0536-1321-97 Compare to the active ingredient in Children's Tylenol* Oral Suspension**

Children's

Acetaminophen 160 mg per 5 mL Oral Suspension

Pain Reliever/Fever Reducer

Fast, effective relief of children's pain and fever

Ibuprofen Free Alcohol Free Aspirin Free



Grape Flavor For Ages 2 to 11 Years

4 FL OZ (118 mL)



Children's

Acetaminophen 160 mg per 5 mL

Oral Suspension

Pain Reliever/Fever Reducer

DIRECTIONS: STEP 1. Shake well before using



STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose dosing cup



IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.

Keep carton, it contains important information.





13026 ST C2

RUGBY CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0536-1321

Route of Administration

ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)			
CARRAGEENAN (UNII: 5C69YCD2YJ)			
D&C RED NO. 33 (UNII: 9DBA0SBB0L)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
GLYCERIN (UNII: PDC6A3C0OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)			
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)			

Product Characteristics			
Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1321- 97	1 in 1 CARTON	10/21/2020	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing In	larketing Information		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/21/2020	

Revised: 5/2024 Rugby Laboratories