# ANTIVERT- meclizine hydrochloride tablet Roerig

-----

**ANTIVERT®** 

Tablets/12.5 mg meclizine HCl

**ANTIVERT®/25** 

Tablets/25 mg meclizine HCl

**ANTIVERT®/50** 

Tablets/50 mg meclizine HCl

#### DESCRIPTION

Chemically, ANTIVERT<sup>®</sup> (meclizine HCl) is 1-(p-chloro- $\alpha$ -phenylbenzyl)-4-(m-methylbenzyl) piperazine dihydrochloride monohydrate.

Inert ingredients for the tablets are: dibasic calcium phosphate; magnesium stearate; polyethylene glycol; starch; sucrose. The 12.5 mg tablets also contain: Blue 1. The 25 mg tablets also contain: Yellow 6 Lake; Yellow 10 Lake. The 50 mg tablets also contain: Blue 1 Lake; Yellow 10 Lake.

#### **CLINICAL PHARMACOLOGY**

ANTIVERT is an antihistamine that shows marked protective activity against nebulized histamine and lethal doses of intravenously injected histamine in guinea pigs. It has a marked effect in blocking the vasodepressor response to histamine, but only a slight blocking action against acetylcholine. Its activity is relatively weak in inhibiting the spasmogenic action of histamine on isolated guinea pig ileum.

#### **Pharmacokinetics**

The available pharmacokinetic information for meclizine following oral administration has been summarized from published literature.

#### **Absorption**

Meclizine is absorbed after oral administration with maximum plasma concentrations reaching at a median Tmax value of 3 hours post-dose (range: 1.5 to 6 hours) for the tablet dosage form.

#### **Distribution**

Drug distribution characteristics for meclizine in humans are unknown.

#### **Metabolis** m

The metabolic fate of meclizine in humans is unknown. In an in vitro metabolic study using human hepatic microsome and recombinant CYP enzyme, CYP2D6 was found to be the dominant enzyme for metabolism of meclizine.

The genetic polymorphism of CYP2D6 that results in extensive-, poor-, intermediate- and ultrarapid metabolizer phenotypes could contribute to large inter-individual variability in meclizine exposure.

#### Elimination

Meclizine has a plasma elimination half-life of about 5-6 hours in humans.

#### **INDICATIONS**

Based on a review of this drug by the National Academy of Sciences - National Research Council and/or other information, FDA has classified the indications as follows:

Effective: Management of nausea and vomiting, and dizziness associated with motion sickness.

Possibly Effective: Management of vertigo associated with diseases affecting the vestibular system.

Final classification of the less than effective indications requires further investigation.

#### CONTRAINDICATIONS

Meclizine HCl is contraindicated in individuals who have shown a previous hypersensitivity to it.

#### WARNINGS

Since drowsiness may, on occasion, occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking this drug.

Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma, or enlargement of the prostate gland.

#### **Usage in Children**

Clinical studies establishing safety and effectiveness in children have not been done; therefore, usage is not recommended in children under 12 years of age.

#### Usage in Pregnancy

Pregnancy Category B. Reproduction studies in rats have shown cleft palates at 25–50 times the human dose. Epidemiological studies in pregnant women, however, do not indicate that meclizine increases the risk of abnormalities when administered during pregnancy. Despite the animal findings, it would appear that the possibility of fetal harm is remote. Nevertheless, meclizine, or any other medication, should be used during pregnancy only if clearly necessary.

#### **PRECAUTIONS**

#### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when meclizine is administered to a nursing woman.

## **Hepatic Impairment**

The effect of hepatic impairment on the pharmacokinetics of meclizine has not been evaluated. As

meclizine undergoes metabolism, hepatic impairment may result in increased systemic exposure of the drug. Treatment with meclizine should be administered with caution in patients with hepatic impairment.

### **Renal Impairment**

The effect of renal impairment on the pharmacokinetics of meclizine has not been evaluated. Due to a potential for drug/metabolite accumulation, meclizine should be administered with caution in patients with renal impairment and in the elderly as renal function generally declines with age.

## **Drug Interactions**

There may be increased CNS depression when meclizine is administered concurrently with other CNS depressants, including alcohol, tranquilizers, and sedatives. (*see* WARNINGS)

Based on in-vitro evaluation, meclizine is metabolized by CYP2D6. Therefore there is a possibility for a drug interaction between meclizine and CYP2D6 inhibitors.

#### ADVERSE REACTIONS

Anaphylactoid reaction, drowsiness, dry mouth, headache, fatigue, vomiting and, on rare occasions, blurred vision have been reported.

#### DOSAGE AND ADMINISTRATION

#### Vertigo

For the control of vertigo associated with diseases affecting the vestibular system, the recommended dose is 25 to 100 mg daily, in divided dosage, depending upon clinical response.

#### **Motion Sickness**

The initial dose of 25 to 50 mg of Antivert should be taken one hour prior to embarkation for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

#### **HOW SUPPLIED**

Antivert: 12.5 mg tablets:

Bottles of 100

(NDC 0049-2100-66)

Antivert/25: 25 mg tablets:

Bottles of 100

(NDC 0049-2110-66)

Antivert/50: 50 mg tablets:

Bottles of 100

(NDC 0049-2140-66)

#### Rx only

### PRINCIPAL DISPLAY PANEL - 12.5 mg Tablet Label

NDC 0049-2100-66

100 Tablets

Antivert®

(meclizine HCl)

12.5 mg

**Pfizer** 

Distributed by

Roerig

Division of Pfizer Inc, NY, NY 10017



## PRINCIPAL DISPLAY PANEL - 25 mg Tablet Label

NDC 0049-2110-66

100 Tablets

Antivert®

(meclizine HCl)

25 mg

Pfizer

Distributed by

Roerig

Division of Pfizer Inc, NY, NY 10017



## PRINCIPAL DISPLAY PANEL - 50 mg Tablet Label

NDC 0049-2140-66

100 Tablets

Antivert® (meclizine HCl)


50 mg


**Pfizer** 

Distributed by

Roerig

Division of Pfizer Inc, NY, NY 10017





Route of Administration

ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

MECLIZINE HYDRO CHLO RIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570) MECLIZINE HYDRO CHLO RIDE | 12.5 mg

## **Inactive Ingredients**

inactive ingredients				
Ingredient Name	Strength			
CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US (UNII: L11K75P92J)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)				
SUCROSE (UNII: C151H8M554)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				

## **Product Characteristics**

Color	BLUE, WHITE	Score	no score
Shape	OVAL	Size	10 mm
Flavor		Imprint Code	Antivert;210
Contains			

## **Packaging**

_	i uckuging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:0049-2100-66	100 in 1 BOTTLE					
2	NDC:0049-2100-82	1000 in 1 BOTTLE					

## **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA010721	04/11/1997	

## **ANTIVERT**

meclizine hydrochloride tablet

D۳	ոժո	ct I	nfo	rm	ation	
РΙ				1 11 11 17	411011	

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0049-2110
Pouts of Administration	ORAL		

Active Ingredient/Active Moiety	Active	Ingred	lient/Ac	tive M	Ioie tv
---------------------------------	--------	--------	----------	--------	---------

- 1			
	Ingredient Name	Basis of Strength	Strength
ı	MECLIZINE HYDDOCHI ODIDE (LINII: HDD7W/4/CIO) (MECLIZINE - LINII: 21 5TO 8/4570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients				
Ingredient Name	Strength			
CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US (UNII: L11K75P92J)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)				
SUCROSE (UNII: C151H8 M554)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
ALUMINUM OXIDE (UNII: LMI26O6933)				

Product Characteristics						
Color	YELLOW, WHITE	Score	no score			
Shape	OVAL	Size	13mm			
Flavor		Imprint Code	Antivert;211			
Contains						

P	Packaging							
#	Item Code	Package Description	Marketing Start Date	Marketing End Date				
1	NDC:0049-2110-66	100 in 1 BOTTLE						
2	NDC:0049-2110-82	1000 in 1 BOTTLE						

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
NDA	NDA0 10 721	04/11/1997				

## **ANTIVERT**

meclizine hydrochloride tablet

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0049-2140	
Route of Administration	ORAL			

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength		
IECT TAINE HANDOCHT ODINE (TINII: HDD7MMCIO) (MECT TAINE	HMH-21 ETO 9 4570)	MECI IZINE UVDDOCUI ODIDE	EO ma	

MECLIZINE HYDRO CHLO RIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570) | MECLIZINE HYDRO CHLO RIDE | 50 mg

Inactive Ingredients	
Ingredient Name	Strength

CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US (UNII: L11K75P92J)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)		
SUCROSE (UNII: C151H8M554)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
ALUMINUM OXIDE (UNII: LMI26O6933)		

Product Characteristics				
Color	YELLOW, BLUE	Score	2 pieces	
Shape	OVAL	Size	16 mm	
Flavor		Imprint Code	Antivert;214	
Contains				

	Packaging				
	# Item Code	Package Description	Marketing Start Date	<b>Marketing End Date</b>	
ı	1 NDC:0049-2140-66	100 in 1 BOTTLE			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA0 10 721	04/11/1997		

# **Labeler** - Roerig (829076996)

Establishment			
Name	Address	ID/FEI	Business Operations
UCB Pharma S.A.		372274485	MANUFACTURE(0049-2140, 0049-2110, 0049-2100)

Establishment				
Name	Address	ID/FEI	Business Operations	
Pfizer Pharmaceuticals LLC		829084545	API MANUFACTURE(0049-2140)	

Revised: 11/2012 Roerig