

MOTION SICKNESS RELIEF- meclizine hcl tablet
Topco Associates, LLC

Topcare 44-403

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- take first dose ½ to 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments?

1-888-423-0139

Principal display panel

+TopCare®
health

NDC 36800-943-19

**COMPARE TO
DRAMAMINE® LESS DROWSY
ACTIVE INGREDIENT***

LESS DROWSY FORMULA

Motion Sickness Relief
MECLIZINE HCl, 25 mg - ANTIEMETIC

LONG LASTING:

- Up to 24 Hours of Protection
- Prevents Nausea,
Dizziness and Vomiting

Ages 12 Years and Over

**UP TO
24
HOURS**

8 TABLETS

actual size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO LNKA0521 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Less Drowsy.

50844 ORG042340319

**QUALITY
GUARANTEED**

UP TO 24 HOURS **LONG LASTING:**
• Up to 24 Hours of Protection
• Prevents Nausea, Dizziness and Vomiting

+TopCare.
health

8 TABLETS

LESS DROWSY FORMULA

Motion Sickness Relief
MECLIZINE HCl, 25 mg - ANTIEMETIC

NDC 36800-943-19

+TopCare.
health

**COMPARE TO
DRAMAMINE® LESS DROWSY
ACTIVE INGREDIENT***

LESS DROWSY FORMULA

Motion Sickness Relief
MECLIZINE HCl, 25 mg - ANTIEMETIC

UP TO 24 HOURS **LONG LASTING:**
• Up to 24 Hours of Protection
• Prevents Nausea,
Dizziness and Vomiting

Ages 12 Years and Over

8 TABLETS

actual size

E1044

8-1910-08004

1-888-423-0139

50844 ORG042340319

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Less Drowsy.

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts (continued)

Drug Facts
(in each tablet)
Meclizine HCl 25 mg
Antiemetic

Purpose
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than directed
take first dose 1/2 to 1 hour before starting activity
adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
protect from heat and humidity
see end flap for expiration date and lot number


Inactive ingredients
colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments? 1-888-423-0139

Drug Facts
Warnings
Do not use for children under 12 years of age unless directed by a doctor.
Ask a doctor before use if you have difficulty in urination due to enlargement of the prostate gland
glaucoma
a breathing problem such as emphysema or chronic bronchitis
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
drowsiness may occur
avoid alcoholic beverages
alcohol, sedatives, and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery

Uses
for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness



TopCare 44-403

0 36800 77027 0

No print/No varnish
Lot & Exp date

Drug Facts
Warnings
Do not use for children under 12 years of age unless directed by a doctor.
Ask a doctor before use if you have difficulty in urination due to enlargement of the prostate gland
glaucoma
a breathing problem such as emphysema or chronic bronchitis
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
drowsiness may occur
avoid alcoholic beverages
alcohol, sedatives, and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery

Drug Facts
Active ingredient
(in each tablet)
Meclizine HCl 25 mg
Antiemetic

Purpose
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than directed
take first dose 1/2 to 1 hour before starting activity
adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
protect from heat and humidity
see end flap for expiration date and lot number

Inactive ingredients
colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments? 1-888-423-0139

TopCare 44-403

0 36800 77027 0

No print/No varnish
Lot & Exp date

Drug Facts
Warnings
Do not use for children under 12 years of age unless directed by a doctor.
Ask a doctor before use if you have difficulty in urination due to enlargement of the prostate gland
glaucoma
a breathing problem such as emphysema or chronic bronchitis
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
drowsiness may occur
avoid alcoholic beverages
alcohol, sedatives, and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery

Drug Facts
Active ingredient
(in each tablet)
Meclizine HCl 25 mg
Antiemetic

Purpose
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than directed
take first dose 1/2 to 1 hour before starting activity
adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
protect from heat and humidity
see end flap for expiration date and lot number

Inactive ingredients
colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments? 1-888-423-0139

TopCare 44-403

0 36800 77027 0

No print/No varnish
Lot & Exp date

MOTION SICKNESS RELIEF

meclizine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-943
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-943-19	1 in 1 CARTON	05/30/2019	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	05/30/2019	

Labeler - Topco Associates, LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-943)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-943) , pack(36800-943)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-943)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(36800-943)